



Royal Netherlands Institute for Sea Research

This is a pre-copyedited, author-produced version of an article accepted for publication, following peer review.

**Penning, E.; Govers, L.L.; Dekker, R.; Piersma, T.** (2021). Advancing presence and changes in body size of brown shrimp *Crangon crangon* on intertidal flats in the western Dutch Wadden Sea, 1984–2018. *Mar. Biol. (Berl.)* 168(11): 160.

Published version: <https://dx.doi.org/10.1007/s00227-021-03967-z>

NIOZ Repository: <http://imis.nioz.nl/imis.php?module=ref&refid=345833>

[Article begins on next page]

The NIOZ Repository gives free access to the digital collection of the work of the Royal Netherlands Institute for Sea Research. This archive is managed according to the principles of the [Open Access Movement](#), and the [Open Archive Initiative](#). Each publication should be cited to its original source - please use the reference as presented.

When using parts of, or whole publications in your own work, permission from the author(s) or copyright holder(s) is always needed.

1 manuscript for *Marine Biology* (dd. 20 August 2021):

2

3 Advancing presence and changes in body size of brown shrimp *Crangon*  
4 *crangon* on intertidal flats in the western Dutch Wadden Sea, 1984-2018

5

6 Emma Penning<sup>1,2</sup>, Laura L. Govers<sup>1,2</sup>, Rob Dekker<sup>1</sup>, Theunis Piersma<sup>1,2</sup>

7

8

9 <sup>1</sup>NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems, P.O. Box 59, 1790  
10 AB Den Burg, Texel, The Netherlands

11 <sup>2</sup>Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University  
12 of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands

13

14

15 Address for correspondence:

16 Emma Penning

17 NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems, P.O. Box 59, 1790

18 AB Den Burg, Texel, The Netherlands

19 E-mail: [emma.penning@nioz.nl](mailto:emma.penning@nioz.nl)

## 20 **Abstract**

21 Upon settlement after a pelagic larval phase, brown shrimp *Crangon crangon* depend on  
22 intertidal flats. During low as well as high tide the young brown shrimp play roles as predators of  
23 meiofauna and as prey for fish and birds. Unlike the biology of the commercially important  
24 adults, knowledge on these juveniles remains sketchy. Here we provide an analysis of 35 years  
25 (1984-2018) of brown shrimp monitoring in May-June on intertidal flats in the westernmost  
26 Dutch Wadden Sea. Intertidal shrimp densities were sampled bi-weekly at three stations during  
27 low tide, using sampling corers. We show that over this 35-year period the appearance of shrimp  
28 on mudflats advanced by 12 days ( $-0.34 \text{ days yr}^{-1}$ ). Simultaneously, densities on 7 May increased  
29 by more than 2.4 times, from 28 shrimp  $\text{m}^{-2}$  in 1984 to 69 shrimp  $\text{m}^{-2}$  in 2018. Across years,  
30 mean shrimp length decreased from 12.6 to 10.7 mm, but length in early May did not change.  
31 The advancement in settlement and the increasing shrimp densities correlated with increases in  
32 the seawater temperatures in April more than during earlier times of the year. We propose four  
33 interpretations of these changes: (1) shrimp settle on the mudflat when they reach a certain  
34 ‘threshold’ length, (2) settlement of shrimp is controlled by a critical period of ‘threshold’  
35 temperature sensitivity, (3) timing of shrimp settlement is a response to food availability on  
36 mudflats or (4) a direct response to inferred predation pressure. The different interpretations will  
37 lead to different scenarios of change in a warming world.

38  
39 **key words:** seasonal timing, life cycle, intertidal foodweb, benthos, Wadden Sea, long-term  
40 monitoring, *Crangon crangon*

41

42


## 43 **Introduction**

44 Intertidal flat systems often support high benthic primary production (Christianen et al. 2017)  
45 and a high number of primary and secondary consumers (Mathot et al. 2019). An abundance of  
46 bivalves, polychaete worms and crustaceans attracts birds and fish to these flats (Piersma et al.  
47 1993). The international Wadden Sea, the most extensive intertidal system in the world, is visited  
48 on a yearly basis by hundreds of thousand shorebirds, who aggregate here to refuel during long-  
49 distance migrations between northern breeding areas and more southerly wintering areas (van  
50 de Kam et al. 2004; Rakhimberdiev et al. 2018). Among the intertidal benthic invertebrates,

51 brown shrimp *Crangon crangon* are important prey for e.g. Eurasian spoonbills *Platalea*  
52 *leucorodia* (Jouta et al. 2018), sanderling *Calidris alba* (Penning et al. unpubl. data) and dunlin  
53 *Calidris alpina* (Nehls and Tiedemann 1993). When submerged, shrimp are prey for offshore  
54 pelagic fish like whiting *Merlangius merlangus* and cod *Gadus morhua* and in the Wadden Sea  
55 for demersal species like plaice *Pleuronectes platessa* and gobies Gobiidae (Tiews 1970; del  
56 Norte Campos and Temming 1994).

57         The Wadden Sea supports a high biomass of brown shrimp which are of ecological and  
58 commercial importance (Campos and van der Veer 2008; Tulp et al. 2016). Shrimp play a central  
59 role in the food web, acting both as key prey and key predators (Pihl and Rosenberg 1984; Oh et  
60 al. 1999). The main food of large shrimp (>20 mm) consists of juvenile plaice (van der Veer and  
61 Bergman 1987; Gibson et al. 1995) and several of the small benthic macrofaunal species (Pihl  
62 and Rosenberg 1984; Jensen and Jensen 1995; van der Veer et al. 1998); on this basis top-down  
63 control of bivalve recruitment by shrimp has been proposed (van der Veer et al. 1998; Philippart  
64 et al. 2003; Beukema and Dekker 2005, 2014; Andresen and van der Meer 2010; van der Heide  
65 et al. 2014). On the mudflats, shrimp smaller than 20 mm mainly eat meiofaunal organisms such  
66 as copepods and ostracods (Pihl and Rosenberg 1984; Boddeke et al. 1985; Jensen and Jensen  
67 1995). Cannibalism of larger on smaller size classes is not uncommon (Marchand 1981; Pihl and  
68 Rosenberg 1984), but the impact on juvenile survival and mortality rates remains unknown  
69 (Campos and van der Veer 2008).

70         The life-cycle of shrimp can be sketched as consisting of three stages (Fig. 1) in which  
71 (1) the larvae are planktonic, (2) post-larvae settle on the intertidal flats, and (3) adults live  
72 demersally in the subtidal zone (Lloyd and Yonge 1947). Females carry the eggs until they  
73 hatch, after which the larvae join the zooplankton (Smaldon 1979). The planktonic larvae go  
74 through different stages before they reach a length of 4.7 mm and settle on intertidal soft  
75 sediments supported by North Sea currents and selective tidal stream transport (Tiews 1970,  
76 Daewel et al. 2011); by now they are called ‘post-larvae’ living in ‘nursery areas’ (Tiews 1970;  
77 Beukema 1992; Campos and van der Veer 2008). Records of shrimp <20 mm in the subtidal are  
78 scarce, but appropriate sampling to catch those small sized shrimp has not been carried out  
79 (Janssen and Kuipers 1980; Boddeke et al. 1985; Beukema 1992). Therefore, it remains  
80 surprisingly unclear if post-larvae make use of the subtidal zone at all.



81  
 82 **Fig. 1** Illustration of the known and described life-cycle of shrimp (based on Tiews 1970,  
 83 Janssen and Kuipers 1980, Kuipers and Dapper 1984 and Campos 2009). Adult females carry  
 84 eggs until they hatch. The hatched eggs go through different larval stages as plankton (up to zoea  
 85 5) before they settle on intertidal mudflats as post-larvae. Here they continue to develop as  
 86 juveniles. Before shrimp become adult, they leave the intertidal and move to deeper waters of the  
 87 subtidal and offshore  
 88

89 In autumn, adults make a seasonal migration to deeper waters, returning in spring with  
 90 the incoming tides to coastal areas such as the Wadden Sea (Broekema 1941; van der Baan 1975;  
 91 Boddeke 1976; Spaargaren 2000). In addition to *seasonal* migrations, adult shrimp show *tidal*  
 92 migrations: they move to the mudflats with the incoming tide and move to deeper waters again  
 93 with the outgoing tide (Hartsuyker 1966; Al-Adhub and Naylor 1975). These movements may be  
 94 triggered by changes in hydrostatic pressure which are sensed by adult shrimp (Tielmann et al.  
 95 2015). No such movements are known for the settling post-larvae and juveniles, which appear to  
 96 remain on the intertidal flats during both low and high tide (Janssen and Kuipers 1980). Large  
 97 juveniles have been reported to gradually leave the intertidal zone when they reach lengths of 20-  
 98 25 mm, moving to the subtidal parts of the Wadden Sea and to the nearshore parts of the North  
 99 Sea (van der Baan 1975; Kuipers and Dapper 1981, 1984; Beukema 1992). However, in an  
 100 experimental setting, shrimp 15 - 20 mm showed selective ebb tide activity, indicating that  
 101 juvenile departure from intertidal mudflats may even start a bit earlier than at lengths of 20-25  
 102 mm (Hufnagl et al. 2014).

103 Male shrimp become sexually mature at smaller lengths (22–43 mm) than females (30–55  
 104 mm) (Lloyd and Yonge 1947; ICES 2015). Spawning can take place in as long a period as 46 of  
 105 the 52 weeks of the year, with egg-bearing females being observed year-round (Tiews 1970;

106 Boddeke 1982; Hünerlage et al. 2019). Nevertheless, there are peaks of egg-bearing females in  
107 winter and early summer (Boddeke and Becker 1979; Boddeke 1982; Kuipers and Dapper 1984;  
108 Siegel et al. 2008; Hünerlage et al. 2019). Egg-bearing females are most abundant in shallow  
109 offshore waters up to 20 m deep (Hünerlage et al. 2019). The average egg size gradually changes  
110 during the spawning season (Hünerlage et al. 2019), with winter eggs being larger than summer  
111 eggs (Boddeke 1982). The peak of post-larvae and juvenile shrimp on mudflats in spring mainly  
112 originates from winter spawning (Temming and Damm 2002). Nevertheless, Temming et al.  
113 (2017) argue that later in spring and early summer, shrimp from summer eggs also contribute.

114 The extent to which the intertidal flats of the Wadden Sea act as nurseries for brown  
115 shrimp has been suggested to be affected by winter seawater temperatures (Beukema 1992),  
116 sediment type (Beukema and Dekker 2005), surface water nutrient loads (Boddeke and Hagel  
117 1991; Boddeke 1996; Philippart et al. 2007), with obvious direct and indirect effects of fisheries  
118 (cf. Tulp et al. 2020). Shrimp may be affected in different ways: cold winters would delay the  
119 settlement of post-larvae (Beukema 1992), shrimp densities may be higher in coarse sediments  
120 than fine sediments (Beukema and Dekker 2005), eutrophication may correlate with an increase  
121 in secondary production (Philippart et al. 2007) boosting shrimp recruitment or growth as a result  
122 (Boddeke and Hagel 1991; Boddeke 1996). The trawling for shrimp disturbs the seafloor and  
123 make it less habitable for juvenile shrimp (Tulp et al. 2020).

124 Between 1983 and 2012, juvenile shrimp densities in spring on the mudflats increased  
125 (Beukema and Dekker 2014). At the same time, long-term trends in the timing of shrimp  
126 settlement on mudflats remain unknown. The timing of settlement may be influenced by water  
127 temperature, as higher temperatures speed up the development of shrimp from eggs to adults  
128 (Criales and Anger 1986). With the mean water temperature of the Dutch Wadden Sea having  
129 increased over the past decades (van Aken 2008), this could have advanced the timing of shrimp  
130 settlement on intertidal flats. To update the situation and re-assess some previous correlations  
131 with environmental factors, we here provide an analysis of 35 years (1984-2018) of monitoring  
132 in May-June of post-larvae and juvenile shrimp on the intertidal flats in the westernmost Dutch  
133 Wadden Sea (Balgzand). We aimed to study shrimp in terms of: (1) timing of appearance on the  
134 intertidal flats, (2) abundance, and (3) body size. A lack of clear predictions and precise enough  
135 relevant environmental data made it impossible to simultaneously assess correlations with factors  
136 such as sediment characteristics, nutrient loadings of the seawater and sediment and fishery

137 pressures. Thus, rather than go out on a ‘fishing expedition’ to survey correlates with all possible  
138 environmental covariates, we made the choice to focus on the most obvious previously  
139 established relationship, i.e. to examine correlations with seawater temperatures over different  
140 time periods preceding the occurrence of post-larval settling shrimp in May-June.

141

## 142 **Methods**

### 143 **Study area**

144 The Balgzand area is a tidal flat system of 50 km<sup>2</sup> in the western part of the Dutch Wadden Sea,  
145 at ~53°N and 5°E. The area is characterized by a semi-diurnal tide with an amplitude of 1.5-2.5  
146 m, depending on exact location, lunar phase and wind conditions (Dapper and van der Veer  
147 1981). Height of intertidal mud- and sandflats ranges from ~ 70 cm above to ~ 80 cm below  
148 mean tide level (MTL). The majority of the area is located under MTL. Details of physical  
149 parameters of Balgzand can be found in Beukema and Cadée (1997).

150 Shrimp densities were measured at three sampling stations in the southwestern part of  
151 Balgzand (Beukema 1992; Beukema and Dekker 2005). For practical reasons, these stations  
152 were chosen to be close to each other but at different heights (from ~0.1 m above to ~0.4 m  
153 below MTL) and with different sediment characteristics (silt content ranging from 2-10%). The  
154 stations were located along a transect roughly perpendicular to the coast, at distances of 0.1-1 km  
155 from the shore and 0.3-1 km from a major tidal channel. The coordinates of the 3 stations are (1):  
156 52°55'02"N, 4°48'40"E, (2): 52°55'18"N, 4°48'51"E and (3): 52°55'26"N, 4°49'02"E.


157

### 158 **Sampling**

159 From 1983 to 2018 each spring, from April to at least late June, samples were collected weekly  
160 or bi-weekly during low tide (Fig. 2). The method, using a sampling corer, was used to include  
161 shrimps <10-15 mm in the samples, which were lost in historic samples due to the 5 mm mesh  
162 size that was commonly used (Beukema 1992). For this study, three stations were visited, at each  
163 of these stations 4 samples consisting of 10 pooled cores of 0.009 m<sup>2</sup> each were taken to a depth  
164 of ca. 5 cm and sieved over a 1 mm mesh (Beukema 1992). Up to and including 2009 material  
165 was taken back alive to the laboratory. From 2010 onwards all samples were preserved directly  
166 after sampling in the field with 4% formaldehyde and in the laboratory stained with rose bengal.

167 After sorting the samples in the laboratory each individual shrimp was measured to the

168 nearest mm from the tip of the scaphocerite to the end of the telson (Beukema 1992). For  
169 calculations, the data on shrimp were pooled per sampling event, resulting in a total area of 1.08  
170 m<sup>2</sup> sampled per sampling day. The first year (1983) was excluded from further analyses as the  
171 sampling period did not match well with those in subsequent years (see Fig. 2). For the analyses,  
172 the time window of 7 May – 24 June was selected as the only one covered in all years from 1984  
173 - 2018. Seawater temperatures were measured at the NIOZ jetty (53°00'06"N, 4°47'21"E) at -1.5  
174 m (relative to Amsterdam Ordnance Datum, NAP). The jetty is located at the tidal inlet nearest to  
175 the sampling stations 9 km away from the center of the three sampling stations (see van Aken  
176 2008). The water that passes the jetty is therefore representative for the water temperature that  
177 the larvae experience. Monthly averages were calculated and the yearly cumulative sum of the  
178 sea water temperatures from November until May. The cumulative sum of sea water  
179 temperatures was calculated with the water temperatures measured at 0800hr every day. Missing  
180 values were replaced by the mean of the measured temperatures before and after the gaps.



181  
182 **Fig. 2** Sampling period per year from 1983-2018. Rectangle marks time window that was used in  
183 the analyses: 7 May– 24 June 1984-2018

184  
185 **Shrimp length**

186 To examine seasonal changes in mean shrimp length, the mean shrimp length per week was  
187 calculated (Fig. S1). After inspecting year by year patterns for consistency and change, we split  
188 the full period (1984-2018) into four decennia: 1984-1989, 1990-1999, 2000-2009, 2010-2018.  
189 All measured lengths were plotted per time period per half month period. To find out if long-  
190 term changes in the average shrimp length happened, the mean shrimp lengths on 7 May, 31 May

191 and 24 June were calculated using linear interpolation between datapoints. The minimal shrimp  
192 length per sampling was calculated to see if changes in the length at settlement could be  
193 detected. The settlement of new shrimp on the mudflat may occur in waves that are related to the  
194 lunar cycle or large amplitude flood tides, as is the case in other Decapoda (Mense et al. 1995;  
195 Christy and Morgan 1998). First, we calculated the median shrimp length per sampling. Then,  
196 we scrutinized the correlations between median shrimp length per sampling and moon phase or  
197 tidal height to see if such a mechanism could exist in shrimp too. The moon phase one day  
198 preceding each sampling date was extracted with software (see Statistics). Tidal heights in Den  
199 Helder (52°57'52"N, 4°44'42"E) were collected by Rijkswaterstaat  
200 (<http://www.rijkswaterstaat.nl>). The water height of the highest high tide of the day preceding  
201 each sampling day in Den Helder was selected to calculate correlations between shrimp length  
202 and tidal height.

203

#### 204 **Densities and cumulative densities per time period**

205 The sum of shrimp per sampling day was divided by the sampling area (1.08 m<sup>2</sup>) to obtain the  
206 density of shrimp per square meter (Fig. S2). In years without a sampling occasion on 7 May or  
207 24 June, the density on these dates was obtained by linear interpolation between the adjacent data  
208 points. To show the timing of shrimps arriving on the sampled mudflats, we computed the  
209 ‘standardized cumulative shrimp density’: the shrimp density of the current week plus the  
210 density of the previous week (except for the first sample). The timing and number of  
211 observations per year are given in Figure 2. Between year comparisons were made using three  
212 computed variables: (1) the date when 50% of the annual cumulative shrimp density is reached,  
213 (2) the cumulative shrimp density on 31 May, halfway the sampling period, and (3) the shrimp  
214 density on 7 May, the starting date of sampling periods. The values were obtained by linearly  
215 interpolating between datapoints. A Pearson’s product-moment correlation test determined the  
216 correlation between the date when 50% of the standardized annual cumulative shrimp density  
217 was reached and the standardized cumulative shrimp density on 31 May.

218

#### 219 **Statistics**

220 All calculations and analyses were carried out using R (R Development Core Team 2020, R  
221 version 3.6.3). The distribution of the data was tested for normality with the

222 Shapiro Wilk test with no significant results. Temporal differences in shrimp density, timing and  
223 length were analyzed with linear models from the lm function in base R with year as explanatory  
224 variable. To check if the change in preservation and staining improved the detection of small  
225 shrimp, a segmented regression analysis was carried out using the R package “segmented”  
226 (Muggeo 2008). With this analysis sudden changes in the slope of a trend, in this case a change  
227 in the mean length from 2010 onwards, could be detected. The results of the segmented  
228 regression were compared with linear regression of the same data. The best model explained  
229 most variation.

230 The existence of settlement waves was studied in relation to the moon phase. The moon  
231 phase one day before each sampling date was extracted using the package “lunar” (Lazaridis  
232 2014). Differences in the median shrimp length between moon phases were analyzed in an  
233 analysis of variance (ANOVA). The association between the median shrimp length and the  
234 height of the tide was analyzed by linear regression.

235 To find out during which time of the year the water temperature has the strongest  
236 association with the shrimp density on 7 May and the 50% date, several linear regression  
237 analyses were carried out. The same was done to assess during which time of year the water  
238 temperature best explained the mean shrimp length on either 7 May, 31 May or 24 June. For  
239 each year, the mean water temperature of the separate months January, February, March and  
240 April were calculated to be used as explanatory variables. Secondly, the mean of the months  
241 January, February and March was calculated per year to represent the mean water temperature in  
242 winter, as in Beukema (1992). Lastly, the cumulative sum of the sea water temperature was calculated  
243 per year for the time period of November until May. The time of year (month, winter or cumulative  
244 sum) of which temperature showed the strongest correlation with shrimp density on 7 May and  
245 50% density date was then further analyzed based on the amount of variation explained. The  
246 relative importance of water temperature and year in explaining (1) the date at which 50% of the  
247 shrimp density was reached and (2) the shrimp density on 7 May, was analyzed with linear  
248 models. Time and temperature were tested separately and together, including and excluding a  
249 two-way interaction. The best model was selected on the basis of Akaike’s Information Criterion  
250 (AIC; Burnham and Anderson 2002).

251 Finally, shrimp density may not be independent of year as density in one year may be  
252 influenced by shrimp density in the preceding year. Therefore, an autocorrelation function was

253 plotted, based on the standardized residuals from the linear model that describes shrimp density.  
254 Visual inspection of this plot did not show indications for autocorrelation. Still, we checked for  
255 temporal autocorrelation by adding two types of autocorrelation structures from the gls function  
256 to the final model. The autocorrelation structures we used were: Compound Symmetry  
257 Correlation Structure (CompSymm) and First Order Autoregressive Structure AR-1 (Zuur et al.  
258 2007).

259


## 260 **Results**

### 261 **Length**

262 From 1984 to 2018 a total of 141 sampling occasions took place (Fig. 2). Over all the years, the  
263 estimated mean length on 7 May, the start of the study season, was  $9.69 \pm 0.091$  mm and the  
264 shrimp length on the last day of the study season 24 June was  $12.93 \pm 0.083$  mm (Fig. 3a). Based  
265 on visual inspection of the histograms in Fig. 3c, in May smaller shrimp occurred more than in  
266 June. However, throughout the years, median lengths as small as 5 mm occurred only on 1% of  
267 the sampling days. Across the 35 year study period, mean shrimp length in this time window  
268 decreased significantly from 12.6 mm to 10.7 mm (linear regression:  $r^2 = 0.22$ ,  $F_{1,33} = 9.49$ ,  $p =$   
269  $0.004$ ) (Fig. 3b). Visual inspection of the boxplots in Fig. 3d did not suggest that such decreases  
270 in length occurred during specific years. Indeed, the mean lengths on either 7 May, 31 May or 24  
271 June did not change over the years (linear regression: April  $F_{1,33} = 2.02$ ,  $p = 0.16$ , May  $F_{1,33} =$   
272  $1.03$ ,  $p = 0.32$ , June  $F_{1,33} = 3.33$ ,  $p = 0.07$ ). Segmented regression did not estimate a break point  
273 in the shrimp length after the year 2010 and the model explained less variation than the simple  
274 linear regression (segmented regression:  $r^2 = 0.19$ , linear regression:  $r^2 = 0.22$ ). Therefore, there  
275 is no indication that the results have been biased by the change in preservation and staining of  
276 samples from 2010 onwards.

277         Lengths on the dates when 50% of cumulative density was reached did not change over  
278 time either (linear regression:  $F_{1,33} = 1.70$ ,  $p = 0.20$ ). Furthermore, mean shrimp length on 7 May  
279 was not significantly correlated with the January temperature, nor the April water temperature  
280 but it was with water temperatures in winter, February and March (Table S3). The minimal  
281 shrimp length did not change over the season (linear regression:  $F_{1,141} = 0.0003$ ,  $p = 0.99$ ) and  
282 the median length per sampling was not correlated with a specific moon phase (ANOVA:  $F_{1,7} =$   
283  $0.75$ ,  $p = 0.63$ ) or the height of the preceding high tide (linear regression:  $F_{1,141} = 0.02$ ,  $p = 0.89$ ).

284 The same analyses were carried out per month separately to see if the effect would be present in  
 285 a specific time period but none of these tests results were significant either.  
 286




287  
 288 **Fig. 3** The shrimp length over time: seasonal and annual changes **a** mean shrimp length during  
 289 the sampling season, general additive model (GAM) was used as a smoother. **b** The mean shrimp  
 290 length per year, linear regression line plotted ( $r^2 = 0.22$ ,  $F_{1,33} = 9.49$ ,  $p = 0.004$ ). **c** The length  
 291 frequency distribution from 1984-2018 in May (light grey, grey border) and June (dark grey,  
 292 black border). **d** boxplots of the shrimp lengths per half month from 1984-2018. The titles give  
 293 the time period that was selected for each graph

294  
 295 **Density**

296 Changes in shrimp density over time were analyzed using the standardized shrimp density and  
 297 the density on 7 May (Fig. 4a). The date when 50% of the standardized annual cumulative  
 298 shrimp density was reached was strongly correlated with the standardized cumulative shrimp  
 299 density on 31 May (Pearson's product-moment correlation coefficient  $r = 0.93$ ,  $n = 35$ ,  $p < 0.001$ ).  
 300 Therefore, we only analyzed the date when 50% of the cumulative shrimp density was reached  
 301 and dropped the cumulative shrimp density on 31 May from further analyses. Mean shrimp

302 density (calculated per year) remained stable over time (linear regression:  $F_{1,33} = 2.89, p = 0.10$ ).  
 303 The date when 50% of the yearly cumulative shrimp density was reached advanced significantly  
 304 in the past 35 years at a slope of 0.34 days per year (Fig. 4b). In total, the 50% cumulative  
 305 shrimp density advanced with 12 days in the 35 years study period. Between 1984 and 2018 the  
 306 shrimp density on 7 May increased significantly by more than 2.4 times, from 28  $m^{-2}$  to 69  $m^{-2}$   
 307 (Fig. 4c).

308




309

310 **Fig. 4 a** The standardized cumulative shrimp density runs (0-1) and is plotted against time of  
 311 year. Each year is plotted separately in a different color. The solid line marks the middle of the  
 312 sampling period on 31 May. The dashed line marks the moment in time when 50% of the yearly  
 313 shrimp density has been reached. **b** The time of year when 50% of the cumulative shrimp density  
 314 has been reached through time (years). Linear regression line plotted ( $r^2 = 0.14, F_{1,33} = 5.22, p =$   
 315  $0.029$ ). **c** Shrimp density on 7 May, the start of the sampling season with the linear regression  
 316 line plotted ( $r^2 = 0.17, F_{1,33} = 6.72, p = 0.014$ ). Shaded area in b and c mark the 95% confidence  
 317 interval (CI)

318

319 **Sea water temperatures**

320 The correlations between seawater temperature and the 50% density date and shrimp density on  
 321 7 May were strongest with temperatures in April (Table S3). April seawater temperature  
 322 increased significantly in the course of the 35-year study period (Fig. 5a; linear regression:  $r^2$ :  
 323 0.35,  $F_{1,33} = 17.58$ ,  $p < 0.001$ ). The date when the cumulative shrimp density reached 50%  
 324 advanced by 3.4 days for every 1°C higher mean April seawater temperature ( $r^2 = 0.24$ ,  $F_{1,33} =$   
 325 10.55,  $p = 0.003$ , Fig. 5b). In line with this, shrimp density on 7 May was significantly higher  
 326 with high April seawater temperature ( $r^2 = 0.44$ ,  $F_{1,33} = 26.06$ ,  $p < 0.001$ , Fig. 5c). Shrimp density  
 327 increased with 14.7 m<sup>-2</sup> per 1°C higher mean April seawater temperature.  
 328



329  
 330 **Fig. 5 a** Average April seawater temperature over time. **b** The time of year when 50% of the  
 331 cumulative shrimp density has been reached. Linear regression line plotted ( $r^2 = 0.24$ ,  $F_{1,33} =$   
 332 10.55,  $p = 0.003$ ). **c** the shrimp density on 7 May, plotted against the mean April seawater  
 333 temperature. Linear regression plotted ( $r^2 = 0.44$ ,  $F_{1,33} = 26.06$ ,  $p < 0.001$ ). In **b** and **c** each point  
 334 represents a year, the shaded areas mark the 95% CI

335  
 336 Based on lowest AIC values, April water temperatures best explained the timing of  
 337 arrival of shrimp on the mudflats (Table 1). For both response variables that were tested (timing  
 338 of 50% of yearly cumulative density and shrimp density on 7 May), year did not contribute  
 339 significantly to the explanation of variation. Models including an autocorrelation structure did

340 not have a lower AIC value (model without autocorrelation structure AIC: 327.6, models with  
 341 autocorrelation structure AIC: 329.6).

342  
 343 **Table 1.** Comparison of linear regression models describing 1) the timing of settlement: date  
 344 when 50% of annual cumulative shrimp density is reached and 2) shrimp density on 7 May.  
 345 Independent variables: year and April seawater temperature. Variance explained ( $r^2$ ) and the p-  
 346 value of the models with the lowest AIC are indicated in bold.

<i>Model terms</i>	<i>ac-structure</i>	<i>AIC</i>	<i>r<sup>2</sup></i>	<i>p</i>
50% day ~ water temp. + year		253.82		
50% day ~ water temp. * year		255.64		
<b>50% day ~ water temp.</b>		<b>252.27</b>	<b>0.24</b>	<b>0.002</b>
50% day ~ year		256.84		
7 May density ~ water temp. + year		324.65		
7 May density ~ water temp. * year		326.16		
<b>7 May density ~ water temp.</b>		<b>322.68</b>	<b>0.44</b>	<b>&lt;0.001</b>
7 May density ~ year		336.57		
50% day ~ water temp. + year		262.88		
50 % day ~ water temp. + year	CompSymm	264.88		
50 % day ~ water temp. + year	AR1	264.88		
7 May density ~ water temp. + year		327.63		
7 May density ~ water temp. + year	CompSymm	329.63		
7 May density ~ water temp. + year	AR1	329.63		

347

348

## 349 **Discussion**

350 We show that in the 35 years from 1984 to 2018 the settlement of juvenile post-larval brown  
 351 shrimp on the intertidal flats of the Dutch Wadden Sea advanced by 12 days. Additionally,  
 352 densities at the start of the study season more than doubled, with an estimated overall average  
 353 shrimp density on 7 May of 69 m<sup>-2</sup>. The mean length of shrimp at the start of the season did not  
 354 change, nor was it correlated with water temperatures in April. Instead, mean shrimp length on 7  
 355 May was correlated strongest with the mean water temperature of the winter months. In

356 summary, in warm springs we find higher shrimp densities earlier in the sampling season and at  
357 that time they are not of a larger size compared to colder springs.

358 The correlation between the advanced appearance of shrimp and sea water temperature  
359 was higher for the temperatures in April than for the temperatures in the whole previous winter  
360 season, and indeed became better as time approached the time of appearance of shrimp, with the  
361 highest correlations for the April seawater temperatures. This is consistent with various known  
362 biological processes such as egg development and larval growth, both of which speed up with  
363 higher water temperatures (Wear 1974, Criales & Anger 1986, Paschke et al. 2004, Hufnagl and  
364 Temming 2011). For the field, such differences in development time also have implications for  
365 the spatial origin of the post-larvae that settle in the Wadden Sea. Shrimp larvae drift passively  
366 with currents until they become post-larvae (Temming and Damm 2002). At low temperatures  
367 the larvae will drift for longer because development is slower (Daewel et al. 2011). The time  
368 period that larvae are exposed to pelagic predators is therefore also extended, possibly leading to  
369 higher mortality rates. Wind speed and direction influence the drifting track as well (Daewel et  
370 al. 2011) and could cause differences in the timing and abundance of post-larvae that settle in the  
371 Wadden Sea.

372 Post-larval settlement of other crustaceans is known to take place in waves correlated  
373 with the moon phase or the amplitude of the high tide. For the post-larval shrimp on the intertidal  
374 flats in the westernmost Dutch Wadden Sea, we did not find that median length was correlated  
375 with the moon phase or the height of the tide. This rejects a hypothesis for post-larval shrimp  
376 settlement to occur in moon or tide-driven waves. The idea that settlement could especially  
377 happen during nocturnal flood tides (Christy and Morgan 1998), could not be evaluated with our  
378 data.

379 Over the course of the sampling season, the mean shrimp length increased from 9.7 mm  
380 to 12.9 mm. This increase might suggest that post-larval shrimp move to the mudflat at larger  
381 sizes. However, this seems unlikely, as we did not find a change in the minimal shrimp length  
382 over the season. Also, the increase in mean length is too low to be attributed to growth. On the  
383 basis of measured individual growth rates of 0.28 mm per day for a 10 mm shrimp in water of 10  
384 °C (Hufnagl and Temming 2011), the individual increase in body length would be expected to be  
385 12.3 mm over the 44 day study period, i.e. almost four times the ‘growth’ in the field. Even with  
386 this conservative scenario (the mean water temperature in May is actually 12 °C), the growth

387 speed is clearly too high to explain the increase in the mean shrimp length. This suggests that the  
388 mean shrimp length decreased because of an increase in the turnover of the population, with the  
389 larger juveniles leaving for deeper waters at ever faster rates. Indeed, departure of shrimp from  
390 intertidal flats may start at 15 mm (Hufnagl et al. 2014) instead of 20 mm (Beukema 1992),  
391 which would have caused a steeper decrease of larger length classes which would have kept the  
392 observed increase in mean shrimp length across the season small. Whether the departure of  
393 juveniles is triggered by changes in the sensitivity to hydrostatic pressure just as in adults, will  
394 require further investigation.

395         The advanced settlement on intertidal flats was best correlated with the seawater  
396 temperatures in April (which increased over the years). The mechanisms underlying this pattern  
397 remain to be established, but on the basis of the observational data presented we propose four  
398 non-mutually exclusive interpretations of the changes in timing of shrimp on the mudflat: (1)  
399 shrimp settle on the mudflat when they reach a certain ‘threshold’ length, (2) the settlement of  
400 shrimp is controlled by a critical period of ‘threshold’ temperature sensitivity, (3) the timing of  
401 shrimp settlement is a response to food availability on the mudflats and/or (4) shrimp settlement  
402 on intertidal flats is behaviour to avoid predation by other species and cannibalism by adult  
403 shrimp. Predictions on climate effects on future shrimp lengths at time of settlement and shrimp  
404 densities in spring will differ between the four different hypotheses.

405         For the first interpretation, assuming a threshold length for settlement of 4.7 mm (i.e. the  
406 length when shrimp reach the post-larval phase; Kuipers and Dapper 1984), this length should be  
407 the dominant length, at least during the sampling early in the season. However, median lengths  
408 as small as 5 mm occurred only on 1% of the sampling days. Either the peak of settlement  
409 already passed by the time that sampling started, or settled post-larval shrimp spend time  
410 elsewhere before moving onto the intertidal flats. Regardless of the precise threshold length at  
411 settlement, in the light of climate change no changes from this threshold length are expected if  
412 body length itself triggers settlement. Advancements in the timing of settlement would occur if  
413 this length would be reached earlier in the year. With increasing water temperatures, growth rates  
414 will increase during all life stages of shrimp (Wear 1974, Criales & Anger 1986, Paschke et al.  
415 2004). This actually may occur, as Beukema and Dekker (1992, 2014) showed advanced shrimp  
416 settlement and higher spring shrimp densities after mild winters.

417                   However, by comparing the correlations between shrimp density on 7 May and  
418 the water temperature in preceding time periods, we discovered that the water temperature in  
419 April shows the strongest correlations. This cannot be explained by greater growth rates at higher  
420 water temperatures, but rather hints at the existence of a threshold temperature during a sensitive  
421 time window. Here, we assume that shrimp go through a developmental period during which  
422 they are sensitive for a threshold in the sea water temperature. When this threshold is reached,  
423 the shrimp settle on the mudflat. Under this interpretation shrimp would settle on the mudflat  
424 when the temperature conditions are right (rather than when they reach a certain length or life  
425 stage). With increasing water temperatures, the temperature threshold may be reached earlier in  
426 the year, leading to the observed advancement in the timing that shrimp settle on intertidal  
427 mudflats. That the water temperature in the preceding months will lead up to the water  
428 temperature in April may then explain why water temperatures during the preceding months also  
429 correlate with shrimp densities in May. In the presence of a temperature threshold, shrimp may  
430 actually settle on the mudflat at the minimal post-larval length of 4.7 mm after warm winters and  
431 at larger sizes after cold winters. Due to the timing of sampling we can not confirm this with our  
432 data. Survival and growth rates are lower at low temperatures, but shrimp may still grow in water  
433 below 10°C (Rochanaburanon and Williamson 1976; Hufnagl and Temming 2011). Therefore,  
434 during a cold spring, the temperature threshold will be reached later, leading to individuals that  
435 settle on the mudflat when they are larger. Again, assuming that in the subtidal shrimp do show  
436 growth, in warm springs the temperature threshold would be reached earlier in the year, resulting  
437 in smaller individuals that settle on the mudflat.

438                   The third interpretation is that the advance in shrimp settlement is based on increases or  
439 peaks in food availability occurring earlier. The diet of shrimp <20 mm mainly consists of  
440 meiofauna (Pihl and Rosenberg 1984; del Norte-Campos and Temming 1994). As a result of  
441 increased water temperatures, the phenology of meiofauna may be advanced. Shrimp may follow  
442 their prey, with advanced settlement on the mudflats as a result. Assuming that with increased  
443 water temperatures the food of shrimp becomes available earlier in the year, shrimp will advance  
444 settlement on the mudflat. Additionally, they could still be of a smaller size, as earlier in the  
445 growing season shrimp will still be smaller.

446                   Our fourth and last alternative interpretation is an advancement of high predation  
447 pressure in the subtidal induces post-larval shrimp to move to intertidal mudflats earlier in the

448 year. Currently, post-larval settled shrimp are thought to be absent in the subtidal - they have  
449 rarely been found there in the past. At the same time, because previously used mesh sizes were  
450 too wide, sampling may have failed to detect small shrimp <10 mm properly (Beukema 1992), or  
451 started too late. Earlier we already concluded that settled post-larval shrimp may spend time  
452 somewhere deeper before moving onto the intertidal flats.

453 Observational studies (including comparisons between different areas in e.g. the Wadden  
454 Sea), and experimental studies on the mechanisms on shrimp settlement, are now required to  
455 elucidate why our measurements of ‘settlement’ are correlated with water temperatures  
456 especially in April. Noting that the different interpretations will lead to different scenarios of  
457 change in a warming world, without them it will be impossible to interpret the striking changes  
458 in phenology. As shrimp are now present in the intertidal earlier in the year, different predators  
459 may benefit from their presence. To fully understand the consequences for higher trophic levels,  
460 it is necessary to look into the juvenile shrimp abundances over a longer time in the year and  
461 identify if new matches or mismatches between shrimp and consumer exist.

462

### 463 **Acknowledgements**

464 This study was part of project ‘Behoud van een dynamisch Griend’ managed by  
465 Natuurmonumenten and funded by Waddenfonds, the provinces of Fryslân, Groningen and  
466 Noord-Holland, and Rijkswaterstaat. LLG was funded by NWO grant 016.Veni.181.087.  
467 Exploratory analyses were carried out by Arjen de Boer and Lies Bosma during their BSc course  
468 in 2019 at the University of Groningen. We would like to thank all assistants who helped with  
469 collecting and sorting the Balgzand dataset. Thanks to Sonja van Leeuwen for maintaining the  
470 NIOZ jetty temperature database. We are grateful to the anonymous reviewers for their most  
471 constructive feedback.

### 472 **Author contributions**

473 EP, LG, TP and RD contributed to the concept and design of the study. RD conducted the data  
474 collection together with Jan Beukema and EP ran the statistical analysis. EP wrote the main  
475 sections of the manuscript supervised by TP and with input from LG and RD. All authors  
476 approved the submitted version.

477 **Funding**

478 EP was funded by Waddenfonds grant WF-223011 to a consortium of Natuurmonumenten,  
479 Royal NIOZ and University of Groningen for the restoration of Griend, the Dutch Wadden Sea  
480 provinces (Friesland, Groningen, Noord-Holland) and Rijkswaterstaat. LLG was funded by  
481 NWO grant 016.Veni.181.087.

482 **Data availability**

483 The datasets generated during and/or analyzed during the current study will become available in  
484 the Zenodo Repository (<http://doi.org/10.5281/zenodo.4792217>).

485 **Compliance with ethical standards**

486 **Conflict of interest**

487 The authors have no conflicts of interests or competing interests to declare.

488

489 **Ethics approval**

490 This research was conducted in accordance with the Netherlands Code of Conduct for Scientific  
491 Practice. All applicable international, national, and institutional guidelines for sampling, care,  
492 and experimental use of organisms for the study have been followed.

493

494 **References**

- 495 Al-Adhub AHY, Naylor E (1975) Emergence rhythms and tidal migrations in the brown shrimp *Crangon crangon*  
496 (L.). J Mar Biol Assoc UK 55:801–810. <https://doi.org/10.1017/S0025315400017720>
- 497 Andresen H, van der Meer J (2010) Brown shrimp (*Crangon crangon*, L.) functional response to density of different  
498 sized juvenile bivalves *Macoma balthica* (L.). J Exp Mar Biol Ecol 390:31–38.  
499 <https://doi.org/10.1016/j.jembe.2010.04.027>
- 500 Beukema JJ (1992) Dynamics of juvenile shrimp *Crangon crangon* in a tidal-flat nursery of the Wadden Sea after  
501 mild and cold winter. Mar Ecol Prog Ser 83:157–165
- 502 Beukema JJ, Dekker R (2005) Decline of recruitment success in cockles and other bivalves in the Wadden Sea:  
503 Possible role of climate change, predation on postlarvae and fisheries. Mar Ecol Prog Ser 287:149–167.  
504 <https://doi.org/10.3354/meps287149>
- 505 Beukema JJ, Dekker R (2014) Variability in predator abundance links winter temperatures and bivalve recruitment:  
506 correlative evidence from long-term data in a tidal flat. Mar Ecol Prog Ser 513:1–15.  
507 <https://doi.org/10.3354/meps10978>
- 508 Boddeke R (1976) The seasonal migration of the brown shrimp *Crangon crangon*. Neth J Sea Res 10:103–130.  
509 [https://doi.org/10.1016/0077-7579\(76\)90006-5](https://doi.org/10.1016/0077-7579(76)90006-5)
- 510 Boddeke R (1982) The occurrence of winter and summer eggs in the brown shrimp (*Crangon crangon*) and the

- 511 pattern of recruitment. *Neth J Sea Res* 162:151–162. [https://doi.org/10.1016/0077-7579\(82\)90026-6](https://doi.org/10.1016/0077-7579(82)90026-6)
- 512 Boddeke R (1996) Changes in the brown shrimp (*Crangon crangon* L.) population off the Dutch coast in relation to  
513 fisheries and phosphate discharge. *ICES J Mar Sci* 53:995–1002. <https://doi.org/10.1006/jmsc.1996.0124>
- 514 Boddeke R, Becker R (1979) A quantitative study of the fluctuations of the stock of brown shrimps along the coast  
515 of the Netherlands. *Rapp P-v Réun Cons Int Explor Mer* 175: 253-258
- 516 Boddeke R, Hagel P (1991) Eutrophication on the North Sea continental zone, a blessing in disguise. ICES Council  
517 Meeting Papers, ICES CM 1991/E:7
- 518 Boddeke R, Driessen G, Doesburg W, Ramaekers G (1985) Settlement and growth of brown shrimp (*Crangon*  
519 *crangon*) in a coastal area. IJmuiden, The Netherlands: International Council for the Exploration of the Sea  
520 Conference Meeting 1985/K 16:8
- 521 Broekema MMM (1941) Seasonal movements and the osmotic behaviour of the shrimp, *Crangon crangon* L. *Arch*  
522 *Néerl Zool* 6:1–102
- 523 Burnham KP, Anderson DR (2002) *Model Selection and Multimodel Inference*. Springer, New York, New York
- 524 Campos J (2009) The eco-geography of the brown shrimp *Crangon crangon* in Europe. Disstertation, Vrije  
525 Universiteit, Amsterdam, The Netherlands, p 171
- 526 Campos J, van der Veer HW (2008) Autecology of *Crangon crangon* (L.) with an emphasis on latitudinal trends.  
527 *Oceanogr Mar Biol* 46:65–104. <https://doi.org/10.1201/9781420065756.ch3>
- 528 Campos J, Freitas V, Pedrosa C, et al. (2009) Latitudinal variation in growth of *Crangon crangon* (L.): Does  
529 counter-gradient growth compensation occur? *J Sea Res* 62:229–237.  
530 <https://doi.org/10.1016/j.seares.2009.04.002>
- 531 Christianen MJA, Middelburg JJ, Holthuijsen SJ, et al. (2017) Benthic primary producers are key to sustain the  
532 Wadden Sea food web: stable carbon isotope analysis at landscape scale. *Ecology* 98:1498–1512.  
533 <https://doi.org/10.1002/ecy.1837>
- 534 Christy JH, Morgan SG (1998) Estuarine immigration by crab postlarvae: mechanisms, reliability and adaptive  
535 significance. *Mar Ecol Prog Ser* 174: 51-65. doi:10.3354/meps174051
- 536 Criales MM, Anger K (1986) Experimental studies on the larval development of the shrimps *Crangon crangon* and  
537 *C. allmanni*. *Helg Meeresunters* 40:241–265. <https://doi.org/10.1007/BF01983735>
- 538 Daewel U, Schrum C, Temming A (2011) Towards a more complete understanding of the life cycle of brown  
539 shrimp (*Crangon crangon*): modelling passive larvae and juvenile transport in combination with physically  
540 forced vertical juvenile migration. *Fish Oceanogr* 20: 479–496
- 541 Dapper R, van der Veer HW (1981) Onderzoek naar de ruimtelijke variatie van de bodemsamenstelling op het  
542 Balgzand. Internal report no. 1981-9. Royal Netherlands Institute for Sea Research, Texel.
- 543 del Norte Campos AGC, Temming A (1994) Daily activity, feeding and rations in gobies and brown shrimp in the  
544 northern Wadden Sea. *Mar Ecol Prog Ser* 115:41–54. <https://doi.org/10.3354/meps115041>
- 545 Freitas V, Campos J, Fonds M, van der Veer H (2007) Potential impact of temperature change on epibenthic predator-  
546 bivalve prey interactions in temperate estuaries. *J Therm Biol* 32:328-340. 10.1016/j.jtherbio.2007.04.004.
- 547 Gibson RN, Yin MC, Robb L (1995) The behavioural basis of predator-prey size relationships between shrimp  
548 (*Crangon crangon*) and juvenile plaice (*Pleuronectes platessa*). *J Mar Biol Assoc UK* 75:337–349.  
549 <https://doi.org/10.1017/S002531540001821X>
- 550 Hartsuyker L (1966) Daily tidal migrations of the shrimp, *Crangon crangon* L. *Neth J Sea Res* 3:52–67.  
551 [https://doi.org/10.1016/0077-7579\(66\)90006-8](https://doi.org/10.1016/0077-7579(66)90006-8)
- 552 Hayward PJ, Ryland JS (2017) *Handbook of the Marine Fauna of North-West Europe.*, Second edi. Oxford  
553 University Press, Oxford

- 554 Hufnagl M, Temming A (2011) Growth in the brown shrimp *Crangon crangon*. II. Meta-analysis and modelling.  
555 Mar Ecol Prog Ser 435: 155-172. <https://doi.org/10.3354/meps09224>
- 556 Hufnagl M, Temming A and Pohlmann T (2014) The missing link: tidal-influenced activity a likely candidate to  
557 close the migration triangle in brown shrimp *Crangon crangon* (Crustacea, Decapoda). Fish Oceanogr 23:  
558 242-257. <https://doi.org/10.1111/fog.12059>
- 559 Hünnerlage K, Siegel V, Saborowski R (2019) Reproduction and recruitment of the brown shrimp *Crangon crangon*  
560 in the inner German Bight (North Sea): An interannual study and critical reappraisal. Fish Oceanogr. 28 708-  
561 722. <https://doi.org/10.1111/fog.12453>ICES (2015) Report of the Working Group on Crangon Fisheries and  
562 Life History (WGCRAN). ICES CM 2015/SSGEPD: 07
- 563 Janssen GM, Kuipers BR (1980) On tidal migration in the shrimp *Crangon crangon*. Neth J Sea Res 14:339–348.  
564 [https://doi.org/10.1016/0077-7579\(80\)90007-1](https://doi.org/10.1016/0077-7579(80)90007-1)
- 565 Jensen KT, Jensen JN (1985) The importance of some epibenthic predators on the density of juvenile benthic  
566 macrofauna in the Danish Wadden Sea. J Exp Mar Biol Ecol 89: 157–174
- 567 Jouta J, de Goeij P, Lok T, et al. (2018) Unexpected dietary preferences of Eurasian spoonbills in the Dutch Wadden  
568 Sea: spoonbills mainly feed on small fish not shrimp. J Ornithol 159:839–849. [https://doi.org/10.1007/s10336-](https://doi.org/10.1007/s10336-018-1551-2)  
569 018-1551-2
- 570 Kuipers BR, Dapper R (1984) Nursery function of Wadden Sea tidal flats for the brown shrimp *Crangon crangon*.  
571 Mar Ecol Prog Ser 17:171–181
- 572 Kuipers BR, Dapper R (1981) Production of *Crangon crangon* in the tidal zone of the Dutch Wadden Sea. Neth J  
573 Sea Res 15:33–53
- 574 Lazaridis E (2014) lunar: Lunar Phase & Distance, Seasons and Other Environmental Factors (Version 0.1-04).  
575 Available from <http://statistics.lazaridis.eu>
- 576 Lloyd AJ, Yonge CM (1947) The biology of *Crangon vulgaris* in the Bristol Channel and Severn Estuary. J Mar  
577 Biol Assoc UK 26:626–661
- 578 Marchand J (1981) Observations sur l'écologie de *Crangon crangon* (Linné) et *Palaemon longirostris* (H. Milne  
579 Edward) (Crustacea, Decapoda, Natantia). Estuaire interne de la Loire (France). Vie Milieu 31:83–92
- 580 Mathot KJ, Piersma T, Elner RW (2019) Shorebirds as integrators and indicators of mudflat ecology. In: Beninger  
581 PG (ed) Mudflat Ecology. Springer-Nature, Cham, pp 309–338
- 582 Mense DJ, Posey MH, West T, Kincheloe K (1995) Settlement of brachyuran postlarvae along the North Carolina  
583 coast. Bull Mar Sci 57: 793-806.
- 584 Muggeo VMR (2008) Segmented: an R package to fit regression models with broken-line relationships. R News 8/1,  
585 20–25.
- 586 Nehls G, Tiedemann R (1993) What determines the densities of feeding birds on tidal flats? A case study on dunlin  
587 *Calidris alpina* in the Wadden Sea. Neth J Sea Res 31:375–384. [https://doi.org/10.1016/0077-7579\(93\)90054-](https://doi.org/10.1016/0077-7579(93)90054-V)  
588 V
- 589 Oh CW, Hartnoll RG, Nash RDM (1999) Population dynamics of the common shrimp, *Crangon crangon* (L.), in  
590 Port Erin Bay, Isle of Man, Irish Sea. ICES J Mar Sci 56:718–733. <https://doi.org/10.1006/jmsc.1999.0501>
- 591 Paschke KA, Gebauer P, Buchholz F, Anger K (2004) Seasonal variation in starvation resistance of early larval  
592 North Sea shrimp *Crangon crangon* (Decapoda: Crangonidae). Mar Ecol Prog Ser 279:183–191
- 593 Philippart CJM, van Aken HM, Beukema JJ, et al. (2003) Climate-related changes in recruitment of the bivalve  
594 *Macoma balthica*. Limnol Oceanogr 48:2171–2185. <https://doi.org/10.4319/lo.2003.48.6.2171>
- 595 Philippart CJM, Beukema JJ, Cadée GC, et al. (2007) Impacts of nutrient reduction on coastal communities.  
596 Ecosystems 10:95–118. <https://doi.org/10.1007/s10021-006-9006-7>

- 597 Piersma T, de Goeij P, Tulp, I (1993) An evaluation of intertidal feeding habitats from a shorebird perspective:  
598 towards comparisons between temperate and tropical mudflats. *Neth J Sea Res* 31: 503-512.  
599 [https://doi.org/10.1016/0077-7579\(93\)90062-W](https://doi.org/10.1016/0077-7579(93)90062-W)
- 600 Piersma T, van Aelst R, Kurk K, Berkhoudt H, Maas LRM (1998) A new pressure sensory mechanism for prey  
601 detection in birds: the use of principles of seabed dynamics? *Proc R Soc Lond B* 265, 1377-1383.  
602 <https://doi.org/10.1098/rspb.1998.0445>
- 603 Pihl L, Rosenberg R (1984) Food selection and consumption of the shrimp *Crangon crangon* in some shallow  
604 marine areas in Western Sweden. *Mar Ecol Prog Ser* 15:159–168. <https://doi.org/10.3354/meps015159>
- 605 R Core Team (2020) R: A language and environment for statistical computing. R Foundation  
606 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 607 Rakhimberdiev E, Duijns S, Karagicheva J, et al. (2018) Fuelling conditions at staging sites can mitigate Arctic  
608 warming effects in a migratory bird. *Nat Commun* 9:1–10. <https://doi.org/10.1038/s41467-018-06673-5>
- 609 Rochanaburanon T, Williamson DI (1976) Laboratory survival of larvae of *Palaemon elegans* Rathke and other  
610 Caridean shrimps in relation to their distribution and ecology. *Estuar Coast Mar Sci* 4: 83-91
- 611 Siegel V, Damm U, Neudecker T (2008) Sex-ratio, seasonality and long-term variation in maturation and spawning  
612 of the brown shrimp *Crangon crangon* (L.) in the German Bight (North Sea). *Helgol Mar Res* 62 339–349.  
613 <https://doi.org/10.1007/s10152-008-0121-z>
- 614 Smaldon G (1979) British Coastal Shrimps and Prawns. Keys and notes for the Identification of the Species.  
615 Synopses of the British Fauna. Academic Press, London
- 616 Spaargaren DH (2000) Seasonal and annual variations in the catches of *Crangon crangon* (L., 1758) (Decapoda,  
617 Natantia) near the coast of Texel, The Netherlands. *Crustaceana* 73:547–563.  
618 <https://doi.org/10.1163/156854000504651>
- 619 Temming A, Damm U (2002) Life cycle of *Crangon crangon* in the North Sea: a simulation of the timing of  
620 recruitment as a function of the seasonal temperature signal. *Fish Oceanogr* 11: 45-58.  
621 <https://doi.org/10.1046/j.1365-2419.2002.00184.x>
- 622 Temming A, Günther C, Rückert C, Hufnagl M (2017) Understanding the life cycle of North Sea brown shrimp  
623 *Crangon crangon*: a simulation model approach. *Mar Ecol Prog Ser* 584:119–143.  
624 <https://doi.org/10.3354/meps12325>
- 625 Tielmann M, Reiser S, Hufnagl M, Herrmann J-P, Eckardt A, Temming A (2015) Hydrostatic pressure affects  
626 selective tidal stream transport in the North Sea brown shrimp (*Crangon crangon*). *J Exp Biol* 218: 3241-  
627 3248. <https://doi.org/10.1242/jeb.125773>
- 628 Tiews K (1970) Synopsis of biological data on the common shrimp *Crangon crangon*. *FAO Fish Rep* 57:1166–1224
- 629 Tulp I, Chen C, Haslob H, et al. (2016) Annual brown shrimp (*Crangon crangon*) biomass production in  
630 Northwestern Europe contrasted to annual landings. *ICES J Mar Sci* 73:fsw141.  
631 <https://doi.org/10.1093/icesjms/fsw141>
- 632 Tulp I, Glorius S, Rippen A, et al. (2020) Dose-response relationship between shrimp trawl fishery and the  
633 macrobenthic fauna community in the coastal zone and Wadden Sea. *J Sea Res* 156:1-13.  
634 <https://doi.org/10.1016/j.seares.2019.101829>
- 635 van Aken H (2008) Variability of the water temperature in the western Wadden Sea on tidal to centennial time  
636 scales. *J Sea Res* 60:227–234. <https://doi.org/10.1016/j.seares.2008.09.001>
- 637 van de Kam J, Ens B, Piersma T, Zwarts L (2004) Shorebirds: an illustrated behavioural ecology. KNNV Publishers,  
638 Utrecht
- 639 van der Baan, S (1975) Migration of *Crangon crangon* in surface waters near the "Texel" lightship. *Neth J Sea Res*  
640 9:287-296. [https://doi.org/10.1016/0077-7579\(75\)90004-6](https://doi.org/10.1016/0077-7579(75)90004-6)

- 641 van der Heide T, Tielens E, van der Zee EM, Weerman EJ, Holthuijsen S, Eriksson BK, Piersma T, van de Koppel J,  
642 Olf, H. (2014) Predation and habitat modification synergistically interact to control bivalve recruitment on  
643 intertidal mudflats. *Biol Cons* 172:163-169. <http://dx.doi.org/10.1016/j.biocon.2014.02.036>
- 644 van der Veer HW, Bergman MJN (1987) Predation by crustaceans on a newly settled 0-group plaice *Pleuronectes*  
645 *platessa* population in the Western Wadden Sea. *Mar Ecol Prog Ser* 35:203–215.  
646 <https://doi.org/10.3354/meps035203>
- 647 van der Veer HW, Feller RJ, Weber A, Witte JI (1998) Importance of predation by crustaceans upon bivalve spat in  
648 the intertidal zone of the Dutch Wadden Sea as revealed by immunological assays of gut contents. *J Exp Mar*  
649 *Biol Ecol* 231:139–157. [https://doi.org/10.1016/S0022-0981\(98\)00090-2](https://doi.org/10.1016/S0022-0981(98)00090-2)
- 650 van Lissa JHL (1977) Aantallen, voedselopname, groei en produktie van de garnaal (*Crangon crangon* L.) in een  
651 getijdengebied, alsmede de voedselopname en groei onder laboratoriumomstandheden. Internal report no.  
652 1977-10. Royal Netherlands Institute for Sea Research, Texel
- 653 Wear RG (1974) Incubation in British decapod crustacea, and the effects of temperature on the rate and success of  
654 embryonic development. *J Mar Biol Assoc UK* 54:745–762. <https://doi.org/10.1017/S0025315400022918>
- 655 Zuur A, Ieno EN, Smith GM (2007) *Analyzing ecological data*. Springer-Verlag, New York

## Supplementary material

### Title:

Advancing presence and changes in body size of brown shrimp *Crangon crangon* on intertidal flats in the western Dutch Wadden Sea, 1984-2018

### Authors:


Emma Penning<sup>1,2,\*</sup>, Laura L. Govers<sup>1,2</sup>, Rob Dekker<sup>1</sup>, Theunis Piersma<sup>1,2</sup>

### Author affiliations:


<sup>1</sup>NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands

<sup>2</sup>Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands

\*Corresponding author: emma.penning@nioz.nl



**Fig. S1** Mean shrimp length per sampling occasion per year. Rectangles mark the time window that was used in the analyses: 7 May– 24 June 1984-2018. The five coldest and the five warmest years are plotted in blue and red respectively, all other years in green



**Fig. S2** Mean shrimp density per sampling day per year. Rectangles mark the time window that was used in the analyses: 7 May– 24 June 1984–2018. The five coldest and the five warmest years are plotted in blue and red respectively, all other years in green

**Table S3** The correlations between the mean seawater temperature and: 1) the mean shrimp length at the start (7 May), middle (31 May) and end (24 June) of the study season, 2) the shrimp density on 7 May, and 3) the date that 50% of the standardized cumulative shrimp density is reached

	Mean shrimp length						Density				
	7-May		31-May		24-Jun		7 May density		date of 50% density		
	$r^2$	$p$	$r^2$	$p$	$r^2$	$p$	$r^2$	$p$	$r^2$	$p$	
Water temp.											
winter	0.18	0.011	0.17	0.015	0.07	0.133	0.37	<0.001	0.06	0.142	
cum. temp.	0.12	0.045	0.11	0.048	0.01	0.527	0.39	<0.001	0.12	0.044	
January	0.10	0.065	0.04	0.280	0.01	0.680	0.23	0.004	0.01	0.595	
February	0.18	0.012	0.10	0.064	0.09	0.082	0.24	0.003	0.03	0.310	
March	0.14	0.030	0.33	<0.001	0.10	0.070	0.40	<0.001	0.16	0.016	
April	0.08	0.104	0.08	0.102	0.06	0.154	0.44	<0.001	0.24	0.003	