



A multi-scale tracking approach for conserving large migratory fish in an open coastal environment

J.E. Edwards^{a,b,*}, A.D. Buijse^{b,c}, H.V. Winter^{b,d}, A. van Leeuwen^a, A.I. Bijleveld^a

^a NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems, PO Box 59, 1790 AB, Den Burg, Texel, the Netherlands

^b Wageningen University & Research, Aquaculture and Fisheries Group, Droevendaalsesteeg 4, 6708 PB, Wageningen, the Netherlands

^c Deltares, Boussinesqweg, 1 2629 HV, Delft, the Netherlands

^d Wageningen Marine Research, Haringkade 1, 1976 CP, IJmuiden, the Netherlands

ARTICLE INFO

Regional index terms:

The Netherlands
Dutch Wadden Sea

Keywords:

Movement ecology
Ecosystem management
Migration
Bio-logging
Biotelemetry
Wadden sea
Chelon labrosus
Dicentrarchus labrax
Salmo trutta
Galeorhinus galeus

ABSTRACT

Coastal habitats serve essential roles in the life cycles of migratory fishes, impacting both local and regional population stability. Conservation efforts for coastal fish often rely on spatial approaches within designated boundaries to mitigate threats and enhance production. However, in open coastal environments, migratory behaviours can extend beyond these protected areas, exposing individuals to potential threats or population bottlenecks elsewhere in their range. To improve conservation outcomes, a comprehensive understanding of movements across the entire migratory range is essential. Aquatic telemetry is a valuable tool for studying these behaviours, but must be adapted to address questions at multiple spatial and temporal scales. In this study we demonstrate how a combination of telemetry techniques can capture both local and regional fish behaviours. We begin by introducing the implications of migratory behaviours for fish conservation in open coastal environments, using the Wadden Sea as an example where additional research and management are needed to address fish declines. We then present a case study which uses the Dutch Wadden Sea to illustrate how a multi-scale telemetry approach can enhance both fundamental knowledge and conservation strategies for migratory fish. Within this case study, we present the general movement strategies exhibited by coastal migrants, alongside an overview of telemetry techniques applicable for open coastal systems. We then apply a size-based assessment using a reference tag to estimate the suitability of common Wadden Sea species for long-term tracking. Drawing from these results, we select four example species to showcase how species-specific understanding of fish life history and abundance can guide tracking studies, accompanied by illustrative examples using telemetry data. Expanding from this case study, we transition to a broader discussion where we provide overarching perspectives on tracking in open coastal ecosystems and offer recommendations to enhance future tracking studies in the Wadden Sea. The integration of telemetry methodologies aligns research and management with fish movement over multiple scales, improving our understanding of fish behaviors and contributing to more effective conservation strategies.

1. Background

1.1. Fish conservation in open coastal ecosystems

Positioned at the intersection of inland fresh waters and the pelagic marine realm, coastal habitats form an innate link between distinct aquatic ecosystems and are therefore of interest for studies on connectivity and the role of migration in aquatic animal life histories. Coastal marine habitats and estuaries play essential roles for numerous

migratory fish species, which typically have life cycles involving use of spatially-discrete habitats at certain life stages or times of year, punctuated by periods of large-scale movement. For migratory fish, coastal habitats serve as seasonal foraging grounds, spawning areas, and nurseries, or as movement corridors (Elliott et al., 2007; Elliott and Hemingway, 2008; Potter et al., 2015). These often shallow, productive waters also face intense disturbance due to anthropogenic factors and rapid climatic changes, posing risks for coastal residents and migratory species alike (Beck et al., 2001; Brown et al., 2018; Jackson et al., 2001).

* Corresponding author. NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems, PO Box 59, 1790 AB, Den Burg, Texel, the Netherlands.

E-mail addresses: jena.edwards@nioz.nl, edwardsj67@gmail.com (J.E. Edwards).

<https://doi.org/10.1016/j.ecss.2024.108737>

Received 7 September 2023; Received in revised form 2 March 2024; Accepted 23 March 2024

Available online 26 March 2024

0272-7714/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Marine conservation and management tools such as those applied in coastal waters and estuaries often rely on spatial approaches designed to protect critical habitats, provide refuge from human activity, and/or prevent overexploitation within a set of jurisdictional boundaries (Edgar et al., 2014). However, in open coastal ecosystems - characterised by high connectivity with adjacent fresh water, estuarine, or marine habitats - spatially-managed areas typically encompass only a portion of the total area used by migratory fish (Breen et al., 2015; Grüss et al., 2011; Palumbi, 2004). The efficacy of coastal management to protect migratory species is therefore influenced by their spatial and temporal distributions, as guided by individual movement behaviors (Elston et al., 2023; Frisk et al., 2019; Heupel and Simpfendorfer, 2005; Santana-Garcon et al., 2014; Speed et al., 2010).

1.2. Fish declines in the Wadden Sea

Spanning approximately 500 km of coastline across the Netherlands, Germany, and Denmark, the Wadden Sea is the world's largest remaining coastal wetland habitat is recognized as a UNESCO World Heritage Site for its "Outstanding Universal Value for present and future generations of all humanity" (Common Wadden Sea Secretariat, 2016; UNESCO, 2023). The region plays a vital role for numerous migratory fishes, including diadromous species, occasional visitors, and seasonal residents, all of which require access to multiple aquatic habitats to complete their life cycles. For these species, the Wadden Sea serves both as a movement corridor, facilitating access to adjacent freshwater or marine habitats (Jager et al., 2009; Tulp et al., 2017), while also supporting essential functions such as feeding, growth, and maturation (Hovenkamp, 1991; Rauck and Zijlstra, 1978; van der Veer et al., 2011, 2015, 2022).

Despite harboring approximately 189 fish species, representing a significant portion of the North Sea fish community, the Wadden Sea has witnessed shifts in community composition and alarming declines in overall fish abundance since the 1980s (Berg et al., 1996; Jager et al., 2009; Tulp et al., 2017; Tulp and Bolle, 2009; van der Veer et al., 2011, 2015) (Fig. 1).

While the exact causes of recent fish declines remain unclear, historical changes in the Wadden Sea fish community have likely stemmed from a multitude of anthropogenic and environmental shifts across the broader North Sea region (Tulp et al., 2008; van der Veer et al., 2015). Rising temperatures, including an increase of 1.5 °C over the last 25 years in the Dutch Wadden Sea (van Aken, 2008), have led to altered fish movement patterns, resulting in northward range shifts in marine fishes and potential delays in fish migration (Perry et al., 2005; van Walraven et al., 2017). Simultaneous threats such as coastal habitat destruction, exploitation, and increased predation have also influenced the Wadden Sea's suitability for fish, contributing to the local extinctions of a




Fig. 1. | Historical catch per unit effort (CPUE) for all fish species captured by the NIOZ fyke net survey between the years 1980 and 2021. Line colours are used to distinguish spring and fall catching periods. For a detailed description of methods see: (van der Meer et al., 1995). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

number of diadromous species (e.g., *Acipenser sturio*, *Salmo salar*, *Coregonus oxyrinchus*, and *Salmo trutta*) (Brasseur et al., 2015; Camphuyzen, 2004; Lotze, 2005, 2007) and degradation of the nursery function for marine juveniles (van der Veer et al., 2011, 2015). Additionally, for migratory fish, persistent barriers to migration success include the obstruction of freshwater routes to the Wadden Sea and destructive practices such as bottom-trawling and sand nourishment in coastal waters (Leewis et al., 2012; Tulp et al., 2020; van der Veer et al., 2015). These multifaceted impacts underscore the broader scale of factors affecting fish communities beyond the Wadden Sea, emphasizing the necessity for comprehensive conservation efforts across larger ecosystems.

1.3. Conservation needs

Due to the complex array of factors influencing recent fish declines and the effect of continuously shifting baselines (van der Veer et al., 2015), disentangling the specific causes poses a challenge. This lack of understanding extends to how fish interact with both human activities and the Wadden Sea environment at the individual level, hindering the operationalization of key management and conservation objectives outlined in the 2009 Quality Status Report for the Wadden Sea, known as the five Trilateral Fish Targets (Jager et al., 2009; SWIMWAY, 2019) (Box 1). While local management actions are typically confined to the Wadden Sea, it is important to recognize that the mobile fish they aim to protect exhibit much broader migratory behaviours, resulting in interactions with various environmental, anthropogenic, and ecological factors. Effective implementation and evaluation of spatial protection measures, such as those currently in place in the Wadden Sea within the Natura2000 network and UNESCO World Heritage List (Sieben et al., 2013), rely on knowledge of the spatial characteristics of fish movements both within the Wadden Sea and across adjacent habitats. Scientific evidence elucidating the key processes driving population dynamics and the identification of bottlenecks along the swimways of Wadden Sea fishes are essential for making the Fish Targets actionable and quantifying the success of proposed conservation measures.

1.4. A movement ecology approach

Aquatic telemetry is a field in which electronic tracking devices are affixed, either externally or via internal implantation, to free-ranging organisms to study their *in situ* movements across various spatial and temporal scales (Harcourt et al., 2019; Hussey et al., 2015; Watanabe and Papastamatiou, 2023). These movement data are valuable for understanding the fundamental drivers of fish movement (Elston et al., 2022; Rous et al., 2017; Sims et al., 2006) and have revealed important details of fish home ranges (Pereira et al., 2017), aggregation sites (Daly et al., 2019), and regional connectivity (Corrigan et al., 2018), and illustrated the probability of encountering threats or disturbances (James et al., 2005; Orrell et al., 2023). These insights have proven invaluable for wildlife and fisheries management thus far, guiding approaches for individual species and populations, and informing the delineation of fisheries stocks and protected areas (Brooks et al., 2019; Brownscombe et al., 2022; Crossin et al., 2017; Nathan et al., 2022). As such, telemetry studies are increasingly used to inform adaptive management strategies for fish and fish habitats (Allen and Singh, 2016; Cooke et al., 2016; Crossin et al., 2017; Lennox et al., 2019; Matley et al., 2022a).

1.5. Applying movement ecology for fish conservation in the Dutch Wadden Sea

To understand the causes and consequences of local behaviors and migration in fish, individual movements must be tracked and characterized over relevant spatial and temporal scales. Specifically, the scales of fish movements, degrees of habitat connectivity, and frequency of

Box 1

Five main management and conservation objectives (Trilateral Fish Targets; Jager et al., 2009; as worded by SWIMWAY, 2019):

1. Viable stocks of populations and a natural reproduction of typical Wadden Sea fish species;
2. Occurrence and abundance of fish species according to the natural dynamics in (a)biotic conditions;
3. Favourable living conditions for endangered fish species;
4. Maintenance of the diversity of natural habitats to provide substratum for spawning and nursery functions for juvenile fish;
5. Maintaining and restoring the possibilities for the passage of migrating fish between the Wadden Sea and inland waters.

transboundary movements all influence population success and must be taken into consideration for effective management at local, regional, and oceanic scales (Barkley et al., 2019; Cooke et al., 2023; Gillanders et al., 2003). Using the Dutch Wadden Sea as a case study, we hereby explore the implementation of a multi-scale telemetry approach to acquire fundamental knowledge and inform the conservation of migratory fish in open coastal ecosystems. In doing so, we address general knowledge gaps hindering effective migratory fish management by presenting questions related to large-scale movements, temporal aspects, and fine-scale behaviors, addressing them with the primary telemetry technologies available for marine environments (Box 2).

Our approach begins with an overview of the general migratory strategies observed in the Wadden Sea environment (section 2.1) and an introduction to the primary telemetry approaches available for their study (section 2.2). Subsequently, we use a reference transmitter in combination with a size-based approach to assess the potential suitability of all common and fairly common Wadden Sea fish species for long-term tracking (sections 2.3 & 2.4). Drawing from the selected species and available techniques, we then demonstrate the application of a multi-scale telemetry approach using four example species with differing life history traits and migration strategies (section 2.5). Based on these species-specific details, we propose a combination of telemetry approaches, incorporating novel data where available, to address our original research questions. Finally, we conclude with a discussion of our proposed methods, proposing future directions for implementing long-term tracking approaches in the Wadden Sea, and offer perspectives on how these lessons can be applied in other coastal ecosystems.

2. Application of a movement ecology approach for the wadden sea

2.1. Migration strategies in open coastal ecosystems

In open coastal environments such as the Wadden Sea, the local and broad-scale movements of migratory fish can occur across three distinct aquatic habitats, simplified here as the offshore marine, coastal, and freshwater realms (Fig. 2). For example, coastal migrants can exhibit wide-ranging movements in coastal or estuarine waters (e.g. *Salmo trutta*) (Degerman et al., 2012). Euryhaline species may move back and forth between marine or brackish coastal waters and freshwater habitats (e.g., *Coregonus oxyrinchus*) (Jensen et al., 2018), while marine fishes may move between coastal areas and deeper pelagic zones (e.g., *Raja clavata*) (Hunter et al., 2005; P. Walker et al., 1997). For diadromous species with large-scale migrations, (e.g., *Anguilla anguilla*) (Verhelst et al., 2018) coastal zones like the Wadden Sea are likely used primarily as movement corridors that link spatially discrete spawning grounds and adult foraging habitats. By generalizing the movement patterns of potential study species, we can help to identify specific research questions that can, in turn, be used to inform spatial management approaches (Box 2). However, to address these questions at the relevant spatial and temporal scales, each question requires its own suitable telemetry technique or combination thereof.

2.2. Telemetry approaches for fish conservation

Aquatic movement ecology is dominated by three categories of electronic devices: acoustic and satellite transmitters, and archival (data storage) tags (Hazen et al., 2012; Matley et al., 2022). While additional methods such as radio telemetry and passive integrated transponders (PIT) remain common in fresh waters, the proximities required for

Box 2

Movement ecology themes and potential research questions:

1. Temporal aspects of movement
 - a. What are the potential long-term (seasonal, annual) environmental/ecological drivers of fish presence?
 - b. Is the timing of arrival/departure influenced by external cues?
 - c. Do individuals exhibit site fidelity to specific regions/locations and, if so, at what temporal scale?
2. Large-scale movements
 - a. Do adult fish move cyclically between distinct habitats on a predictable (seasonal, annual, interannual) basis?
 - b. Do specific individuals, populations, or species make use of unique migratory pathways?
 - c. Do individual fish movements cross jurisdictional boundaries with varying levels of protection?
3. Fine-scale behaviours and habitat use
 - a. How do local environmental/ecological conditions influence the fine-scale movements and habitat use of individual fish?
 - b. Do individuals show site-specific resident or transient behaviours?
 - c. Are fine-scale individual behaviours consistent over different temporal scales?




Fig. 2. | Generalised movements exhibited by migratory fish species that inhabit (A) coastal waters, and (B) more specifically, in the context of the Dutch Wadden Sea. Arrows depict movement patterns characterised for: 1) coastal/marine migrants, 2) euryhaline species, 3) diadromous species, and 4) coastal/estuarine residents.

detection and ineffectiveness in salt water prohibits their use in coastal environments (Lennox et al., 2017). Despite rapid advancements in this field, technological and logistical constraints continue to affect the spatial and temporal scales addressed by telemetry studies (Harcourt et al., 2019). Tag size, lifespan, and the resolution and recovery of acquired data are all important considerations for selecting appropriate techniques for specific study questions and species. For each of the three dominant tag types (acoustic, satellite, archival), a brief summary of these factors is provided in the following subsections. With these details in mind, we then consider how a combination of these tagging approaches can be used to study the movements of migratory fish both within coastal habitats, and over broader spatial and temporal scales.

2.2.1. Acoustic telemetry

Acoustic telemetry uses autonomous receivers to detect and decode unique acoustic signals emitted by animal-borne transmitters, enabling tracking of fish movements in both freshwater and marine environments (Donaldson et al., 2014; Heupel and Webber, 2012; Matley et al., 2022b). Acoustic tag lifespans are dictated by transmission settings and battery size, with examples ranging from 100 days for miniature transmitters suitable for juvenile fish (Lingard et al., 2023) and the largest tags lasting for periods of up to 10 years (Edwards et al., 2022a). However, long-term tracking tagged fish requires the maintenance of acoustic receivers over equivalent time periods. Using a network of receivers, known as an acoustic array, researchers can analyse the timing of animal detections to understand fish presence in relation to temporal cues such as diurnal, seasonal, or interannual variations in environmental conditions (Box 1). Different array configurations can address research questions at various spatial and temporal scales (Heupel et al., 2006). Acoustic positioning methods, employing synchronization tags and positioning models (e.g., using time-difference-of-arrival or time-of-arrival) allow fine-scale positioning in localized areas (Fig. 3d), but require dense arrays of close-proximity receivers and prior knowledge of fine-scale residency by target species (Baktoft et al., 2017; Lennox et al., 2023a; Orrell and Hussey, 2022; van der Knaap et al., 2020). In contrast, coarser detection data from receiver gates can indicate fish occurrence, mortality rates, and passage success at key locations along migratory pathways (Fig. 3e) (Chaput et al., 2019; Laroque et al., 2020; Verhelst et al., 2018). Meanwhile other forms of presence-absence arrays (e.g., grids) can effectively illustrate habitat selection, coastal residency, and site fidelity over broader, defined study areas (Fig. 3c) (Able et al., 2014; Novak et al., 2020; Reyier et al., 2023). In each of these approaches, acoustic detections confirm the presence of an individual within the detection range of a receiver, whereas accurate estimates of animal positions remain unknown. Detection ranges are

also highly variable over time and space due to the influence of a variety of factors on the transmission and detection of acoustic signals (Kessel et al., 2014). At the largest spatial scales, collaborative telemetry networks, such as the Ocean Tracking Network (OTN) (Iverson et al., 2019), the European Tracking Network (ETN) (Abecasis et al., 2018), the Florida Telemetry network (FACT) (Young et al., 2020), the Integrated Marine Observing System's Animal Tracking Facility in Australia (IMOS ATF) (Hoenner et al., 2018), extend the reach of acoustic studies to regional and international ranges by linking independently managed arrays and facilitating data sharing and collaboration (Fig. 3b) (Abecasis et al., 2018).

2.2.2. Satellite and archival tags

Satellite transmitters (SAT tags) use platforms such as GPS (Global Positioning System, e.g., GPS and Fasloc GPS tags) and ARGOS (see <https://www.argos-system.org/>) to remotely transmit stored data and animal positions over vast spatial scales, but are suitable only for animals that spend considerable time at the surface such as marine mammals, sharks, and marine reptiles (Hussey et al., 2015). Tag lifespans are dictated by battery life and attachment success, but are typically programmed for periods of >1 year (Hammerschlag et al., 2011), limiting their application to study migration cycles or repeated movements over longer timespans. By comparison, archival tags – also known as data storage tags or bio-loggers – store information on environmental conditions (e.g., temperature, pressure, light levels) and/or the host's animal's physiological, or biological status (e.g., heart rate, acceleration) to onboard memory, and are ideal for studying smaller, non-surfacing species (Harcourt et al., 2019). Without the energy costs of data transmission, archival tags can be deployed for multi-year periods, but must be physically recovered to access stored data, prompting their preferential use for species with a high likelihood of recapture (e.g., commercially-targeted species or those with predictable distribution patterns) (Watanabe and Papastamatiou, 2023). PSATs (pop-up satellite archival tags) are an exception, wherein summarised or binned subsets of stored data - including depth, temperature, and or light levels - can be transmitted to satellites, along with a GPS position of the tag's location at the surface following pre-programmed release from the host animal (Block et al., 1998; Skomal et al., 2009). Due to difficulties related to tag attachment and premature release from the host animal, PSATs tend to have short deployment periods, typically limited to <1 year (Hammerschlag et al., 2011; Musyl et al., 2011). While lacking continuous positional data, both archival tags and PSATs (if recovered) store high-frequency environmental data useful for revealing individual environmental preferences and retroactively modeling movement trajectories using methods like hidden Markov models (HMMs) (Braun




Fig. 3. | Spatiotemporal scales and potential applications of acoustic telemetry systems in coastal and offshore marine habitats exemplified for the Dutch Wadden Sea. Panels depict a variety of research techniques available using acoustic telemetry devices, including: A) the selection of receiver attachment locations from available structures (e.g., Rijkswaterstaat navigational buoys – <https://www.rijkswaterstaat.nl>), B) the incorporation of independently-managed acoustic arrays into an international telemetry network (e.g., the European Tracking Network – <https://www.europeantrackingnetwork.org>), where coloured circles each represent acoustic arrays comprised of multiple receiver stations, C) a presence-absence array design showing nearby receivers with non-overlapping detection ranges (where R indicates detection range), D) a fine-scale acoustic positioning array using receivers in close proximity to track fish trajectories, and E) acoustic receivers in gate/curtain formation. Panel captions describe the temporal scale (TS) and spatial scale (SS) of the illustrated telemetry method (M) and list corresponding research questions (Q) as listed in Box 2.

et al., 2018; Thygesen et al., 2009).

2.2.3. Combined approaches for multi-scale tracking

In coastal systems and estuaries, telemetry studies have historically taken place in geographically-constrained locations such as bays, fjords, or lagoons (Grothues, 2009). Access to study animals and receiver deployment sites, alongside a desire to maximize detection or recapture frequencies, have likely contributed to such decisions, but may preclude the consideration of other factors important for migratory fish populations, such as broad-scale connectivity. Conversely, telemetry techniques equipped to study large-scale animal movements may be limited in temporal or spatial resolution, with the use of some tag types further restricted by prohibitive lifespans or sizes. For migratory fish inhabiting open coastal systems, the combination of multiple telemetry approaches will be necessary to capture the complete range of movement behaviours exhibited across various life stages and spatial scales.

Recent advancements in tracking technologies have introduced new capabilities, such as the incorporation of a variety of sensors into acoustic tags to record and, in some cases, transmit information such as

animal acceleration, environmental conditions, and predation events (Goossens et al., 2023; Klinard et al., 2019; Lennox et al., 2023b). These innovations are expanding research opportunities to investigate various factors influencing individual fish behavior. Additionally, double-tagging, which involves the attachment of two tags to one individual (e.g., acoustic transmitters and bio-loggers), increases the breadth of data acquired for each tagged animal. While combination tags and double-tagging can increase tag burden and typically require larger individuals, the separate deployment of standard acoustic and archival tags on individuals within a population can also provide valuable insights, particularly for species frequenting both coastal and offshore waters. These complementary approaches are particularly useful in open coastal systems where acoustic receiver coverage is typically distributed close to shore, whereas environmental reference fields and archival tag data can be used to reconstruct movements in more sparsely covered offshore regions.

Modern telemetry techniques can be integrated with conventional methods, such as biological sampling and mark-recapture, to provide a comprehensive understanding of the causes and consequences of fish movement at both individual and population levels. While conventional capture-mark-recapture studies provide insights into movement behaviours at coarse spatial and temporal scales (Queiroz et al., 2005), the modelling approaches they employ can also be used in combination with electronic tagging to estimate demographic parameters crucial for fisheries management, including survival rates, abundance estimates, and scales of movement (Bacheler et al., 2009; Brooks et al., 2019; Dudgeon et al., 2015; Lees et al., 2021). Furthermore, the integration of telemetry data and non-lethal biological samples collected during tagging (e.g., fin clips, scales, blood) offers additional insights into eco-physiological factors such as ageing, genetics, stable isotopes, microchemistry, and hormone levels (Borcherding et al., 2008; Biton-porsmoguer, 2022; Brownscombe et al., 2022b; Matley et al., 2023). These aggregated datasets contribute to a deeper understanding of fish diets, physiological status, migration histories, and population structure.

2.3. Criteria for reference tag selection and species suitability estimates

In the face of multi-species declines, such as those observed in the Wadden Sea, many factors may be used to determine the suitability of a species for tagging. To demonstrate a potential size-based approach, we examined 76 potential study species whose frequency of occurrence in the Wadden Sea was classified as common or fairly common (Jager et al., 2009). To maintain our focus on large-scale connectivity, one strictly freshwater species (*Abramis brama*) was excluded from this list. Scientific names were listed as referred to by (Jager et al., 2009) and common names as they appear in FishBase (Froese and Pauly, 2023).

As a simplified approach, we then used the 2% body weight rule (Jepsen et al., 2005) and the weight of a standard reference tag to estimate the minimum fish body size required for multi-year tracking. Several considerations were used to select a reference tag that would both minimize tag burden on target individuals, while maximizing important factors such as detection range, equipment compatibility, and study duration. Due to the wide-scale adoption of acoustic telemetry, its potential for use across coastal, offshore, and freshwater environments, and the potential to achieve tag lifespans of multiple years, we chose an acoustic transmitter as our reference tag. As a hypothetical approximation of the minimum tag size required to achieve sufficient detection ranges for open coastal areas, equipment compatibility, and study durations of >1 y, we selected a 9 mm 69 kHz acoustic transmitter weighing approximately 3.6 g in air. As many commercially-available DSTs are of comparable size when used without a float, the species suitability estimates listed in Table 1 and Appendix I are deemed applicable to both tag types.

Based on our chosen reference tag, the minimum body weight for fish tagging was 180 g. However, as individual fish weights are often unavailable at capture, we used biological information from FishBase

Table 1

Common and fairly common marine Wadden Sea fish species considered to reach suitable (S) or potentially suitable (P) sizes for multi-year tracking with internally implanted acoustic transmitters. Lengths are listed in cm for male/unsexed individuals (M/U) and correspond to total length (TL), standard length (SL), fork length (FL), or disc width (WD). IUCN Red List status includes: Not Evaluated (NE), Data Deficient (DD), Least Concern (LC), Near Threatened (NT), Vulnerable (VU), Endangered (EN), and Critically Endangered (CR). For species whose abundance trends for the Dutch Wadden Sea (DWS) have been assessed, recent trends were either stable, decreasing, or uncertain. Wadden Sea guilds are listed as diadromous (D), marine juvenile (MJ), or estuarine resident (ER). Species (potentially) large enough to support tagging with PSATs are denoted by an asterisk.

Scientific name	English name	Length at maturity (cm)	Max length (M/U) (cm)	Length unit	Max published weight (kg)	Max reported age (y)	IUCN Red List Status	DWS trend (2011–2020)	WS guild	Suitable for telemetry
<i>Alosa alosa</i>	Allis shad	47.8	69	TL	4	10	LC			S*
<i>Alosa fallax</i>	Twaite shad	32.5	60	TL	1.5	25	LC	Decreasing	D	S
<i>Anguilla anguilla</i>	European eel	55	122	TL	6.6	23	CR	Uncertain/ Decreasing ^a	D	S*
<i>Belone belone</i>	Garfish	U	104	TL	1.4		LC			P
<i>Chelon labrosus</i>	Thicklip grey mullet	29.5	75	SL	4.5	25	LC			S
<i>Cyclopterus lumpus</i>	Lumpsucker	U	61	TL	9.5	13				P
<i>Dasyatis pastinaca</i>	Common stingray	U	69.5	WD	10.2		VU			P
<i>Dicentrarchus labrax</i>	European sea bass	36.1	103	TL	12	30	LC	Decreasing	MJ	S
<i>Gadus morhua</i>	Atlantic cod	65.4	200	TL	96	25	VU	Uncertain	MJ	S*
<i>Gaidropsarus vulgaris</i>	Three-bearded rockling	U	60	TL			LC			P
<i>Galeorhinus galeus</i>	Tope shark	144.1	193	TL	44.7	55	CR			S*
<i>Glyptocephalus cynoglossus</i>	Witch flounder	30.4	60	TL	2.5	25	VU			S
<i>Hippoglossoides platessoides</i>	American plaice	35	82.6	TL	6.4	30	EN			S
<i>Lampetra fluviatilis</i>	River lamprey	34.8	50	TL	0.15	10	LC	Stable	D	S
<i>Merlangius merlangus</i>	Whiting	28.2	91.5	TL	3.1	20	LC	Stable	MJ	S
<i>Merluccius merluccius</i>	European hake	42.3	140	TL	15	20	LC			S
<i>Microstomus kitt</i>	Lemon sole	29.3	65	TL	3	23	LC			S
<i>Petromyzon marinus</i>	Sea lamprey	U	120	TL	2.5	11	LC	Decreasing	D	P
<i>Pleuronectes platessa</i>	European plaice	32.7	100	SL	7	50	LC	Stable	MJ	S
<i>Pollachius pollachius</i>	Pollack	41	130	TL	18.1	8	LC			S
<i>Pollachius virens</i>	Saithe	39.1	130	TL	32	25	NE			S
<i>Psetta maxima</i>	Turbot	40.8	100	SL	25	25	LC			S
<i>Salmo salar</i>	Atlantic salmon	73.1	150	TL	46.8	13	NT		D	S*
<i>Salmo trutta</i>	Sea trout	40.9	140	SL	50	38	LC	Decreasing	D	S
<i>Scomber scombrus</i>	Atlantic mackerel	28.7	60	FL	3.4	17	LC			S
<i>Scophthalmus rhombus</i>	Brill	37	75	TL	8	6	LC			S
<i>Solea solea</i>	Common sole	30.3	70	SL	3	26	DD	Stable	MJ	S
<i>Stizostedion lucioperca</i>	Pike-perch	37.2	100	SL	20	17	LC			S

^a Decreasing trend for juveniles.

(Froese and Pauly, 2023) to convert from a weight-based to a size-based evaluation. Using the length-weight relationship of European sea bass as an example (Froese and Pauly, 2023), we set our length criteria for tagging suitability to a minimum total length of 27 cm. To maintain consistency across species, we then used length at maturity (L_m) as a reference for this assessment. Length data for each fish species was obtained from FishBase (Froese and Pauly, 2023) and are listed in Appendix I. For those species where L_m was not available, maximum length was used as the reference value with the minimum criteria being 60 cm (based on the maximum lengths of all other species determined to be suitable). However, due to the level of uncertainty surrounding these length data, species for which the maximum length was used were deemed 'potentially suitable' for tracking. To aid in assessing the priority of potential Wadden Sea species for study, population trends for the Dutch Wadden Sea over the period of 2011–2020 are included in the table, as well as the guild assigned to each species (Tulp et al., 2021).

To determine which species have the potential for study with larger

PSAT tags, a minimum length at maturity of ≥ 47 cm was selected (Table 1). This value was chosen as a cautious estimate as per the results of (Lynch et al., 2017) which found no effect of PSAT tagging on the metabolic rates and swimming kinematics of juvenile sandbar sharks within this size range. It should be noted that a variety of factors may influence the effect of tag burden on short-term swimming performance and long-distance migration (e.g., tag shape and buoyancy as well as the swimming kinematics of the tagged individual) and are, for many species, not well understood (Lynch et al., 2017; Svendsen et al., 2021). For a thorough understanding of species-specific tag effects, similar results must be obtained via additional targeted studies.

2.4. Potential target species for long-term tracking

Using our size-based assessment, 28 (fairly) common Wadden Sea fish species were deemed suitable (23 species) or potentially suitable (5 species) for multi-year telemetry studies (Table 1). Only 5 species were

of sufficient size (or had sufficient size information available) to be deemed suitable for tagging with PSATs. This included 3 species for which dedicated assessments of the effects of satellite tagging on migratory behaviours and swimming performance have previously been conducted (Atlantic salmon, European eel, and Atlantic cod) (Hedger et al., 2017; Methling et al., 2011; Nielsen et al., 2023; Økland et al., 2013). The majority of species included were classified by the IUCN as being of Least Concern (18 species), while 6 species are classified as Vulnerable (common stingray, Atlantic cod, witch flounder), Endangered (American plaice), or Critically Endangered (tope shark and European eel). Despite their common occurrence in the Dutch Wadden Sea, information on local population status was available for only 10 species in total, of which less than half were classified as stable. Maximum reported ages of these fish also indicated long life spans often exceeding 20 years (14 species) and even reaching 50 years for the tope shark and European plaice.

While this size-based assessment was used to narrow down the list of potential candidate species, additional factors like catchability, vulnerability to capture-induced stress, and protective measures for threatened species are also vital considerations for fish tracking studies. Additionally, local and broadscale population trends, threats, and life history traits affecting population resilience (e.g., long lifespans, low fecundity) are all useful for determining the necessity of tracking studies. Given the potential for long-lived species to exhibit migration cycles extending over multiple years, animal longevity should also be considered in terms of tag selection and programming to prevent mismatches in temporal scale between tracking studies and target behaviours (Edwards et al., 2022b). Furthermore, additional information on species-specific size distributions and capacity for tagging could result in additions to this list. Reductions in tag size and advances in tagging methodologies will

also continue to improve survival rates and reduce tag burden in the future, with benefits for animal welfare and data collection (Lennox et al., 2017).

2.5. Applying the approach to four example species

Knowledge of a species' life cycle can guide research in movement ecology, particularly for migratory fish that move between various habitats at different life stages. To illustrate the application of a life cycle approach for movement ecology, we selected four fish species from Table 1 that differ in their population trends (Fig. 4) and migration strategies (Fig. 5), as well as in our level of knowledge of each. These example species are: thicklip (or thick-lipped) grey mullet (*Chelon labrosus*), European sea bass (*Dicentrarchus labrax*), sea trout (*Salmo trutta*), and tope shark (*Galeorhinus galeus*). For each of these four species, sufficient knowledge is available to make generalised predictions of where individuals are likely to occur (i.e., in the Wadden Sea or in adjacent freshwater or marine habitats) on a seasonal basis (Fig. 5, Table 2). Furthermore, movement data from the Wadden Sea or adjacent waters are available for each of these four species, highlighting the various telemetry approaches described in this study. Rather than acting as representatives of broader ecological groups, these species were selected as examples with which to demonstrate how species-specific knowledge of fish life histories can be used to inform research questions and telemetry approaches. It should be noted that the multi-year periodicity of migratory behaviours of these species within the described habitats is still poorly understood. Whether migrations between these regions occur annually or interannually, and whether these temporal aspects differ among individuals, are questions that can still be addressed using the telemetry techniques discussed in the following sections. To




Fig. 4. | Historical catch per unit effort (CPUE) and total catches (n) of four large fish species from the NIOZ fyke net survey, which are used in the examples: A) thicklip grey mullet, B) European sea bass, C) sea trout, and D) tope shark. For a detailed description of fishing methodology see: (van der Meer et al., 1995).




Fig. 5. | Generalised movements/migration strategies exhibited by four migratory fish species within and between offshore marine, coastal (e.g., the Wadden Sea), and freshwater habitats. A) thicklip grey mullet, B) European sea bass, C) sea trout, and D) tote shark. The size of the fish indicates the life stage and eggs are shown as white circles.

Table 2

| Current knowledge of broad-scale movement behaviours seasonal distributions of adult individuals from four migratory fish species that use the Wadden Sea. Life history information from FishBase (Froese and Pauly, 2023) was used for all species, along with the listed references.

Species	Winter	Spring	Summer	Autumn	Refs
<i>C. labrosus</i>	Spawning, overwintering Offshore; location unknown, timing unknown	Migrating Offshore → Wadden Sea	Foraging Coastal waters; duration uncertain, spatial scale uncertain	Migrating Wadden Sea → Offshore	(Cardona, 2006; Hickling, 1970; Kennedy and Fitzmaurice, 1969, 1972; Walsh et al., 1994)
<i>D. labrax</i>	Spawning, overwintering Offshore; location unknown, timing unknown	Migrating Offshore → Wadden Sea	Foraging Coastal waters; duration uncertain, spatial scale uncertain	Migrating Wadden Sea → Offshore	(de Pontual et al., 2019; Doyle et al., 2017; Holden and Williams, 1974; Kennedy and Fitzmaurice, 1972; Pawson et al., 1987, 2007)
<i>S. trutta</i>	Spawning, overwintering Fresh water	Migrating Fresh water → Wadden Sea → Offshore	Foraging Coastal waters; duration uncertain, spatial scale uncertain	Migrating Offshore → Wadden Sea → Fresh water	(Aldvén et al., 2015; Bendall et al., 2005; Bij De Vaate et al., 2003; Eldøy et al., 2015; Flaten et al., 2016; Jensen et al., 2018; Thorstad et al., 2016)
<i>G. galeus</i>	Unknown Presumed offshore; scale of movements unknown	Unknown Presumed offshore; scale of movements unknown	Pupping Coastal waters; precise timing and locations unknown	Unknown Presumed offshore; scale of movements unknown	(Capapé et al., 2005; Colloca et al., 2019; Fitzmaurice et al., 2003; Holden and Horrod, 1979; Schaber et al., 2022; Stevens, 1990; Thorburn et al., 2019)

understand the complete life cycles of these and other Wadden Sea fish species, and to unravel the potential causes of local declines in this region, it will be necessary to increase the scope of movement studies to

ecologically-relevant scales by combining available telemetry techniques and complementary data sources.

2.5.1. Sea bass and thicklip mullet – seasonal residents

2.5.1.1. Knowledge gaps hindering management. As adults, both European sea bass and thicklip grey mullet are known as seasonal visitors to the Wadden Sea (Boer, 1971; Cardoso et al., 2015), with juveniles and adults of both species migrating toward coastal waters in spring and adults departing in autumn to spawn and overwinter in offshore waters (Fig. 5a and b) (Hickling, 1970; Pawson et al., 1987). Despite their similar life histories (Fig. 5a and b), historical survey data suggest that sea bass and thicklip mullet populations in the Dutch Wadden Sea have experienced contrasting trends throughout the latter part of the 20th century (Fig. 4a and b). While the cause for this disparity remains unclear, examining differences in the ecological roles of these two seasonal residents may provide insight.

Sea bass and thicklip mullet rely on a distinct set of environmental conditions, including preferred temperatures, substrates, and feeding ecology (Poiesz et al., 2020, 2023). This suggests that local and regional environmental conditions might impact the growth and survival of each species differently, potentially influencing their population trends. However, the precise conditions that individuals encounter and their behavioural responses require further study. For thicklip mullet in particular, whether individuals move between marine and freshwater habitats or represent two distinct populations remains completely unknown (Fig. 5), but could have important implications for population stability. Similarly, anecdotal evidence suggests that a portion of the local sea bass population may remain resident in Dutch coastal waters year-round, however, this has yet to be confirmed by the scientific literature.

For both species, the influence of Wadden Sea conditions (e.g., temperature, substrate type, food availability) and species-specific biology (e.g., thermal and salinity tolerance, diet) on migration timing, residency duration, and home range size are still unknown, but could indicate the suitability of Wadden Sea habitats for fish from distinct ecological guilds, informing local conservation measures. Additionally, little is known about the destinations and migration routes of individuals leaving the Wadden Sea (e.g., spawning or overwintering habitats), obscuring the role of offshore conditions and threats on reproductive success and survival. By addressing these uncertainties, conservation efforts can better target specific threats occurring at both local and regional scales.

2.5.1.2. Local and large-scale tracking with acoustic and archival telemetry. Within the boundaries of the Dutch Wadden Sea, acoustic telemetry offers a means to investigate the nature and drivers of local behaviours and migration timing for sea bass and thicklip mullet. As seasonal residents, individuals can be targeted and tagged during the summer feeding period when they are known to be locally present. Presence/absence data from coastal acoustic receivers – such as those in the local Swimway array (<https://swimway.nl/hoef-gebruiken-grote-vissen-de-waddenzee/>) – can then be used to analyse their space use in relation to potential environmental influences such as depth, temperature, and food availability (Fig. 6). These correlations can provide valuable insights for ecological models that predict how long-term environmental changes might affect the distribution and success of fish populations under future climate and fisheries management scenarios.

Moreover, movement trajectories and departure of tagged fish can be determined by the sequentially arranging individual fish detections within the acoustic network (Fig. 6). These trajectories help define the scales of space use, regional connectivity, and high activity areas at both individual and population levels. Individual movement data can rapidly reveal behavioural responses to changing environmental conditions (Breece et al., 2018; Brownscombe et al., 2022; Lédée et al., 2015), indicating the abilities of sea bass and thicklip mullet to use behavioural flexibility as an adaptive mechanism in the face of climate change.




Fig. 6. | Local movement trajectory of an individual thicklip grey mullet (animal ID: TKM21) tagged and tracked via acoustic telemetry in the western Dutch Wadden Sea. Hollow circles indicate the locations of all acoustic receiver stations in the Swimway array where the tagged individual was not detected. Filled circles are colour by date from June 24, 2021 to July 29, 2021 and connected by interpolated shortest-distance paths. Wadden Sea bathymetry data are publicly available via: (Baptist et al., 2019); DOI:10.17632/27-mysx289g.1. Depth codes are classified as follows: deep sublittoral (depth < -5 m mean low water spring tide), shallow sublittoral (depth \geq -5 m mean low water spring tide, <4% mean exposure), low littoral (4% \leq mean exposure <25%), middle littoral (25% \leq mean exposure <75%), high littoral (75% \leq mean exposure <85%), supralittoral (mean exposure \geq 85%), salt marsh (vegetated).

Using a combination of acoustic detections from the European Tracking Network and archival data from DSTs, the destinations and migration pathways of sea bass and thicklip mullets can also be addressed. Fish can either be tagged with a single acoustic tag, a single DST or ADST, or - provided that an individual is of adequate size (estimated for sea bass to be \geq 50 cm) – internally double-tagged with both an acoustic transmitter and DST. For acoustic-tagged fish that depart the Dutch Wadden Sea (i.e., the Swimway array), data for individuals detected by acoustic arrays in neighbouring regions (e.g., German or Danish Wadden Sea regions to the east, or Belgian waters and the English Channel to the west) can be uploaded to the European Tracking Network by their managing researchers. As a member of the ETN, the tag owner can then access and offload these data to uncover the presence of these tagged fish across a much broader study area (Fig. 7). As knowledge of the distributions of most seasonally migrating Wadden Sea species is limited to only parts of the year when they are locally resident, these data could reveal connectivity between discretely managed stocks as well as highlighting offshore aggregation sites potentially used for spawning, foraging, or overwintering.

2.5.2. Sea trout – A diadromous migrant

2.5.2.1. Knowledge gaps hindering management. Sea trout are iteroparous, returning to freshwater habitats to spawn multiple times throughout their adult life and exhibiting an array of movement behaviours spanning three habitat types (Fig. 5c). However, reductions in habitat connectivity and the subsequent implications for migrant mortality could jeopardize the stability of local sea trout populations (van Puijenbroek et al., 2019; Wright et al., 2014). Although the Wadden Sea likely acts as a movement corridor or temporary foraging ground for sea trout, the existence and location of common migratory pathways through the system remain unclear. This information is valuable for prioritising or improving fish passages for enhanced migration speed and survival. For instance, in Dutch waters, sea trout catches have




Fig. 7. | Large scale movement trajectories outside the Wadden Sea for two adult sea bass (animal IDs: 19840, 19872) tagged with acoustic transmitters in the western Dutch Wadden Sea. Filled circles indicate locations of fish detections and are scaled by the number of detections recorded. Points are connected by interpolated shortest-distance paths which have been curved to prevent overlap. Colours indicate the date of detections ranging from July 9, 2021 to July 17, 2022 (ID: 19840) and July 21, 2021 to May 8, 2022 (ID: 19872), respectively. In the Wadden Sea, detections were recorded by receivers in the SWIMWAY array. Additional data from FISHINTEL (Sheehan et al., 2021) and BPNS (Reubens et al., 2017) receivers were sourced from the European Tracking Network data portal (<http://www.lifewatch.be/etn/>), developed by the Flanders Marine Institute as part of the Flemish contribution to LifeWatch. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

remained low over the past several decades (Fig. 4), and continued obstruction by the Afsluitdijk dam limits the passage of sea trout entering the Rhine basin from the western Wadden Sea (Bij de Vaate et al., 2003). In contrast, in Danish rivers surrounding the Wadden Sea, measures have been implemented to enhance connectivity for migrating trout, including net fishing restrictions and major river restoration efforts, resulting in positive outcomes such as improved and expanded wild salmonid habitat and increased juvenile production (Tulp et al., 2022).


Despite high mortality rates incurred both at-sea and during migration (Aarestrup et al., 2015), sea trout movements both within and between marine habitats have received limited attention compared to the breadth of fresh water studies (Eldøy et al., 2015; Thorstad et al., 2016). As sea trout are known to use coastal habitats affected by anthropogenic activities, including aquaculture, renewable energy production, and other coastal infrastructure, there is potential for overlap in marine behaviours and a variety of possible threats (Eldøy et al., 2017). The lack of information on sea trout behaviours, survival, and residency in marine habitats are therefore major gaps in our knowledge of when and where population bottlenecks occur, hindering effective conservation and management.

2.5.2.2. Assessing connectivity and marine movements with acoustic and archival telemetry. As with the seasonal residents, acoustic telemetry can be used to monitor the movements of diadromous migrants like the sea trout through fresh, transitional, and coastal marine waters (Aldvén et al., 2015; Bendall et al., 2005; Finstad et al., 2005). Strategic placement of single acoustic receivers or receiver gates at marine-freshwater transition points can inform rates of passage on both seaward and return migrations, providing critical insight on migration success and the effectiveness of conservation measures promoting fish passage (Aldvén et al., 2015) (Fig. 2e). In restricted waterways and boundary waters, small acoustic tags with inherently limited transmission ranges can capture the first migrations of the smallest migratory life stages (e.g., post-smolts) (Flatén et al., 2016). In open coastal habitats, receiver gates and grid arrays (Fig. 2c–e) and longer-range transmitters can then be used to identify movement corridors and monitor individual residency to determine the role of coastal regions like the Wadden Sea for growth and survival (Aarestrup et al., 2015; Eldøy et al., 2015). Tags equipped with depth and/or temperature sensors can highlight individual preferences in relation to environmental conditions (Eldøy et al., 2017) or biological factors such as parasite infestation (Gjelland and Hedger, 2017).

For regions with limited coverage by acoustic arrays, including offshore marine habitats, archival data from DSTs can be used alongside hidden Markov models (HMMs) to reconstruct the large-scale movement pathways of adult fish (Braun et al., 2018; Pedersen, 2007; Thygesen et al., 2009). HMMs have already been employed to reconstruct the offshore migrations of two Wadden Sea fish species, the European sea bass and sea trout, which were tagged with data storage tags in coastal European waters (de Pontual et al., 2019; Kristensen et al., 2018, 2019) (Fig. 8). These data highlight the spatial and temporal extent of marine movements, as well as revealing behavioural shifts in response to elevated water temperatures as a means of optimizing growth at sea (Kristensen et al., 2018, 2019). For migrating sea trout in particular, further investigation of the balance between the metabolic costs and benefits of migration could prove useful for understanding the potential consequences of environmental change (Kristensen et al., 2019).

2.5.3. Tope shark – an occasional visitor

2.5.3.1. Knowledge gaps hindering management. Tope sharks are capable of conducting extensive migrations across deep pelagic waters in the northeast Atlantic, occasionally moving into the western Mediterranean (Colloca et al., 2019; Fitzmaurice et al., 2003; Holden and Horrod, 1979) or as far west as Iceland (Stevens, 1990; Thorburn et al., 2019). However, whether these broad-scale movements represent repeated migratory behaviours or occasional dispersion is still the subject of speculation (Colloca et al., 2019). Incidental catches of adult females and neonates from within the Wadden Sea and along the northern Dutch coastline suggest that pupping occurs in these areas in mid-summer (Batsleer et al., 2020) (Fig. 5d). However, the use of these coastal waters by gravid females and their potential nursery function for pups have yet to be determined. Due to a lack of directed angling for tope sharks and their absence from standard fish monitoring programs, their presence can easily remain undetected, leaving insufficient data to indicate local abundance (Fig. 4). Importantly, northeast Atlantic tope sharks have been deemed Critically Endangered by the IUCN's Red List (T. I. Walker et al., 2020), but are still vulnerable to capture by French and English fishing fleets in coastal waters of the English Channel (Biton-porsmo-guer, 2022). Given the scale of horizontal movements exhibited by this species, tope sharks may act as indicators of regional connectivity at an ocean basin scale. Furthermore, if specific biological needs are met by discrete locations throughout its vast range (e.g., pupping in shallow coastal waters), broad-scale connectivity between juvenile and adult habitats may be critical for their population success.



(Adapted from Kristensen et al., 2018; 2019)

Fig. 8. | Data types required for the reconstruction of fish trajectories from archival tags (*i.e.*, data storage tags, bio-loggers): A) Sea surface temperatures representing a single timepoint for the North Sea and Baltic Sea region: data from (Van der Molen et al., 2021), B) bathymetry of the North Sea and Baltic Sea region: data from (Perluka et al., 2006), C) From Kristensen et al. (2018): example of marine behaviour with depth (blue) and temperature (red) time series recorded by a data storage tag (DST) deployed on an adult sea trout, D) From: Kristensen et al. (2019): reconstructed track of the most probable movements of an adult sea trout (Fish 8). Line colour refers to the month of tracking as follows: April (green), May (blue), June (yellow), July (red), August (pink). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

2.5.3.2. Acoustic telemetry networks and PSATs to capture long-range movements. Acoustic telemetry can provide insight into the nursery function of coastal and estuarine habitats and indicate the efficacy of spatial protections such as Shark Refuge Areas and fisheries closures (McAllister et al., 2015). In place of catch data, long-term acoustic monitoring can provide estimates of juvenile mortality for improved population modeling as well as demonstrating population recovery (Heupel and Simpfendorfer, 2002; McAllister et al., 2015). Receiver gates positioned across the tidal inlets could also indicate the timing of emigration toward offshore waters for juveniles tagged in the Wadden Sea (Heupel, 2007; Wada et al., 2017). However, as juveniles begin emigrating from nursery areas as early as 1–2 y of age with low rates of return (McAllister et al., 2015), fixed coastal receiver arrays will likely capture only glimpses of adult behaviours. Gravid females have also been shown to demonstrate partial migration, with some individuals using offshore migration routes where they may bypass protected coastal areas and face higher fishing pressures (McMillan et al., 2019). Strategic use of acoustic networks can thereby help elucidate connectivity between protected and unprotected coastal and offshore waters for both juveniles and adults, allowing the evaluation and refinement of spatial management boundaries (Espinoza et al., 2021; McAllister et al., 2015).

Due to their large body size, adult tope sharks can also be equipped

with large devices such as PSATs that record high-resolution data for track reconstruction and examination of vertical movement behaviours (Schaber et al., 2022; Thorburn et al., 2019) (Table 1, Fig. 9). Similarly, this species is potentially suitable for double tagging with PSATs and long-lifespan acoustic transmitters, combining the benefits of short-term high-resolution data with the potential for opportunistic detections over longer time spans. Further evidence of large-scale regional connectivity in the north Atlantic could have implications for tope shark population structure and resilience, signifying a need for broad-scale cooperation among European fisheries managers to prevent further declines in this Critically Endangered species.

3. Synthesis

3.1. Summary of findings

The interconnected nature of open coastal systems means that fish tagged in these regions are not necessarily restricted to a defined study area. The timing and duration of fish presence in coastal waters, as well as the scale of local movements, environmental drivers, and habitat associations, can all influence the effectiveness of local management actions and are relevant for identifying potential causes of fish declines and informing conservation approaches. In this research, we considered




Fig. 9. | Data on the large-scale movements of four tope sharks (IDs: 153233, 168495, 168499, 168500) in the North Sea and Northeast Atlantic from (Schaber et al., 2022) (both panels). A) Most likely movement trajectories-geolocation from GPE3 state-space model between deployment start (S) and end (E) locations (black paths). Shaded zones around the most likely tracks indicate the 95% location probabilities and are colour-coded by tag ID/Ptt. B) Time series of tope shark depth measurements from four PSAT deployments with corresponding bathymetry along migration path. For complete details on both figures, please refer to the original manuscript. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

the importance of studying fish behaviours and developing management strategies at ecologically-relevant spatial and temporal scales. Given the variety of movement behaviours exhibited by coastal migrants, and the potential for threats and bottlenecks to be encountered at any point within a species' migratory range, we propose that movement studies will ultimately require a combination of telemetry approaches designed to match these behavioural scales to the data resolution provided by commercially available tags.

Using the Dutch Wadden Sea as an example study environment, we suggested a size-based approach as a first means of evaluating the suitability of fish species for long-term tracking, and illustrated how telemetry approaches can be selected and combined to address questions relating to species-specific knowledge gaps and migratory strategies. Specifically, acoustic telemetry provides options for fish tracking at various spatial and temporal scales (Fig. 2), but is limited by the spatial coverage of receivers in offshore marine waters and cooperation among telemetry researchers. While lacking positional information, sensor data from acoustic and archival tags and PSATs can be used to understand the environmental drivers of movement within and beyond coastal waters, and provide opportunities to model movement trajectories beyond the extent of acoustic arrays (Figs. 8 and 9).

While each of these methods are already available for use, they are not always employed to their full potential. We here discuss some of the limitations that should be considered and addressed for open coastal ecosystems in general and propose future directions for research in the Dutch Wadden Sea.

3.2. Perspectives for long-term tracking in open coastal ecosystems

For many migratory species, current knowledge of coastal behaviours may be insufficient to combat locally-observed declines. Despite more holistic efforts to monitor fish behaviours across broader spatial scales, acoustic receiver arrays are typically geographically-limited and rarely extend beyond national borders (Abecasis et al., 2018). Acoustic telemetry networks expand the scope of animal tracking studies to ecologically-relevant scales, but require commitment from independent researchers to ensure equipment compatibility, prompt and regular inputs of data to online databases, and coordinated placement of acoustic arrays (Reubens et al., 2021). At the ETN, these issues are addressed via annual meetings that foster collaborations, promote equipment and data compatibility, such as via the introduction of Open Protocols (Reubens et al., 2021) (<https://www.europeantrackingnetwork.org/en/open-protocol>), and outreach activities, including workshops for telemetry researchers and online tutorials (<https://www.youtube.com/@europeantrackingnetwork>). Researchers tagging migratory fish in the Wadden Sea can benefit from existing platforms such as this to prioritise collaborations at regional, national, and international levels.

In an era of rapid environmental change, long-term datasets may act as valuable indicators of shifts in fish communities, including species replacements, modified behaviours, or altered distributions (Daufresne et al., 2004; Hoegh-Guldberg and Bruno, 2010; Perry et al., 2005; Rodriguez-Dominguez et al., 2019). However, most fish tracking studies are conducted over a period of only a few years, limiting their practicality for long-term monitoring. For example, acoustic receiver arrays managed by independent research institutes are often deployed temporarily, resulting in changes to receiver coverage within or across coastal regions between years, and potentially obscuring infrequent or unexpected fish behaviours that may be of high ecological importance (Lennox et al., 2024). Acoustic arrays are both costly and time consuming to establish and maintain, prohibiting extensive coverage, and should therefore be designed with the aim of maximizing data output and applicability (Lennox et al., 2024). As such, array managers should coordinate across at least, regional, and at best, international scales to maximize the potential for animal detection and address relevant study questions. Recent studies have already proposed the deployment of receiver gates at important transition points and

jurisdictional borders (Abecasis et al., 2018; Lennox et al., 2024), a task which is currently being undertaken by a pan-European network covering four key locations in Europe: 1) the Danish Straits, 2) the North Channel, 3) the Strait of Gibraltar, and 4) the Strait of Bosphorus and Dardanelles (<https://www.europeantrackingnetwork.org/en/straits>). Networks of permanent or semi-permanent arrays will be critical for identifying long-term trends in population- or community-level responses to environmental change.

Limited study durations are also problematic for methods such as archival tagging in which tag recovery rates may be low - particularly for non-commercial species - and in which significant delays between tag deployment and recovery may prolong the collection of sufficient data for statistical analyses (Domeier et al., 2018). By comparison, PSATs provide more predictable data returns over shorter time periods (Hussey et al., 2015), but are more cost-prohibitive and unsuitable for animals below a threshold body size. Furthermore, archived data are binned prior to transmission to satellites, leading to lower resolutions compared to the data records available upon tag recovery (Edwards et al., 2019). As such, the potential value of archival tag recoveries should not be overlooked. Instead, the time periods predicted for sufficient data recovery should be factored into project proposals and funding schemes.

3.3. Recommendations for future tracking in the Wadden Sea

The Dutch Wadden Sea was selected as a prime example of an open coastal system requiring enhanced understanding of local and regional fish mobility to manage severe population declines. Using this system as a case study, highlighted the importance of existing knowledge of population trends, species-specific life histories, biological parameters (e.g., size and morphology) for defining research questions and identifying appropriate telemetry techniques. We identified 28 fish species that commonly occur in the Wadden Sea as being potentially suitable for tagging (Table 1), and suggest this as a starting point for future telemetry studies. We also recommend the consideration of aspects such as population status, catchability - including availability of skilled and knowledgeable fisherman -, and vulnerability to capture and handling when selecting potential study species and associated telemetry devices.

In this study, we aimed to showcase the application of acoustic, archival, and PSAT technologies to examine the full range of migratory fish behaviours and identify population bottlenecks across coastal, marine, and fresh waters. Specific guidelines for where and when these techniques should be employed, and for which species, are highly dependant on individual research questions and are therefore difficult to provide. Instead we hope that this work will aid in selecting appropriate telemetry techniques for a predetermined research question and/or species. Furthermore, we advocate for the selection of management-driven research objectives, particularly in the face of significant population declines (Matley et al., 2022). For example, the importance of coastal environments for occasional visitors might be difficult to discern due to low observation rates or unpredictability, but could be crucial for overall population success. Large-scale acoustic telemetry networks and environmental data recorded by archival tags and PSATs could thereby be essential for revealing broad migration pathways and highlighting interregional connectivity between adult and juvenile habitats and subpopulations (McMillan et al., 2019). For seasonal residents, acoustic receiver arrays and long-lifespan tags can be used to define periods and individual fish occurrence over seasonal and multi-year periods and illustrate fine-scale behaviours within coastal waters, informing the delineation and refinement of local protections and area closures (Kohler et al., 2023; Nemeth et al., 2023). Meanwhile, tracks reconstructed using archival data from double-tagged fish or other individuals within a population, alongside acoustic telemetry networks, can fill in knowledge gaps in offshore marine regions, other coastal habitats, or adjacent freshwaters (Goossens et al., 2023; Thorburn et al., 2019). For diadromous species that are often required pass through a variety of anthropogenic barriers en route between freshwater to offshore marine

environments, acoustic receivers deployed strategically at habitat boundaries and transition zones can pinpoint barriers to regional connectivity and direct restoration efforts (Verhelst et al., 2018). Acoustic detection data can be supplemented by environmental records derived from animal-borne data storage tags, sensors, or data repositories to infer the factors driving fine-scale behaviours and habitat use across a spatially and temporally variable environment. Ultimately, the management of migratory fish populations will require a thorough understanding of the ecological role of coastal habitats in fish life cycles to determine when and where spatial management actions can be applied most effectively.

Declaration of generative Ai in scientific Writing

During the preparation of this work the author(s) used ChatGPT, a language generation tool developed by OpenAI, in order to assist in diversifying word choice and grammar. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

CRedit authorship contribution statement

J.E. Edwards: Conceptualization, Data curation, Investigation, Visualization, Writing – original draft, Writing – review & editing. **A.D. Buijse:** Conceptualization, Supervision, Writing – review & editing. **H. V. Winter:** Conceptualization, Data curation, Supervision, Writing – review & editing. **A. van Leeuwen:** Data curation, Investigation, Supervision, Writing – review & editing. **A.I. Bijleveld:** Conceptualization, Data curation, Supervision, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We would like to thank all those dedicated to maintaining the long-standing NIOZ fyke sampling program as well as the many local fishers, students, and colleagues who donated their expertise and sleepless nights while taking part in the capture and tagging of sea bass and mullet exemplified in this study. We would like to acknowledge the invaluable contributions of our project partners, Sportvisserij Nederland and Rijkswaterstaat for facilitating fieldwork related to both fish tagging and acoustic receiver deployments. Thank you also to the journal editor and anonymous reviewer for providing essential feedback for the improvement of this manuscript.

Funding for this study was provided by Waddenfonds, Rijkswaterstaat, the provinces of North Holland, Friesland, and Groningen, and the Dutch Ministry of Agriculture, Nature, and Food Quality as part of the Waddentools Swimway Wadden Sea project.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecss.2024.108737>.

References

- Aarestrup, K., Baktoft, H., Thorstad, E.B., Svendsen, J.C., Höjesjö, J., Koed, A., 2015. Survival and progression rates of anadromous brown trout kelts *Salmo trutta* during downstream migration in freshwater and at sea. *Mar. Ecol. Prog. Ser.* 535, 185–195.
- Abecasis, D., Steckenreuter, A., Reubens, J., Aarestrup, K., Alós, J., Badalamenti, F., Bajona, L., Boylan, P., Deneudt, K., Greenberg, L., Brevé, N., Hernández, F., Humphries, N., Meyer, C., Sims, D., Thorstad, E.B., Walker, A.M., Whoriskey, F., Afonso, P., 2018. A review of acoustic telemetry in Europe and the need for a regional aquatic telemetry network. *Animal Biotelemetry* 6 (1), 12. <https://doi.org/10.1186/s40317-018-0156-0>.
- Able, K.W., Grothues, T.M., Turnure, J.T., Malone, M.A., Henkes, G.A., 2014. Dynamics of residency and egress in selected estuarine fishes: evidence from acoustic telemetry. *Environ. Biol. Fish.* 97 (1), 91–102. <https://doi.org/10.1007/s10641-013-0126-6>.
- Aldvén, D., Hedger, R., Økland, F., Rivinoja, P., Höjesjö, J., 2015. Migration speed, routes, and mortality rates of anadromous brown trout *Salmo trutta* during outward migration through a complex coastal habitat. *Mar. Ecol. Prog. Ser.* 541 (January 2016), 151–163. <https://doi.org/10.3354/meps11535>.
- Allen, A.M., Singh, N.J., 2016. Linking movement ecology with wildlife management and conservation. *Frontiers in Ecology and Evolution* 3, 155.
- Bachelor, N.M., Buckel, J.A., Hightower, J.E., Paramore, L.M., Pollock, K.H., 2009. A combined telemetry – tag return approach to estimate fishing and natural mortality rates of an estuarine fish. *Can. J. Fish. Aquat. Sci.* 66 (8), 1230–1244. <https://doi.org/10.1139/F09-076>.
- Baktoft, H., Gjelland, K.Ø., Økland, F., Thygesen, U.H., 2017. Positioning of aquatic animals based on time-of-arrival and random walk models using YAPS (Yet Another Positioning Solver). *Sci. Rep.* 7 (1), 14294.
- Baptist, M.J., van der Wal, J.T., Folmer, E.O., Gräwe, U., Elschot, K., 2019. An ecotope map of the trilateral Wadden Sea. *J. Sea Res.* 152 (November 2018), 101761. <https://doi.org/10.1016/j.seares.2019.05.003>.
- Barkley, A.N., Gollock, M., Samoilys, M., Llewellyn, F., Shivji, M., Wetherbee, B., Hussey, N.E., 2019. Complex transboundary movements of marine megafauna in the Western Indian Ocean. *Anim. Conserv.* 22 (5), 420–431.
- Batsleer, J., Bleeker, K., Brunel, T., van Hal, R., Staat, L., 2020. Overzicht beschikbare gegevens ten behoeve van Nederlandse beleidsdoelen voor haaien en roggen.
- Beck, M.W., Heck, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M., Halpern, B., Hays, C.G., Hoshino, K., Minello, T.J., Orth, R.J., Sheridan, P.F., Weinstein, M.P., 2001. The Identification, Conservation, and Management of Estuarine and Marine Nurseries for Fish and Invertebrates: a better understanding of the habitats that serve as nurseries for marine species and the factors that create site-specific variability in nursery quality will improve conservation and management of these areas. *Bioscience* 51 (8), 633–641. [https://doi.org/10.1641/0006-3568\(2001\)051\[0633:TICAMO\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0633:TICAMO]2.0.CO;2).
- Bendall, B., Moore, A., Quayle, V., 2005. The post-spawning movements of migratory brown trout *Salmo trutta* L. *J. Fish. Biol.* 67, 809–822. <https://doi.org/10.1111/j.1095-8649.2005.00786.x>.
- Berg, S., Krog, C., Muus, B., Nielsen, J., Fricke, R., Berghahn, R., Neudecker, Th., Wolff, W.J., 1996. IX. Red List of lampreys and marine fishes of the Wadden Sea. *Helgol. Meeresunters.* 50 (Suppl. L.), 101–105. <https://doi.org/10.1007/bf02366178>.
- Bij De Vaate, A., Breukelaar, A.W., Vriese, T., De Laak, G., Dijkers, C., 2003. Sea trout migration in the Rhine delta. *J. Fish. Biol.* 63 (4), 892–908. <https://doi.org/10.1046/j.1095-8649.2003.00198.x>.
- Biton-porsmoguer, S., 2022. Diet strategies of starry smooth-hound *Mustelus asterias* and tope shark *Galeorhinus galeus* (Carcharhiniformes: Triakidae) in the Eastern English Channel: implication for conservation. *Can. Biol. Mar.* 63 (2), 129–137.
- Block, B.A., Dewar, H., Farwell, C., Prince, E.D., 1998. A new satellite technology for tracking the movements of Atlantic bluefin tuna. *Proc. Natl. Acad. Sci. USA* 95 (16), 9384–9389. <https://doi.org/10.1073/pnas.95.16.9384>.
- Boer, P., 1971. De andere harder. *Levende Nat.* 74 (3), 62–65.
- Borcherding, J., Pichardt, C., Winter, H.V., Becker, J.S., 2008. Migration history of North Sea houting (*Coregonus oxyrinchus* L.) caught in Lake IJsselmeer (The Netherlands) inferred from scale transects of 88 Sr: 44 Ca ratios. *Aquat. Sci.* 70, 47–56.
- Brasseur, S.M., van Polanen Petel, T.D., Gerrodette, T., Meesters, E.H., Reijnders, P.J., Aarts, G., 2015. Rapid recovery of Dutch gray seal colonies fueled by immigration. *Mar. Mamm. Sci.* 31 (2), 405–426.
- Braun, C.D., Galuardi, B., Thorrold, S.R., 2018. HMMocean: an R package for improved geolocation of archival-tagged fishes using a hidden Markov method. *Methods Ecol. Evol.* 9 (5), 1212–1220.
- Breece, M.W., Fox, D.A., Oliver, M.J., 2018. Environmental drivers of adult Atlantic sturgeon movement and residency in the Delaware Bay. *Marine and Coastal Fisheries* 10 (2), 269–280.
- Breen, P., Posen, P., Righton, D., 2015. Temperate Marine Protected Areas and highly mobile fish: a review. *Ocean Coast Manag.* 105, 75–83. <https://doi.org/10.1016/j.ocecoaman.2014.12.021>.
- Brooks, J.L., Chapman, J.M., Barkley, A.N., Kessel, S.T., Hussey, N.E., Hinch, S.G., Patterson, D.A., Hedges, K.J., Cooke, S.J., Fisk, A.T., Gruber, S.H., Nguyen, V.M., 2019. Biotelemetry informing management: case studies exploring successful integration of biotelemetry data into fisheries and habitat management. *Can. J. Fish. Aquat. Sci.* 76 (7), 1238–1252. <https://doi.org/10.1139/cjfas-2017-0530>.
- Brown, E.J., Vasconcelos, R.P., Wennhage, H., Bergström, U., Støttrup, J.G., van de Wolfshaar, K., Millisenda, G., Colloca, F., Le Pape, O., 2018. Conflicts in the coastal zone: human impacts on commercially important fish species utilizing coastal

- habitat. *ICES (Int. Council. Explor. Sea) J. Mar. Sci.* 75 (4), 1203–1213. <https://doi.org/10.1093/icesjms/lsx237>.
- Brownscombe, J.W., Griffin, L.P., Brooks, J.L., Danylchuk, A.J., Cooke, S.J., Midwood, J. D., 2022a. Applications of telemetry to fish habitat science and management. *Can. J. Fish. Aquat. Sci.* 79 (8), 1347–1359.
- Brownscombe, J.W., Shipley, O.N., Griffin, L.P., Morley, D., Acosta, A., Adams, A.J., Boucek, R., Danylchuk, A.J., Cooke, S.J., Power, M., 2022b. Application of telemetry and stable isotope analyses to inform the resource ecology and management of a marine fish. *J. Appl. Ecol.* 59 (4), 1110–1121.
- Camphuysen, C.J., 2004. The return of the harbour porpoise (*Phocoena phocoena*) in Dutch coastal waters. *Lutra* 47 (2), 113–122.
- Capapé, C., Ben Souissi, J., Méjri, H., Guélorget, O., Hemida, F., 2005. The reproductive biology of the school shark, *Galeorhinus galeus* Linnaeus 1758 (Chondrichthyes: Triakidae), from the Maghreb shore (southern Mediterranean). *Acta Adriat.: Int. J. Mar. Sci.* 46 (2), 109–124.
- Cardona, L., 2006. Habitat selection by grey mullets (*Osteichthyes: mugilidae*) in Mediterranean estuaries: the role of salinity. *Sci. Mar.* 70 (September), 443–455.
- Cardoso, J.F.M.F., Freitas, V., Quilez, I., Jouta, J., Witte, J.I., van der Veer, H.W., 2015. The European sea bass *Dicentrarchus labrax* in the Dutch Wadden Sea: from visitor to resident species. *J. Mar. Biol. Assoc. U. K.* 95 (4), 839–850. <https://doi.org/10.1017/S0025315414001714>.
- Chaput, G., Carr, J., Daniels, J., Tinker, S., Jonsen, I., Whoriskey, F., 2019. Atlantic salmon (*Salmo salar*) smolt and early post-smolt migration and survival inferred from multi-year and multi-stock acoustic telemetry studies in the Gulf of St. Lawrence, northwest Atlantic. *ICES (Int. Council. Explor. Sea) J. Mar. Sci.* 76 (4), 1107–1121. <https://doi.org/10.1093/icesjms/fsy156>.
- Colloca, F., Scannella, D., Geraci, M.L., Falson, F., Giusto, B., Vitale, S., Di Lorenzo, M., Bono, G., 2019. British sharks in Sicily: records of long-distance migration of tope shark (*Galeorhinus galeus*) from the north-eastern Atlantic to the Mediterranean Sea. *Mediterr. Mar. Sci.* 20 (2), 309–313. <https://doi.org/10.12681/MMS.18121>.
- Common Wadden Sea Secretariat, 2016. Report on the State of Conservation of the World Heritage Property “The Wadden Sea. *NI314*).
- Cooke, S.J., Auld, H.L., Birnie-Gauvin, K., Elvidge, C.K., Piczak, M.L., Twardek, W.M., Raby, G.D., Brownscombe, J.W., Midwood, J.D., Lennox, R.J., Madliger, C., Wilson, A.D.M., Binder, T.R., Schreck, C.B., McLaughlin, R.L., Grant, J., Muir, A.M., 2023. On the relevance of animal behavior to the management and conservation of fishes and fisheries. *Environ. Biol. Fish.* 106 (5), 785–810. <https://doi.org/10.1007/s10641-022-01255-3>.
- Cooke, S.J., Martins, E.G., Struthers, D.P., Gutowsky, L.F.G., Power, M., Doka, S.E., Dettmers, J.M., Crook, D.A., Lucas, M.C., Holbrook, C.M., Krueger, C.C., 2016. A moving target—incorporating knowledge of the spatial ecology of fish into the assessment and management of freshwater fish populations. *Environ. Monit. Assess.* 44 (0), 19.
- Corrigan, S., Lowther, A.D., Beheregaray, L.B., Bruce, B.D., Cliff, G., Duffy, C.A., Foulis, A., Francis, M.P., Goldsworthy, S.D., Hyde, J.R., Jabado, R.W., Kacev, D., Marshall, L., Mucientes, G.R., Naylor, G.J.P., Pepperell, J.G., Queiroz, N., White, W. T., Wintner, S.P., Rogers, P.J., 2018. Population connectivity of the highly migratory shortfin mako (*Isurus oxyrinchus Rafinesque* 1810) and implications for management in the Southern Hemisphere. *Frontiers in Ecology and Evolution* 6 (NOV), 1–15. <https://doi.org/10.3389/fevo.2018.00187>.
- Crossin, G.T., Heupel, M.R., Holbrook, C.M., Hussey, N.E., Lowerre-Barbieri, S.K., Nguyen, V.M., Raby, G.D., Cooke, S.J., 2017. Acoustic telemetry and fisheries management. *Ecol. Appl.* 27 (4), 1031–1049. <https://doi.org/10.1002/eap.1533>.
- Daly, R., Fimalter, J.D., Daly, C.A.K., Bennett, R.H., Pereira, M.A.M., Mann, B.Q., Dunlop, S.W., Cowley, P.D., 2019. Acoustic telemetry reveals multi-seasonal spatiotemporal dynamics of a giant trevally *Caranx ignobilis* aggregation. *Mar. Ecol. Prog. Ser.* 621, 185–197. <https://doi.org/10.3354/meps12975>.
- Daufresne, M., Roger, M.C., Capra, H., Lamouroux, N., 2004. Long-term changes within the invertebrate and fish communities of the Upper Rhône River: effects of climatic factors. *Global Change Biol.* 10 (1), 124–140. <https://doi.org/10.1046/j.1529-8817.2003.00720.x>.
- de Pontual, H., Lalire, M., Fablet, R., Laspougeas, C., Garren, F., Martin, S., Drogou, M., Woillez, M., 2019. New insights into behavioural ecology of European seabass off the West Coast of France: implications at local and population scales. *ICES (Int. Council. Explor. Sea) J. Mar. Sci.* 76 (2), 501–515.
- Degerman, E., Leonardsson, K., Lundqvist, H., 2012. Coastal migrations, temporary use of neighbouring rivers, and growth of sea trout (*Salmo trutta*) from nine northern Baltic Sea rivers. *ICES (Int. Council. Explor. Sea) J. Mar. Sci.* 69 (6), 971–980. <https://doi.org/10.1093/icesjms/fss073>.
- Domeier, M.L., Ortega-García, S., Nasby-Lucas, N., Offield, P., 2018. First marlin archival tagging study suggests new direction for research. *Mar. Freshw. Res.* 70 (4), 603–608.
- Donaldson, M.R., Hinch, S.G., Suski, C.D., Fisk, A.T., Heupel, M.R., Cooke, S.J., 2014. Making connections in aquatic ecosystems with acoustic telemetry monitoring. *Front. Ecol. Environ.* 12 (10), 565–573. <https://doi.org/10.1890/130283>.
- Doyle, T.K., Haberlin, D., Clohessy, J., Bennison, A., Jessopp, M., 2017. Localised residency and inter-annual fidelity to coastal foraging areas may place sea bass at risk to local depletion. *Sci. Rep.* 7 (1), 45841.
- Dudgeon, C.L., Pollock, K.H., Braccini, J.M., Semmens, J.M., Barnett, A., 2015. Integrating acoustic telemetry into mark-recapture models to improve the precision of apparent survival and abundance estimates. *Oecologia* 178 (3), 761–772. <https://doi.org/10.1007/s00442-015-3280-z>.
- Edgar, G.J., Stuart-Smith, R.D., Willis, T.J., Kininmonth, S., Baker, S.C., Banks, S., Barrett, N.S., Becerro, M.A., Bernard, A.T.F., Berkhout, J., Buxton, C.D., Campbell, S. J., Cooper, A.T., Davey, M., Edgar, S.C., Försterra, G., Galván, D.E., Irigoyen, A.J., Kushner, D.J., et al., 2014. Global conservation outcomes depend on marine protected areas with five key features. *Nature* 506 (7487). <https://doi.org/10.1038/nature13022>. Article 7487.
- Edwards, J.E., Hedges, K.J., Hussey, N.E., 2022a. Seasonal residency, activity space, and use of deep-water channels by Greenland sharks (*Somniosus microcephalus*) in an Arctic fjord system. *Can. J. Fish. Aquat. Sci.* 79 (2), 314–330.
- Edwards, J.E., Hedges, K.J., Kessel, S.T., Hussey, N.E., 2022b. Multi-year acoustic tracking reveals transient movements, recurring hotspots, and apparent seasonality in the coastal-offshore presence of Greenland sharks (*Somniosus microcephalus*). *Front. Mar. Sci.* 9. <https://www.frontiersin.org/articles/10.3389/fmars.2022.902854>.
- Edwards, J.E., Pratt, J., Tress, N., Hussey, N.E., 2019. Thinking deeper: uncovering the mysteries of animal movement in the deep sea. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 146 <https://doi.org/10.1016/j.dsr.2019.02.006>.
- Eldoy, S.H., Davidsen, J.G., Thorstad, E.B., Whoriskey, F., Aarestrup, K., Næsje, T.F., Rønning, L., Sjørnsen, A.D., Rikardsen, A.H., Arnekleiv, J.V., 2015. Marine migration and habitat use of anadromous brown trout (*Salmo trutta*). *Can. J. Fish. Aquat. Sci.* 72 (9), 1366–1378. <https://doi.org/10.1139/cjfas-2014-0560>.
- Eldoy, S.H., Davidsen, J.G., Thorstad, E.B., Whoriskey, F.G., Aarestrup, K., Næsje, T.F., Rønning, L., Sjørnsen, A.D., Rikardsen, A.H., Arnekleiv, J.V., 2017. Marine depth use of sea trout *Salmo trutta* in fjord areas of central Norway. *J. Fish. Biol.* 91 (5), 1268–1283.
- Elliott, M., Hemingway, K.L., 2008. *Fishes in Estuaries*. John Wiley & Sons.
- Elliott, M., Whitfield, A.K., Potter, I.C., Blaber, S.J.M., Cyrus, D.P., Nordlie, F.G., Harrison, T.D., 2007. The guild approach to categorizing estuarine fish assemblages: a global review. *Fish. Fish.* 8 (3), 241–268. <https://doi.org/10.1111/j.1467-2679.2007.00253.x>.
- Elston, C., Cowley, P.D., Murray, T.S., Parkinson, M.C., 2023. Novel insights into coastal site affinity and habitat connectivity of a benthic stingray with implications for management. *Biodivers. Conserv.* 32 (1), 181–202.
- Elston, C., Cowley, P.D., von Brandis, R.G., Lea, J., 2022. Stingray habitat use is dynamically influenced by temperature and tides. *Front. Mar. Sci.* 8. <https://www.frontiersin.org/articles/10.3389/fmars.2021.754404>.
- Espinoza, M., Lédée, E.J.I., Smoothey, A.F., Heupel, M.R., Peddemors, V.M., Tobin, A.J., Simpfordorfer, C.A., 2021. Intra-specific variation in movement and habitat connectivity of a mobile predator revealed by acoustic telemetry and network analyses. *Mar. Biol.* 168 (6), 80. <https://doi.org/10.1007/s00227-021-03886-z>.
- Finstad, B., Økland, F., Thorstad, E.B., Bjørn, P.A., McKinley, R.S., 2005. Migration of hatchery-reared Atlantic salmon and wild anadromous brown trout post-smolts in a Norwegian fjord system. *J. Fish. Biol.* 66 (1), 86–96.
- Fitzmaurice, P., Keirse, G., Green, P., Clarke, M., 2003. *Tagging in Irish Waters (1970–2002)*. Central Fisheries Board.
- Flaten, A.C., Davidsen, J.G., Thorstad, E.B., Whoriskey, F., Rønning, L., Sjørnsen, A.D., Rikardsen, A.H., Arnekleiv, J.V., 2016. The first months at sea: marine migration and habitat use of sea trout *Salmo trutta* post-smolts. *J. Fish. Biol.* 89 (3), 1624–1640. <https://doi.org/10.1111/jfb.13065>.
- Frisk, M.G., Shipley, O.N., Martinez, C.M., McKown, K.A., Zacharias, J.P., Dunton, K.J., 2019. First observations of long-distance migration in a large skate species, the winter skate: implications for population connectivity, ecosystem dynamics, and management. *Marine and Coastal Fisheries* 11 (2), 202–212.
- Froese, R., Pauly, D., 2023. FishBase. <https://www.fishbase.org/summary/citation.php>.
- Gillanders, B.M., Able, K.W., Brown, J.A., Eggleston, D.B., Sheridan, P.F., 2003. Evidence of connectivity between juvenile and adult habitats for mobile marine fauna: an important component of nurseries. *Mar. Ecol. Prog. Ser.* 247, 281–295. <https://doi.org/10.3354/meps247281>.
- Gjelland, K.Ø., Hedger, R.D., 2017. On the parameterization of acoustic detection probability models. *Methods Ecol. Evol.* 8 (10), 1302–1304.
- Goossens, J., Woillez, M., LeBris, A., Verhelst, P., Moens, T., Torrelee, E., Reubens, J., 2023. Acoustic and archival technologies join forces: A combination tag. *Methods Ecol. Evol.* 14 (3), 860–866.
- Grothues, T.M., 2009. A Review of Acoustic Telemetry Technology and a Perspective on its Diversification Relative to Coastal Tracking Arrays. In: Nielsen, J.L., Arrizabalaga, H., Fragoso, N., Hobday, A., Lutcavage, M., Sibert, J. (Eds.), *Tagging and Tracking of Marine Animals with Electronic Devices*. Springer Netherlands, pp. 77–90. https://doi.org/10.1007/978-1-4020-9640-2_5.
- Grüss, A., Kaplan, D.M., Hart, D.R., 2011. Relative impacts of adult movement, larval dispersal and harvester movement on the effectiveness of reserve networks. *PLoS One* 6 (5), e19960. <https://doi.org/10.1371/journal.pone.0019960>.
- Hammerschlag, N., Gallagher, A.J., Lazarre, D.M., 2011. A review of shark satellite tagging studies. *J. Exp. Mar. Biol. Ecol.* 398 (1–2), 1–8.
- Harcourt, R., Sequeira, A.M.M., Zhang, X., Roquet, F., Komatsu, K., Heupel, M., McMahon, C., Whoriskey, F., Meekan, M., Carroll, G., Brodie, S., Simpfordorfer, C., Hindell, M., Jonsen, I., Costa, D.P., Block, B., Muelbert, M., Woodward, B., Weise, M., et al., 2019. Animal-borne telemetry: an integral component of the Ocean observing toolkit. *Front. Mar. Sci.* 6. <https://www.frontiersin.org/articles/10.3389/fmars.2019.00326>.
- Hazen, E.L., Maxwell, S.M., Bailey, H., Bograd, S.J., Hamann, M., Gaspar, P., Godley, B. J., Shillinger, G.L., 2012. Ontogeny in marine tagging and tracking science: technologies and data gaps. *Mar. Ecol. Prog. Ser.* 457, 221–240. <https://doi.org/10.3354/meps09857>.
- Hedger, R.D., Rikardsen, A.H., Thorstad, E.B., 2017. Pop-up satellite archival tag effects on the diving behaviour, growth and survival of adult Atlantic salmon *Salmo salar* at sea. *J. Fish. Biol.* 90 (1), 294–310. <https://doi.org/10.1111/jfb.13174>.
- Heupel, M.R., 2007. Exiting Terra Ceia Bay: an examination of cues stimulating migration from a summer nursery area. *Am. Fish. Soc. Symp.* 50, 265. <https://fisheries.org/docs/books/54050P/17.pdf>.

- Heupel, M.R., Semmens, J.M., Hobday, A.J., 2006. Automated acoustic tracking of aquatic animals: scales, design and deployment of listening station arrays. *Mar. Freshw. Res.* 500, 1–13.
- Heupel, M.R., Simpfendorfer, C.A., 2002. Estimation of mortality of juvenile blacktip sharks, *Carcharhinus limbatus*, within a nursery area using telemetry data. *Can. J. Fish. Aquat. Sci.* 59 (4), 624–632.
- Heupel, M.R., Simpfendorfer, C.A., 2005. Using acoustic monitoring to evaluate MPAs for shark nursery areas: the importance of long-term data. *Mar. Technol. Soc. J.* 39, 10–18.
- Heupel, M.R., Webber, D.M., 2012. Trends in acoustic tracking: where are the fish going and how will we follow them. *Advances in fish tagging and marking technology* 76.
- Hickling, C.F., 1970. A contribution to the natural history of the English grey mullets [pisces, mugilidae]. *J. Mar. Biol. Assoc. U. K.* 50 (3), 609–633. <https://doi.org/10.1017/S0025315400004914>.
- Hoegh-Guldberg, O., Bruno, J.F., 2010. The impact of climate change on the world's marine ecosystems. *Science* 328 (5985), 1523–1528. <https://doi.org/10.1126/science.1189930>.
- Hoenner, X., Huveneres, C., Steckenreuter, A., Simpfendorfer, C., Tattersall, K., Jaïne, F., Atkins, N., Babcock, R., Brodie, S., Burgess, J., 2018. Australia's continental-scale acoustic tracking database and its automated quality control process. *Sci. Data* 5 (1), 1–10.
- Holden, M.J., Horrod, R.G., 1979. The migrations of tope, *galeorhinus galeus* (L), in the eastern north Atlantic as determined by tagging. *ICES (Int. Counc. Explor. Sea) J. Mar. Sci.* 38 (3), 314–317. <https://doi.org/10.1093/icesjms/38.3.314>.
- Holden, M.J., Williams, T., 1974. The biology, movements and population dynamics of bass, *Dicentrarchus labrax*, in English waters. *J. Mar. Biol. Assoc. U. K.* 70, 91–107.
- Hovenkamp, F., 1991. Immigration of larval plaice (*Pleuronectes platessa* L.) into the western wadden sea: a question of timing. *Neth. J. Sea Res.* 27 (3), 287–296. [https://doi.org/10.1016/0077-7579\(91\)90031-U](https://doi.org/10.1016/0077-7579(91)90031-U).
- Hunter, E., Buckley, A.A., Stewart, C., Metcalfe, J.D., 2005. Migratory behaviour of the thornback ray, *Raja clavata*, in the southern North Sea. *JMBA-Journal of the Marine Biological Association of the United Kingdom* 85 (5), 1095–1106.
- Hussey, N.E., Kessel, S.T., Aarestrup, K., Cooke, S.J., Cowley, P.D., Fisk, A.T., Harcourt, R.G., Holland, K.N., Iverson, S.J., Kocik, J.F., Flemming, J.E.M., Whoriskey, F.G., 2015. Aquatic animal telemetry: a panoramic window into the underwater world. *Science* 348 (6240), 1255642. <https://doi.org/10.1126/science.1255642>.
- Iverson, S.J., Fisk, A.T., Hinch, S.G., Mills Flemming, J., Cooke, S.J., Whoriskey, F.G., 2019. The Ocean Tracking Network: advancing frontiers in aquatic science and management. *Can. J. Fish. Aquat. Sci.* 76 (7), 1041–1051.
- Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjørndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H., Cooke, R., Erlanson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R.R., 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293 (5530), 629–637. <https://doi.org/10.1126/science.1059199>.
- Jager, Z., Bolle, L.J., Neudecker, T., Vorberg, R., Damm, U., Diederichs, B., Jager, Z., Scholle, J., Dänhardt, A., Luerssen, G., Marencic, H., 2009. Trends in Wadden Sea Fish Fauna, Part I: Trilateral Cooperation.
- James, M.C., Ottensmeyer, C.A., Myers, R.A., 2005. Identification of high-use habitat and threats to leatherback sea turtles in northern waters: new directions for conservation. *Ecol. Lett.* 8 (2), 195–201. <https://doi.org/10.1111/j.1461-0248.2004.00710.x>.
- Jensen, L.F., Rognon, P., Aarestrup, K., Böttcher, J.W., Pertoldi, C., Thomsen, S.N., Hertz, M., Winde, J., Svendsen, J.C., 2018. Evidence of cormorant-induced mortality, disparate migration strategies and repeatable circadian rhythm in the endangered North Sea houting (*Coregonus oxyrinchus*): a telemetry study mapping the postspawning migration. *Ecol. Freshw. Fish* 27 (3), 672–685.
- Jepsen, N., Schreck, C., Clements, S., Thorstad, E.B., 2005. A brief discussion on the 2% tag/bodymass rule of thumb. *Aquatic Telemetry: Advances and Applications* 255–259.
- Kennedy, M., Fitzmaurice, P., 1969. Age and growth of thick-lipped grey mullet *Crenimugil labrosus* in Irish waters. *J. Mar. Biol. Assoc. U. K.* 49 (3), 683–699.
- Kennedy, M., Fitzmaurice, P., 1972. The biology of the bass, *Dicentrarchus labrax*, in Irish waters. *J. Mar. Biol. Assoc. U. K.* 52 (3), 557–597.
- Kessel, S.T., Cooke, S.J., Heupel, M.R., Hussey, N.E., Simpfendorfer, C.A., Vagle, S., Fisk, A.T., 2014. A review of detection range testing in aquatic passive acoustic telemetry studies. *Rev. Fish Biol. Fish.* 24 (1), 199–218. <https://doi.org/10.1007/s11160-013-9328-4>.
- Klinard, N.V., Matley, J.K., Fisk, A.T., Johnson, T.B., 2019. Long-term retention of acoustic telemetry transmitters in temperate predators revealed by predation tags implanted in wild prey fish. *J. Fish. Biol.* 95 (6), 1512–1516.
- Kohler, J., Gore, M., Ormond, R., Johnson, B., Austin, T., 2023. Individual residency behaviours and seasonal long-distance movements in acoustically tagged Caribbean reef sharks in the Cayman Islands. *PLoS One* 18 (11), e0293884. <https://doi.org/10.1371/journal.pone.0293884>.
- Kristensen, M.L., Pedersen, M.W., Thygesen, U.H., Del Villar-Guerra, D., Baktoft, H., Aarestrup, K., 2019. Migration routes and habitat use of a highly adaptable salmonid (sea trout, *Salmo trutta*) in a complex marine area. *Animal Biotelemetry* 7 (1). <https://doi.org/10.1186/s40317-019-0185-3>.
- Kristensen, M.L., Righton, D., Del Villar-Guerra, D., Baktoft, H., Aarestrup, K., 2018. Temperature and depth preferences of adult sea trout *Salmo trutta* during the marine migration phase. *Mar. Ecol. Prog. Ser.* 599 (September), 209–224. <https://doi.org/10.3354/meps12618>.
- Larocque, S.M., Johnson, T.B., Fisk, A.T., 2020. Survival and migration patterns of naturally and hatchery-reared Atlantic salmon (*Salmo salar*) smolts in a Lake Ontario tributary using acoustic telemetry. *Freshw. Biol.* 65 (5), 835–848. <https://doi.org/10.1111/fwb.13467>.
- Lédée, E.J., Heupel, M.R., Tobin, A.J., Simpfendorfer, C.A., 2015. Movements and space use of giant trevally in coral reef habitats and the importance of environmental drivers. *Animal Biotelemetry* 3, 1–14.
- Lees, K.J., MacNeil, M.A., Hedges, K.J., Hussey, N.E., 2021. Estimating demographic parameters for fisheries management using acoustic telemetry. *Rev. Fish Biol. Fish.* 31 (1), 25–51. <https://doi.org/10.1007/s11160-020-09626-8>.
- Leewis, L., van Bodegom, P.M., Rozema, J., Janssen, G.M., 2012. Does beach nourishment have long-term effects on intertidal macroinvertebrate species abundance? *Estuar. Coast Shelf Sci.* 113, 172–181.
- Lennox, R.J., Aarestrup, K., Alós, J., Arlinghaus, R., Aspillaga, E., Bertram, M.G., Birnie-Gauvin, K., Brodin, T., Cooke, S.J., Dahlmo, L.S., Dhellemmes, F., Gjelland, K.Ø., Hellström, G., Hershey, H., Holbrook, C., Klefoth, T., Lowerre-Barbieri, S., Monk, C. T., Nilsen, C.I., et al., 2023a. Positioning aquatic animals with acoustic transmitters. *Methods Ecol. Evol.* 14 (10), 2514–2530. <https://doi.org/10.1111/2041-210X.14191>.
- Lennox, R.J., Aarestrup, K., Cooke, S.J., Cowley, P.D., Deng, Z.D., Fisk, A.T., Harcourt, R. G., Heupel, M., Hinch, S.G., Holland, K.N., 2017. Envisioning the future of aquatic animal tracking: technology, science, and application. *Bioscience* 67 (10), 884–896.
- Lennox, R.J., Eldøy, S.H., Dahlmo, L.S., Matley, J.K., Vollset, K.W., 2023b. Acoustic accelerometer transmitters and their growing relevance to aquatic science. *Movement Ecology* 11 (1), 45. <https://doi.org/10.1186/s40462-023-00403-3>.
- Lennox, R.J., Engler-Palma, C., Kowarski, K., Filous, A., Whitlock, R., Cooke, S.J., Auger-Méthé, M., 2019. Optimizing marine spatial plans with animal tracking data. *Can. J. Fish. Aquat. Sci.* 76 (3), 497–509. <https://doi.org/10.1139/cjfas-2017-0495>.
- Lennox, R.J., Whoriskey, F.G., Verhelst, P., Vandergoot, C.S., Soria, M., Reubens, J., Rechisky, E.L., Power, M., Murray, T., Mulder, I., Markham, J.L., Lowerre-Barbieri, S.K., Lindley, S.T., Knott, N.A., Kessel, S.T., Iverson, S., Huveneres, C., Heidemeyer, M., Harcourt, R., et al., 2024. Globally coordinated acoustic aquatic animal tracking reveals unexpected, ecologically important movements across oceans, lakes and rivers. *Ecography* 2024 (1), e06801. <https://doi.org/10.1111/ecog.06801>.
- Lingard, S.A., Bass, A.L., Cook, K.V., Fortier, M., Price, G.G., Hinch, S.G., 2023. Evaluating the influence of environmental and biological factors on migration behavior and residence duration of wild subyearling Chinook Salmon in a fjord estuary using miniature acoustic transmitters. *Trans. Am. Fish. Soc.* 152 (5), 610–631. <https://doi.org/10.1002/tafs.10429>.
- Lotze, H.K., 2005. Radical changes in the Wadden Sea fauna and flora over the last 2,000 years. *Helgol. Mar. Res.* 59 (1), 71–83. <https://doi.org/10.1007/s10152-004-0208-0>.
- Lotze, H.K., 2007. Rise and fall of fishing and marine resource use in the Wadden Sea, southern North Sea. *Fish. Res.* 87 (2–3), 208–218. <https://doi.org/10.1016/j.fishres.2006.12.009>.
- Lynch, S.D., Marce, B.J., Marshall, H.M., Bushnell, P.G., Bernal, D., Brill, R.W., 2017. The effects of pop-up satellite archival tags (PSATs) on the metabolic rate and swimming kinematics of juvenile sandbar shark *Carcharhinus plumbeus*. *Fish. Res.* 186, 205–215. <https://doi.org/10.1016/j.fishres.2016.08.013>.
- Matley, J.K., Klinard, N.V., Barbosa Martins, A.P., Aarestrup, K., Aspillaga, E., Cooke, S. J., Cowley, P.D., Heupel, M.R., Lowe, C.G., Lowerre-Barbieri, S.K., Mitamura, H., Moore, J.-S., Simpfendorfer, C.A., Stokesbury, M.J.W., Taylor, M.D., Thorstad, E.B., Vandergoot, C.S., Fisk, A.T., 2022a. Global trends in aquatic animal tracking with acoustic telemetry. *Trends Ecol. Evol.* 37 (1), 79–94. <https://doi.org/10.1016/j.tree.2021.09.001>.
- Matley, J.K., Klinard, N.V., Larocque, S.M., McLean, M.F., Brownscombe, J.W., Raby, G. D., Nguyen, V.M., Barbosa Martins, A.P., 2023. Making the most of aquatic animal tracking: a review of complementary methods to bolster acoustic telemetry. *Rev. Fish Biol. Fish.* 33 (1), 35–54. <https://doi.org/10.1007/s11160-022-09738-3>.
- Matley, J.K., Klinard, N.V., Martins, A.P.B., Aarestrup, K., Aspillaga, E., Cooke, S.J., Cowley, P.D., Heupel, M.R., Lowe, C.G., Lowerre-Barbieri, S.K., 2022b. Global trends in aquatic animal tracking with acoustic telemetry. *Trends Ecol. Evol.* 37 (1), 79–94.
- McAllister, J.D., Barnett, A., Lyle, J.M., Semmens, J.M., 2015. Examining the functional role of current area closures used for the conservation of an overexploited and highly mobile fishery species. *ICES (Int. Counc. Explor. Sea) J. Mar. Sci.* 72 (8), 2234–2244.
- McMillan, M.N., Huveneres, C., Semmens, J.M., Gillanders, B.M., 2019. Partial female migration and cool-water migration pathways in an overfished shark. *ICES (Int. Counc. Explor. Sea) J. Mar. Sci.* 76 (4), 1083–1093. <https://doi.org/10.1093/icesjms/fsy181>.
- Methling, C., Tudorache, C., Skov, P.V., Steffensen, J.F., 2011. Pop up satellite tags impair swimming performance and energetics of the European eel (*Anguilla anguilla*). *PLoS One* 6 (6), e20797. <https://doi.org/10.1371/journal.pone.0020797>.
- Musyl, M.K., Domeier, M.L., Nasby-Lucas, N., Brill, R.W., McNaughton, L.M., Swimmer, J.Y., Lutcavage, M.S., Wilson, S.G., Galuardi, B., Liddle, J.B., 2011. Performance of pop-up satellite archival tags. *Mar. Ecol. Prog. Ser.* 433, 1–28.
- Nathan, R., Monk, C.T., Arlinghaus, R., Adam, T., Alós, J., Assaf, M., Baktoft, H., Beardsworth, C.E., Bertram, M.G., Bijleveld, A.I., 2022. Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science* 375 (6582), eabg1780.
- Nemeth, R.S., Kadison, E., Jossart, J., Shivji, M., Wetherbee, B.M., Matley, J.K., 2023. Acoustic telemetry provides insights for improving conservation and management at a spawning aggregation site of the endangered Nassau grouper (*Epinephelus striatus*). *Front. Mar. Sci.* 10. <https://www.frontiersin.org/articles/10.3389/fmars.2023.1154689>.
- Nielsen, J., Estévez-Barcia, D., Post, S., Christensen, H.T., Retzel, A., Meire, L., Rigét, F., Strøm, J.F., Rikardsen, A., Hedeholm, R., 2023. Validation of pop-up satellite

- archival tags (PSATs) on Atlantic cod (*Gadus morhua*) in a Greenland fjord. *Fish. Res.* 266, 106782 <https://doi.org/10.1016/j.fishres.2023.106782>.
- Novak, A.J., Becker, S.L., Finn, J.T., Danylchuk, A.J., Pollock, C.G., Hillis-Starr, Z., Jordaan, A., 2020. Inferring residency and movement patterns of horse-eye jack *Caranx latus* in relation to a Caribbean marine protected area acoustic telemetry array. *Animal Biotelemetry* 8 (1), 12. <https://doi.org/10.1186/s40317-020-00199-8>.
- Økland, F., Thorstad, E.B., Westerberg, H., Aarestrup, K., Metcalfe, J.D., 2013. Development and testing of attachment methods for pop-up satellite archival transmitters in European eel. *Animal Biotelemetry* 1 (1), 3. <https://doi.org/10.1186/2050-3385-1-3>.
- Orrell, D.L., Hussey, N.E., 2022. Using the VEMCO Positioning System (VPS) to explore fine-scale movements of aquatic species: applications, analytical approaches and future directions. *Mar. Ecol. Prog. Ser.* 687, 195–216. <https://doi.org/10.3354/meps14003>.
- Orrell, D.L., Webber, D., Hussey, N.E., 2023. A standardised framework for the design and application of fine-scale acoustic tracking studies in aquatic environments. *Mar. Ecol. Prog. Ser.* 706, 125–151. <https://doi.org/10.3354/meps14254>.
- Palumbi, S.R., 2004. Marine reserves and ocean neighborhoods: the spatial scale of marine populations and their management. *Annu. Rev. Environ. Resour.* 29 (1), 31–68. <https://doi.org/10.1146/annurev.energy.29.062403.102254>.
- Pawson, M.G., Kelley, D.F., Pickett, G.D., 1987. The distribution and migrations of bass, *Dicentrarchus labrax* L., in waters around England and Wales as shown by tagging. *J. Mar. Biol. Assoc. U. K.* 67 (1), 183–217. <https://doi.org/10.1017/S0025315400026448>.
- Pawson, M.G., Pickett, G.D., Leballeur, J., Brown, M., Fritsch, M., 2007. Migrations, fishery interactions, and management units of sea bass (*Dicentrarchus labrax*) in Northwest Europe. *ICES (Int. Coun. Explor. Sea) J. Mar. Sci.* 64 (2), 332–345. <https://doi.org/10.1093/icesjms/fsi035>.
- Pedersen, M.W., 2007. *Hidden Markov Models for Geolocation of Fish* [Master's Thesis]. Technical University of Denmark, DTU, Lyngby, Denmark. DK-2800 Kgs.
- Pereira, T.J., Silva, A.F., De Almeida, P.R., Belo, A.F., Costa, J.L., Castro, N., Quintella, B. R., 2017. Assessing the size adequacy of a small no-take marine protected area (MPA) for Mediterranean moray and European conger. *Mar. Ecol. Prog. Ser.* 584, 213–227. <https://doi.org/10.3354/meps12379>.
- Perluka, R., Wiegmann, E.B., Jordans, R.W.L., Swart, L.M.T., 2006. *Opmatetechnieken waddenzee*. Rijkswaterstaat Adviesdienst Geo Informatie En ICT, Delft (AGI-2006-GPMP-004).
- Perry, A.L., Low, P.J., Ellis, J.R., Reynolds, J.D., 2005. Climate change and distribution shifts in marine fishes. *Science* 308 (5730), 1912–1915.
- Poiesz, S.S.H., Witte, J.I.J., van der Veer, H.W., 2020. Only a few key prey species fuel a temperate coastal fish food web. *Mar. Ecol. Prog. Ser.* 653 (Zijlstra 1972), 153–166. <https://doi.org/10.3354/meps13472>.
- Poiesz, S.S.H., Witte, J.I., van der Meer, M.T., Jager, Z., Soetaert, K.E., van der Heide, T., van der Veer, H.W., 2023. Stomach content and stable isotopes illustrate large spatial similarity in the Wadden Sea fish food-web structure. *Mar. Ecol. Prog. Ser.* 707, 57–76.
- Potter, I.C., Tweedley, J.R., Elliott, M., Whitfield, A.K., 2015. The ways in which fish use estuaries: a refinement and expansion of the guild approach. *Fish Fish.* 16 (2), 230–239.
- Queiroz, N., p Lima, F., Maia, A., a Ribeiro, P., p Correia, J., Santos, A.M., 2005. Movement of blue shark, *Prionace glauca*, in the north-east Atlantic based on mark-recapture data. *Marine Biological Association of the United Kingdom. Journal of the Marine Biological Association of the United Kingdom* 85 (5), 1107.
- Rauck, G., Zijlstra, J.J., 1978. On the nursery-aspects of the Waddensea for some commercial fishing species and possible long-term changes. *Rapp. P.-V. Reun. Cons. Int. Explor. Mer.* 172 (1), 266–275.
- Reubens, J., Aarestrup, K., Meyer, C., Moore, A., Okland, F., Afonso, P., 2021. Compatibility in acoustic telemetry. *Animal Biotelemetry* 9 (1), 1–6.
- Reubens, J., Hernandez, F., Deneudt, K., 2017. LifeWatch observatory—permanent acoustic receiver network in the Belgian part of the North sea. VLIZ Flanders Marine Institute - Platform for Marine Research. <https://www.vliz.be/en/imis?module=dataset&dataset=5843>.
- Reyier, E., Ahr, B., Iafraite, J., Scheidt, D., Lowers, R., Watwood, S., Back, B., 2023. Sharks associated with a large sand shoal complex: community insights from longline and acoustic telemetry surveys. *PLoS One* 18 (6), e0286664. <https://doi.org/10.1371/journal.pone.0286664>.
- Rodriguez-Dominguez, A., Connell, S.D., Leung, J.Y.S., Nagelkerken, I., 2019. Adaptive responses of fishes to climate change: feedback between physiology and behaviour. *Sci. Total Environ.* 692, 1242–1249. <https://doi.org/10.1016/j.scitotenv.2019.07.226>.
- Rous, A.M., Midwood, J.D., Gutowsky, L.F.G., Lapointe, N.W.R., Portiss, R., Sciscione, T., Wells, M.G., Doka, S.E., Cooke, S.J., 2017. Telemetry-determined habitat use informs multi-species habitat management in an urban harbour. *Environ. Manag.* 59 (1), 118–128. <https://doi.org/10.1007/s00267-016-0775-2>.
- Santana-Garcon, J., Newman, S.J., Langlois, T.J., Harvey, E.S., 2014. Effects of a spatial closure on highly mobile fish species: an assessment using pelagic stereo-BRUVs. *J. Exp. Mar. Biol. Ecol.* 460, 153–161. <https://doi.org/10.1016/j.jembe.2014.07.003>.
- Schaber, M., Gastauer, S., Cisewski, B., Hielscher, N., Janke, M., Peña, M., Sakinan, S., Thorburn, J., 2022. Extensive oceanic mesopelagic habitat use of a migratory continental shark species. *Sci. Rep.* 12 (1), 2047.
- Sheehan, E., Couturier, L., Hawkes, L., Hooper, T., Horton, T., Laurans, M., Muñoz, C., Reubens, J., Stamp, T., Witt, M., Woillez, M., 2021. FISHINTEL: fisheries Innovation for sustainable SHARED INTerchannEL resources (FISH INTEL). VLIZ Flanders Marine Institute - Platform for Marine Research. <https://www.vliz.be/en/imis?module=dataset&dataset=7881>.
- Sieben, C., Gascoigne, J., Nehls, G., Ramirez-Monsalve, P., Raakjaer, J., 2013. Sustainable Fisheries in the Trilateral Wadden Sea.
- Sims, D.W., Wearmouth, V.J., Southall, E.J., Hill, J.M., Moore, P., Rawlinson, K., Hutchinson, N., Budd, G.C., Righton, D., Metcalfe, J.D., 2006. Hunt warm, rest cool: bioenergetic strategy underlying diel vertical migration of a benthic shark. *J. Anim. Ecol.* 75 (1), 176–190.
- Skomal, G.B., Zeeman, S.I., Chisholm, J.H., Summers, E.L., Walsh, H.J., McMahon, K.W., Thorrold, S.R., 2009. Transequatorial migrations by basking sharks in the western Atlantic ocean. *Curr. Biol.* 19 (12), 1019–1022. <https://doi.org/10.1016/j.cub.2009.04.019>.
- Speed, C.W., Field, I.C., Meek, M.G., Bradshaw, C.J., 2010. Complexities of coastal shark movements and their implications for management. *Mar. Ecol. Prog. Ser.* 408, 275–293.
- Stevens, J.D., 1990. Further results from a tagging study of pelagic sharks in the north-east Atlantic. *J. Mar. Biol. Assoc. U. K.* 70 (4), 707–720. <https://doi.org/10.1017/S0025315400058999>.
- Svendsen, E., Fore, M., Økland, F., Gräns, A., Hedger, R.D., Alfredsen, J.A., Uglem, I., Rosten, C.M., Frank, K., Erikson, U., 2021. Heart rate and swimming activity as stress indicators for Atlantic salmon (*Salmo salar*). *Aquaculture* 531, 735804.
- SWIMWAY, 2019. *TRILATERAL WADDEN SEA SWIMWAY VISION ACTION PROGRAMME*, 1.1.
- Thorburn, J., Neat, F., Burrett, I., Henry, L.-A., Bailey, D.M., Jones, C.S., Noble, L.R., 2019. Ontogenetic variation in movements and depth use, and evidence of partial migration in a benthopelagic elasmobranch. *Frontiers in Ecology and Evolution* 7, 353.
- Thorstad, E.B., Todd, C.D., Uglem, I., Bjørn, P.A., Gargan, P.G., Vollset, K.W., Halttunen, E., Kälås, S., Berg, M., Finstad, B., 2016. Marine life of the sea trout. *Mar. Biol.* 163 (3), 1–19. <https://doi.org/10.1007/s00227-016-2820-3>.
- Thygesen, U.H., Pedersen, M.W., Madsen, H., 2009. Geolocating fish using hidden Markov models and data storage tags. *Tagging and Tracking of Marine Animals with Electronic Devices*, pp. 277–293.
- Tulp, I., Bolle, L., Chen, C., Daenhardt, A., Haslob, H., Jepsen, N., Van Leeuwen, A., Poiesz, S., Scholle, J., Vrooman, J., 2021. Quality status Report: trends in Wadden Sea fish. 15th International Scientific Wadden Sea Symposium.
- Tulp, I., Bolle, L.J., 2009. Trends in Wadden Sea Fish Fauna, Part II: Dutch Demersal Fish Survey (DFS). *IMARES*.
- Tulp, I., Bolle, L.J., Dänhardt, A., de Vries, P., Haslob, H., Jepsen, N., Scholle, J., van der Veer, H.W., 2017a. Fish. Wadden sea quality status Report 2017. Wadden Sea Ecosyst. 9. <http://cwss.www.de/TMAP/Qsr99/Qsr99.html>.
- Tulp, I., Bolle, L.J., Rijnsdorp, A.D., 2008. Signals from the shallows: in search of common patterns in long-term trends in Dutch estuarine and coastal fish. *J. Sea Res.* 60 (1–2), 54–73. <https://doi.org/10.1016/j.seares.2008.04.004>.
- Tulp, I., Bolle, L.J., Chen, C., Dänhardt, A., Haslob, H., Jepsen, N., van Leeuwen, A., Poiesz, S.S.H., Scholle, J., Vrooman, J., Vorberg, R., Walker, P., 2022. *Wadden Sea quality status report—fish* (Wadden Sea quality status Report). Common Wadden Sea Secretariat. <https://qsr.waddensea-worldheritage.org/reports/fish>.
- Tulp, I., Glorius, S., Rippen, A., Looije, D., Craeymeersch, J., 2020. Dose-response relationship between shrimp trawl fishery and the macrobenthic fauna community in the coastal zone and Wadden Sea. *J. Sea Res.* 156, 101829.
- Tulp, I., Van Der Veer, H.W., Walker, P., Van Walraven, L., Bolle, L.J., 2017b. Can guild- or site-specific contrasts in trends or phenology explain the changed role of the Dutch Wadden Sea for fish? *J. Sea Res.* 127, 150–163.
- UNESCO, 2023. *Operational Guidelines for the Implementation of the World Heritage Convention*, WHC.23/01.
- van Aken, H.M., 2008. Variability of the water temperature in the western Wadden Sea on tidal to centennial time scales. *J. Sea Res.* 60 (4), 227–234. <https://doi.org/10.1016/j.seares.2008.09.001>.
- van der Knaap, I., Slabbekoorn, H., Winter, H.V., Moens, T., Reubens, J., 2020. Evaluating Receiver Contributions to Acoustic Positional Telemetry: A Case Study on Atlantic Cod Around Wind Turbines in the North Sea.
- van der Meer, J., Witte, J.I., van der Veer, H.W., 1995. The suitability of a single intertidal fish trap for the assessment of long-term trends in fish and epibenthic invertebrate populations. *Environ. Monit. Assess.* 36 (2), 139–148. <https://doi.org/10.1007/BF00546786>.
- Van der Molen, J., Van Leeuwen, S.M., Govers, L.L., Van der Heide, T., Olf, H., 2021. Potential micro-plastics dispersal and accumulation in the North Sea, with application to the MSC Zoe incident. *Front. Mar. Sci.* 8, 607203.
- van der Veer, H.W., Dapper, R., Henderson, P.A., Jung, A.S., Philippart, C.J., Witte, J.I., Zuur, A.F., 2015. Changes over 50 years in fish fauna of a temperate coastal sea: degradation of trophic structure and nursery function. *Estuar. Coast Shelf Sci.* 155, 156–166.
- van der Veer, H.W., Koot, J., Aarts, G., Dekker, R., Diderich, W., Freitas, V., Witte, J.I., 2011. Long-term trends in juvenile flatfish indicate a dramatic reduction in nursery function of the Balgzand intertidal, Dutch Wadden Sea. *Mar. Ecol. Prog. Ser.* 434, 143–154.
- van der Veer, H.W., Tulp, I., Witte, J.I., Poiesz, S.S., Bolle, L.J., 2022. Changes in functioning of the largest coastal North Sea flatfish nursery, the Wadden Sea, over the past half century. *Mar. Ecol. Prog. Ser.* 693, 183–201.
- van Puijenbroek, P.J., Buijs, A.D., Kraak, M.H., Verdonchot, P.F., 2019. Species and river specific effects of river fragmentation on European anadromous fish species. *River Res. Appl.* 35 (1), 68–77.
- van Walraven, L., Dapper, R., Nauw, J.J., Tulp, I., Witte, J.I., van der Veer, H.W., 2017. Long-term patterns in fish phenology in the western Dutch Wadden Sea in relation to

- climate change. *J. Sea Res.* 127 (May 2016), 173–181. <https://doi.org/10.1016/j.seares.2017.04.001>.
- Verhelst, P., Buysse, D., Reubens, J., Pauwels, I., Aelterman, B., Van Hoey, S., Goethals, P., Coeck, J., Moens, T., Mouton, A., 2018. Downstream migration of European eel (*Anguilla anguilla* L.) in an anthropogenically regulated freshwater system: implications for management. *Fish. Res.* 199, 252–262. <https://doi.org/10.1016/j.fishres.2017.10.018>.
- Wada, T., Kamiyama, K., Mitamura, H., Arai, N., 2017. Horizontal movement and emigration of juvenile spotted halibut *Verasper variegatus* released in a shallow brackish lagoon: matsukawa-ura, northeastern Japan, revealed by acoustic telemetry. *Fish. Sci.* 83 (4), 573–585. <https://doi.org/10.1007/s12562-017-1099-8>.
- Walker, P., Howlett, G., Millner, R., 1997. Distribution, movement and stock structure of three ray species in the North Sea and eastern English Channel. *ICES (Int. Council. Explor. Sea) J. Mar. Sci.* 54 (5), 797–808.
- Walker, T.I., Rigby, C.L., Pacoureaux, N., Ellis, J.R., Kulka, D., Chiaramonte, G.E., Herman, K., 2020. Galeorhinus galeus: IUCN Red List of Threatened Species. *IUCN Red List of Threatened Species*, e.T39352A2907336. <https://dx.doi.org/10.2305/IUCN.UK.2020-2.RLTS.T39352A2907336.en>.
- Walsh, A., Reay, P., O'Halloran, J., Cahill, K., 1994. The growth of grey mullet in a rural and urbanized Irish estuary. *J. Fish. Biol.* 45 (5), 889–897.
- Watanabe, Y.Y., Papastamatiou, Y.P., 2023. Biologging and biotelemetry: tools for understanding the lives and environments of marine animals. *Annual Review of Animal Biosciences* 11 (1), 247–267. <https://doi.org/10.1146/annurev-animal-050322-073657>.
- Wright, G.V., Wright, R.M., Kemp, P.S., 2014. Impact of tide gates on the migration of juvenile sea trout, *Salmo trutta*. *Ecol. Eng.* 71, 615–622.
- Young, J.M., Bowers, M.E., Reyier, E.A., Morley, D., Ault, E.R., Pye, J.D., Gallagher, R.M., Ellis, R.D., 2020. The FACT Network: philosophy, evolution, and management of a collaborative coastal tracking network. *Marine and Coastal Fisheries* 12 (5), 258–271.