Geophysical Research Abstracts Vol. 13, EGU2011-9222, 2011 EGU General Assembly 2011 © Author(s) 2011



Ocean Chemistry reflected in Cold Water Coral Reef Occurrences

Wolf-Christian Dullo (1), Sascha Flögel (1), Andres Rüggeberg (1,2), and Furu Mienis (3) (1) IFM-GEOMAR, Paleooceanology, Kiel, Germany (cdullo@ifm-geomar.de, + 49 431 6002925), (2) Renard Centre of Marine Geology, Ghent University, Krijgslaan 281, S8, 9000 Gent, Belgium, (3) MARUM - Zentrum für Marine Umweltwissenschaften der Universität Bremen, Leobener Str., 28359 Breme

Physical and chemical parameters were measured in five different regions of the Northeast Atlantic with known occurrences of cold-water coral reefs and mounds and in the Mediterranean, where these corals form living carpets over existing morphologies. In this study we analyzed 282 bottom water samples regarding δ 13CDIC, δ 18O, and DIC. The hydrochemical data reveal characteristic patterns and differences for cold-water coral sites with living coral communities and ongoing reef and mound growth at the Irish and Norwegian sites. While the localities in the Mediterranean, in the Gulf of Cadiz, and off Mauritania show only patchy coral growth on mound-like reliefs and various substrates.

The analysis of $\delta 13C/\delta 18O$ reveals distinct clusters for the different regions and the respective bottom water masses bathing the corals. The analysis of relationships between salinity, temperature, $\delta 18O$, and especially between $\delta 13CDIC$ and DIC shows that DIC is a parameter with high sensitivity to the mixing of bottom water masses. It varies distinctively between sites with living reefs/mounds and sites with restricted patchy growth or dead corals. Results suggest that DIC and $\delta 13CDIC$ can provide additional insights into the mixing of bottom water masses.

Prolific cold-water coral growth forming giant biogenic structures plot into a narrow geochemical window characterized by a variation of δ 13CDIC between 0.45 and 0.79 %0 being associated with the water mass having a density of sigma-theta of 27.5 \pm 0.15 kg m-3.