

First records and new information on the associations of echinoderms with other phyla in the rocky reefs of northern Chocó, Colombian Pacific

María Juliana Vanegas González¹, Giomar Helena Borrero-Pérez¹

¹ Instituto de Investigaciones Marinas y Costeras José Benito vives de “Andreis” – INVEMAR. Calle 25 No. 2–55, Playa Salguero, Rodadero, Santa Marta, Colombia

Corresponding author: María Juliana Vanegas González (mjulianavanegas@gmail.com)

Academic editor: A. Martynov | Received 2 January 2019 | Accepted 3 January 2020 | Published 24 March 2020

<http://zoobank.org/0D22F959-F085-467E-96FC-C5C2ECA37AA3>

Citation: Vanegas González MJ, Borrero-Pérez GH (2020) First records and new information on the associations of echinoderms with other phyla in the rocky reefs of northern Chocó, Colombian Pacific. *ZooKeys* 921: 1–22. <https://doi.org/10.3897/zookeys.921.32802>

Abstract

Rocky reefs of the northern Colombian Pacific (Chocó) are diverse ecosystems that are poorly studied. Echinoderms are one of the principal groups in these ecosystems with associations to different species, including benthic organisms in which they live and other species that use them as hosts. These relationships include fishes, sponges, seaweeds, cnidarians, polychaetes, bryozoans, crustaceans, mollusks, and other echinoderms. For this area, 22 associations were registered, including commensalism, epibionts and parasitism. This work constitutes the first report for the associations of *Eucidaris thouarsii* with *Suberites* aff. *ficus*, *E. thouarsii* with *Ophiothela mirabilis*, and *Holothuria* (*Thymiosicia*) *impatiens* with *Encheliophis vermicularis*. Associations of *Pentaceraster cumingi* with *Zenopontonia soror*, and *Ophionereis annulata* with *Malmgreniella* cf. *variegata* are new records for Colombia. This work also expands the range of hosts previously described for *Ophiothela mirabilis* and expands the distribution of the association between *Diadema mexicanum* and *Echineulima* cf. *robusta*.

Keywords

Commensalism, epibionts, parasitism, Asteroidea, Echinoidea, Holothuroidea, Ophiuroidea, Riscuales

Introduction

Echinoderms are distributed in all coastal environments from tidal pools to rocky and coral reefs, in which they share space and refuge areas with members of their own phylum and other taxa (Sotelo-Casas and Rodríguez-Troncoso 2014). Due to this closeness, different types of interactions have been developed; within these associations are found the ones in which echinoderms depends on other organisms such as sponges and octocorals for protection against predators and for easy access to food (Henkel and Pawlik 2005, Marin et al. 2005). Others in which echinoderms interact with other organisms and the substrate, for example species that depend on the fixing cavities constructed by sea urchins such as *Echinometra lucunter lucunter* (Linnaeus, 1758) (Schoppe 1991). The cavities of this sea urchin, from the Caribbean Sea, are used by the brittle star *Ophiothrix synoecina* (Schoppe, 1996) which is obligated symbiont of *E. lucunter lucunter* (Schoppe 1996; Schoppe and Werding 1996). Finally, the mutualism occurring between different species of detritivorous sea cucumbers that share their inhabiting spaces and adopt different schedules for feeding and positions allowing other species to take advantage of the food (Rupert and Barnes 1996).

Relationships occur in all echinoderm classes; for example, the starfishes are frequently inhabited by symbionts of various taxonomic groups such as polychaetes, copepods, and mollusks (Jangoux 1990, Antokhina et al. 2012), with some species that are obligate symbionts (i.e., *Hololepidella millari*, *Doridicola echinasteris*) (Antokhina and Britayev 2012). For sea cucumbers, interactions with at least nine phyla have been described including diatoms, protozoans, flat worms (i.e., Xenacoelomorphos), polychaetes, mollusks, crustaceans, fish, and even other echinoderms (Jangoux 1990, Eeckhaut et al. 2004, Purcell et al. 2016). Related to sea urchins, different types of relationships have also been established, including commensalism with animals as the crab *Stenorhynchus debilis* (Smith, 1871) and a fish of the genus *Apogon* for which the sea urchin spines served as a refuge (Sotelo-Casas and Rodríguez-Troncoso 2014), and sponges that use the spines of sea urchins as an attachment substrate (Hétérier and De Ridder 2004, Aguirre et al. 2011). Although studies on crinoids are very limited, the association and dependence of many of the myxostomid species (Annelida) with this group has been recorded; approximately 100 of the 150 species of myxostomid currently described live above or within crinoids during their adult stage (Summers and Rouse 2014). For brittle stars, interactions with different kind of organisms have been reported, by having different adaptations in color and the brittle stars *Ophionereis* behavior to simulate the host, this is the case of the polynoid *Harmothroe lunulata*, *Ophionereis reticulata* and *O. annulata* (Millott 1953, Granja-Fernández et al. 2013, Gómez-Maduro and Díaz-Díaz 2017). Finally, many associations with benthic organisms such as sponges have been described in relation to ophiuroids (Bejarano et al. 2004, Marin et al. 2005), in some cases they depend specifically on other organisms for their development (Pardo et al. 1988).

The most studied marine groups related to their interactions with echinoderms are Mollusca and Crustacea. Mainly bivalve and gastropod relationships have been recognized (Caullery 1952), with more than 30 species of prosobranchs recorded as parasites of echinoderms (Caullery 1952), especially echinoids (Hyman 1955). It has also been found that most groups of crustaceans have some type of association with echinoderms, for example cirripedes are considered endo- and ectosymbionts of species such as *Dendrogaster* spp. (Caullery 1952), isopods have both obligate and non-specific relationships with all the five classes of echinoderms, and shrimps include species that are obligated commensals of some echinoderms (Ross 1983).

Relationships between echinoderms and different types of organisms have been widely registered throughout the world, but these are poorly studied and understood in Colombian waters and in the entire Tropical Eastern Pacific. To this end, relationships of the echinoderms with other phyla were recorded during a project developed to characterize the biodiversity of the rocky reefs of Chocó Norte in Colombia.

Materials and methods

Individuals were collected during two expeditions carried out on April and October 2016 in the northern area of the Colombian Pacific, Chocó Department (Figure 1), between Cabo Corrientes in the south (5°29'N, 77°32'W) and Cabo Marzo in the north (6°49'N, 77°41'W). Those expeditions were performed to increase the knowledge in terms of diversity and distribution of the marine biodiversity of the rocky reefs of the area. In the Tropical Eastern Pacific, rocky reefs (called “Riscales” and “Morros” in Chocó), represent habitats for many invertebrate species, including corals, sea fans, and fishes, making them important and productive ecosystems for artisanal fishing (Rubio and Angulo 2003). Some of the rocky reefs are submerged, rising above the level of low tide or forming small permanently emerged islets (Díaz et al. 2016).

Samplings were made using scuba diving, and a direct and random collection of echinoderms was made by sweeping each station looking for all potential habitats in different areas of the rocky reef, at all depths starting at 25 m and finishing at 5 m, the top of the submerged reefs. All field information was recorded for each individual collected, including depth and habitat. The collected echinoderms were relaxed using magnesium chloride dissolved with sea water ($MgCl_2 \cdot 6H_2O$) and fixed in 96% ethanol. Specimens were morphologically reviewed, photographed, and identified using stereoscope and microscopes. In order to correctly identify sea cucumbers, body wall ossicles were examined.

During the study, all the associated organisms were photographed in field, separated from the echinoderm when possible, and identified by expert taxonomists of each of the groups. The photographs taken in field were reviewed in order to complement the information of the associations, especially those that included octocorals to confirm the distribution range of the associations. Posterior samplings in the same




Figure 1. A Locations along northern Colombian Pacific (Chocó) where associations of echinoderms with other phyla were collected. Locations names from north to south P.ERO: Piedra de Eroito, FOCA: la foca, P.ROD: Piedra de Rodrigo, MINA: La mina, PARN: Parguera norte, VIUD: la Viuda, CHIC: Chicocora, P.ZAP: Piedra Zapata, P.NOR: Punta norte, MORN: Morromico norte, JURU: Jurubidá, P.ORI: Punta Orión, P.BON: Piedra bonita, PARS: Parguera sur, PARU: Punta Arusí, P.JAI: Piedra de Jairo, ROÑO: Roñosa, AMGR: Amargal, COLO: Coló. The line between FOCA and FARO represents the proximity between both stations.

area, carried out during 2017 and 2018 and focused mainly on octocoral biodiversity of the same locations, allowed us to expand the information on these relationships and are included in this work too.




Figure 1. Continued: **B** Locations where associations between *O. mirabilis* and octocorals were registered. Location names from north to south: P.ERO: Piedra de Eroitto, FOCA: la foca, P.PROD: Piedra de Rodrigo, MINA: La mina, PARN: Parguera norte, VIUD: la Viuda, CHIC: Chicocora, P.ZAP: Piedra Zapata, P.NOR: Punta norte, MORN: Morromico norte, JURU: Jurubidá, P.ORI: Punta Orión, P.BON: Piedra bonita, PARS: Parguera sur, PARU: Punta Arusí, P.JAI: Piedra de Jairo, ROÑO: Roñosa, AMGR: Amargal, COLO: Coló. The lines among several locations represent the proximity between them.

All the collected material was deposited in the biological collections from the Museo de Historia Natural Marina de Colombia (MHNMC) – Makuriwa of INVEMAR.

Results and discussion

A total of 22 relationships were registered between echinoderms and other organisms in the rocky reefs of the northern Chocó in the Colombian Pacific. These relationships include fish, sponges, cnidarians, polychaetes, bryozoans, crustaceans, and mollusks (Table 1). All the individuals were identified to the lowest possible taxonomic level, excepting some specimens that were registered only in photographs.

Pentaceraster cumingi (Gray, 1840) – *Zenopontonia soror* (Nobili, 1904)

Figure 2

Material: one specimen of *Pentaceraster cumingi* (INV EQU4283) was collected with two shrimps (*Zenopontonia soror*) located in the oral part (Figure 2). Shrimps were orange, the same color that the ambulacral feet of the sea star. This relationship was recorded in Piedra de Rodrigo (P.ROD) (Figure 1).

Table 1. Relationships between echinoderms and other marine groups found in the rocky reefs of northern Chocó, Colombian Pacific. Key for the relationships C: Commensalism, E: Epibiont, P: Parasitism. Key for Micro-habitats 1: Exposed in the Rocky reef, 2: Exposed in other living organisms, 3: Under rocks in contact with sand, 4: Partially exposed in the Rocky reef, 5: Partially exposed between rocks.

Echinoderm	Other organisms	Relationship	Micro-habitat	Depth (m)	Figure	
<i>Pentaceraster cumingi</i>	<i>Zenopontonia soror</i>	C	1	19	Figure 2	
<i>Ophiotbela mirabilis</i>	<i>Euclidaris thourarsii</i>	E	2	7	Figure 3C	
	<i>Leptogorgia alba</i>	C	2	3-19	Figure 3A	
	<i>Leptogorgia</i> sp.	C	2	3-19		
	<i>Pacifigorgia adamsi</i>	C	2	7-19	Figure 3H	
	<i>Pacifigorgia bayeri</i>	C	2	5-19	Figure 3C	
	<i>Pacifigorgia eximia</i>	C	2	7-19	Figure 3E	
	<i>Pacifigorgia irene</i>	C	2	7-19	Figure 3B	
	<i>Pacifigorgia rubicunda</i>	C	2	15	Figure 3F	
	<i>Pacifigorgia stenobrochis</i>	C	2	7-19		
	<i>Pacifigorgia sculpta</i>	C	2	25		
	<i>Samogorgia</i> sp.	C	2	7-19		
	<i>Muricea squarrosa</i>	C	2	7		
		Cnidarians	E	2	5	
		Seaweeds	E	2		
<i>Ophionereis annulata</i>	<i>Malmgreniella</i> cf. <i>variegata</i>	C	3	15	Figure 4	
<i>Euclidaris thourarsii</i>	Bryozoans	E	4		Figure 5	
	<i>Suberites</i> aff. <i>fuscus</i>	E	5	9	Figure 5B	
	Barnacles	E	4		Figure 5	
	<i>Ophiotbela mirabilis</i>	E	4		Figure 5C	
<i>Diadema mexicanum</i>	<i>Echineulima</i> cf. <i>robusta</i>	P	4	19	Figure 6	
<i>Holothuria impatiens</i>	<i>Encheliophis vermicularis</i>	P	3	7	Figure 7	




Figure 2. **A** *Pentaceraster cumingi* aboral view **B** *P. cumingi* oral view **C** *Zenopontonia soror*. The shrimp lost the color after fixing in 96% ethanol. Scale bars: 100 mm (**A, B**), 2 mm (**C**).

Starfish are frequently inhabited by several symbiotic animals (Jangoux 1990). Within the genus *Pentaceraster* the association with *Zenopontonia soror* has been commonly reported world-wide. This commensal shrimp belongs to the family Palaemonidae and is known for being a specialized and obligate symbiont of starfishes (Antokhina and Britayev 2012). *Zenopontonia soror* was initially described in the Red Sea (Nobili 1904), but Edmonson (1935) and Gordon (1939) reported its association with some species of asteroids of the Indo-Pacific, and finally Bruce (1976) reported it in Australia associated with *Plectaster decanus* (Müller & Troschel, 1843) and in Pacific of Panama with *Pentaceraster cumingi* (as *Oreaster occidentalis*) (Bruce 1976), both sea stars being common in reefs. *Zenopontonia soror* is currently reported associated with 21 asteroids species, including five species of the genus *Pentaceraster* (Antokhina and Britayev 2012). This is the first report of the association between *Zenopontonia soror* and *Pentaceraster cumingi* for the Colombian Pacific.

***Ophiothela mirabilis* Verrill, 1867 and several symbiosis associations**

Figure 3

Material: *Ophiothela mirabilis* was registered on different hosts: *Eucidaris thouarsii* (INV EQU4218) (Figure 5C), cnidarians (Figure 3I), seaweeds, and octocorals: *Leptogorgia alba* (Duchassaing y Michelotti, 1864) (INV EQU4251) (Figure 3A, D), *Leptogorgia* sp., *Pacifigorgia rubicunda* Breedy y Guzman, 2003 (Figure 3B), *Pacifigorgia eximia* (Verrill, 1868) (Figure 3C), *Pacifigorgia irene* Bayer, 1951 (Figure 3E), *Pacifigorgia stenobrochis* (Valenciennes, 1846) (Figure 3F), *Pacifigorgia sculpta* Breedy & Guzman, 2004 (Figure 3G), *Pacifigorgia bayeri* Breedy, 2001 (Figure 3H), *Pacifigorgia adamsi* (Verrill, 1868), *Samogorgia* sp. and *Muricea squarrosa* Verrill, 1869. One cnidarian morphotype, possibly a hydrozoan, and the seaweeds could not be identified because they were not collected. *Ophiothela mirabilis*' relationship with octocorals was observed in 16 stations during April (2016) including Chicocora (CHIC), la Foca (FOCA), la Mina (MINA), Parguera (PARN), Piedra de Eroito (P.ERO), Punta norte (P.NOR), Piedra de Rodrigo (P.ROD), la viuda (VIUD), Coló (COLO), Morromico norte (MORN), Jurubidá (JURU), la Roñosa (ROÑO), Amargal (AMGR), Parguera norte (PARN2), Piedra bonita (P.BON) and Punta Orión (P.ORI). In October (2016) no octocoral was registered with this brittle star; however, the relationship was observed in subsequent samplings developed in October 2017 (not included on this work) and 2018.

Brittle stars are usually associated with organisms such as sponges, cnidarians, sea urchins, and even algae which provide shelter and a place to feed (Bejarano et al. 2004, Hendler et al. 2012). Some authors like Neira and Cantera (2005) and Lawley et al. (2018) (as *Ophiothela* cf. *mirabilis*) have found that *O. mirabilis* has a preference for octocoral hosts. However, Mantelatto et al. (2016) indicated that in Brazil, area where it is invasive, *O. mirabilis* is an opportunistic and generalist species in relation to host species selection, finding it related to 20 different invertebrates (i.e., *Dysidea etheria* – sponge, *Isostichopus badiionotus* – sea cucumber). The relationship between this brittle star and its hosts is considered as commensalism; however, it has been suggested that there may be a negative effect to the host due to the high densities that *O. mirabilis* can present (Mantelatto et al. 2016). Those negative effects may not be directly caused by its feeding on the host, but some authors suggest that it may be related to structural damage due to the increase in the weight of commensals that the host must support, and in the case of the cnidarians with the possible interruption of the extension of the polyps making it difficult to feed and compromising the ability of the octocoral to obtain nutrients (Mantelatto et al. 2016, Thé de Araújo et al. 2018). In northern Chocó, high densities of *O. mirabilis* on the octocorals were observed during April 2016 and in subsequent samplings in 2017 and 2018, but further studies are needed to elucidate the possible negative effect of this relationship to the octocorals in this area. For the Colombian Pacific, *O. mirabilis* has been reported to be associated mainly with the octocoral *Leptogorgia alba* (Cantera et al. 1987, Pardo et al. 1988, Neira and




Figure 3. *Ophiothela mirabilis* on different cnidarians **A** *Leptogorgia alba* **B** *Pacifigorgia rubicunda* **C** *P. eximia* **D** *Ophiothela mirabilis* collected on *L. alba* **E** *O. mirabilis* collected on *Pacifigorgia irene* **F** *P. stenobrochis* **G** *P. sculpta* **H** *P. bayeri* **I** Cnidarian. Scale bars: 2 mm (**D, E**).

Cantera 2005). However, the results presented here expand its range of hosts to include other octocoral species: *Pacifigorgia rubicunda*, *P. adamsi*, *P. eximia*, *P. irene*, *P. bayeri*, *P. sculpta*, *P. stenobrochis*, *Muricea squarrosa*, *Samogorgia* sp., and *Leptogorgia* sp., leafy algae, and *E. thouarsii*.

Several individuals of *O. mirabilis* were found using *E. thouarsii* spines as a fixing substrate, but there are no studies of *O. mirabilis* as a sea urchin epibiont in the Tropical Eastern Pacific; however, for the Brazilian Caribbean, where is an invasive species, *O. mirabilis*, has been reported living in high densities in *Echinometra lucunter* spines (Mantelatto et al. 2016). This constitutes the first report of the relationship between *O. mirabilis* and *E. thouarsii*, expanding the range of hosts for *O. mirabilis*.

Beside the groups mentioned above, others organisms have also been reported as *O. mirabilis* hosts, especially in the Mexican Pacific where association with scleractinian corals and sponges have been reported (Granja-Fernández and López-Pérez 2011), and in the Caribbean (Brazil), others groups were found including sponges, ascidians, and bryozoans (Mantelatto et al. 2016). So far there are no other reports of *O. mirabilis* living in seaweeds. Although the species of this group was not identified to species level, this would be the first record of the relationship between *O. mirabilis* and seaweeds.

Ophionereis annulata* (Le Conte, 1851) – *Malmgreniella* cf. *variegata

Figure 4

Material: two polychaetes of the family Polynoidae identified as *Malmgreniella* cf. *variegata* were found living on specimens of *Ophionereis annulata* collected in Chicocora (CHIC) (INV EQU4370) and Parguera (PARN) (INV EQU4208) (Figure 1). The polychaetes had a similar coloration pattern to the dorsal side of the ophiuroid arms (Figure 4).

The association between polychaetes of the family Polynoidae with brittle stars belonging to the genus *Ophionereis* has been reported for the Pacific Ocean and the Caribbean Sea. *Malmgreniella variegata* (Treadwell, 1917) distributed mainly in the western Atlantic, including the Caribbean Sea (GBIF 2019, OBIS 2018) has a relationship with *Ophionereis reticulata* (Say, 1825) (Hendler et al. 1995). This association has been well described for Brazil (Santa-Isabel et al. 1996) and recently for Venezuela (Gómez-Maduro and Díaz-Díaz 2017). *Malmgreniella variegata* has a similar coloration and a banded pattern as the ophiuroid arms, which allows it to go unnoticed on the host (Pettibone 1993, Gómez-Maduro and Díaz-Díaz 2017). *Malmgreniella* cf. *variegata* has been reported mainly associated with the aboral part of the disk of the ophiuroid, using it as a refuge (Granja-Fernández et al. 2013). Although *M. variegata* is distributed in the Atlantic Ocean, some authors such as Pettibone (1993) reported the species living on *Ophionereis annulata* in the Gulf of Panama. In the Mexican Pacific, Granja-Fernandez et al. (2013) described a similar commensal relationship between *O. annulata* and a polynoid polychaete, which, after reviewing its morphology, was identified as *M. cf. variegata*, due to differences in the color pattern of the elytra and the notochaetae from *M. variegata* from the Caribbean Sea. After examination of the specimens collected in northern Chocó we found all the taxonomic characteristics described by Salazar-Silva (2009) for the Atlantic polychaete *Malmgreniella variegata* except for the color pattern of the elytra, similar to the results described by Granja-Fernandez et al. (2013). The identification of these polychaetes needs to be revised using additional evidence, such as molecular analysis. This is the first report of the presence of *M. cf. variegata* and its association with *O. annulata* in the Colombian Pacific.



Figure 4. *Malmgreniella* cf. *variegata* (left) and *Ophionereis annulata* (right). Scale bar: 5 mm.

***Eucidaris thouarsii* (L. Agassiz y Desor, 1846) – Bryozoa, Cirripedia, *Suberites* aff. *ficus*, and *Ophiothela mirabilis* Verrill, 1867**

Figure 5

Material: Four types of epibionts were found inhabiting on *Eucidaris thouarsii* spines in different stations (Figure 1), including cirripede barnacles (INV EQU4218), several species of bryozoans (INV EQU4528, INV EQU4293, INV EQU4299) (Figure 5A), and the sponge *Suberites* aff. *ficus* (INV EQU4301) in Piedra Zapata (PZAP) (Figure 5B). The ophiuroid *O. mirabilis* was collected on the sea urchin (INV EQU4218) in Coló (COLO) and Piedra de Eroito (PERO) (Figure 5C, D).

Only one sea urchin morphotype belonging to the family Cidaridae was found in the rocky reefs in northern Chocó, identified as *Eucidaris thouarsii*, mostly due to its thick spines (Figure 5). Different organisms use *E. thouarsii* spines as an attachment substrate, including sessile (bryozoans, sponges) and mobile animals (*O. mirabilis*). There are not many published papers related to the epibionts species of sea urchins; however, it has been reported that sponges, especially species belonging to the genus *Clathria*, grow on the spines and the discs of *E. thouarsii* (David et al. 2009, Aguirre et al. 2011). For other species of sea urchin, different associations have been reported: for *H. asteriscus* more than 20 species of epifauna have been identified, including nine



Figure 5. Epibionts on *Eucidaris thouarsii* spines **A** Bryozoans **B** *Suberites* aff. *ficus* **C** *Ophiothela mirabilis* **D** close up to *O. mirabilis* collected from *E. thouarsii* spine.

species of polychaetes, five species of bryozoans, three species of mollusks, three species of crustaceans, two sponge species, and a single species each of protozoan, cnidarian, nematode, and echinoderm (Salazar and López 1983). The reason sea urchins of the family Cidaridae are commonly used as hosts is that they have spines made of muscle and collagen in their basal part which allows the settlement of epibiont fauna, unlike other echinoids which have antifouling compounds (David et al. 2009, Aguirre et al. 2011). In this work, the sponge that settled on *E. thouarsii* spines was identified as *Suberites* aff. *ficus* (Figure 5B) because of the spicules. *Suberites ficus* (Johnston, 1842) is the given name for a complex of species with megascleres, tylostyles, microscleres, strongyles, and oxeas, originally from the North Atlantic Ocean but with two records on the eastern Pacific (Dickinson 1945, Bakus and Green 1987). The tylostyles and

microscleres from the Colombian Pacific specimen are smaller (tylostyles: 90–221 × 3–8 µm; microscleres: 17–42 × 1–3 µm) than the ones reported in specimens from other areas of the Eastern Pacific, such as lower California, Mexico (tylostyles: 340 × 10 µm; microscleres: 18–36 × 1–3 µm; De Laubenfels 1932) and south California (tylostyles: 120–680 × 1–11 µm; microscleres: 20–48 × 1–2.5 µm; Bakus and Green 1987). Until now, *Clathria* was the only sponge genus reported growing on *E. thouarsii* (Aguirre et al. 2011), but *S. ficus* has been reported living on other organisms such as gastropod shells (Bakus and Green 1987). This report constitutes the first record of *S. aff. ficus* and its association with *E. thouarsii* for the Colombian Pacific. At least five unidentified bryozoan species (Figure 5A) have been observed on the spines of *E. thouarsii*, which were collected in parts of the study area (INV EQU4528, INV EQU4293, INV EQU4299).

The phylum Crustacea is another of the groups reportedly associated with sea urchins; different species, especially of crabs and shrimps, have been found living between the spines and, in some cases, attached to the spines (Macía and Robinson 2012, Britayev et al. 2013), using the spines as protection and on occasion benefiting from the sea urchin's grazing (Morton 1988). However, there is not much information about other crustaceans, especially about barnacles symbiotic with sea urchins. Some species that have been reported living in sea urchin spines are *Balanus trigonus* and *Paralepas percarinata* (as *Heteralepas percarinata*) (Werner 1967, Aldous 1970).

***Diadema mexicanum* A. Agassiz, 1863 – *Echineulima* cf. *robusta* (Pease, 1860)**

Figure 6

Material: Three specimens of *Echineulima* cf. *robusta* were found on the oral portion of one specimen of the sea urchin *Diadema mexicanum* (INV EQU4292), and five organisms were found on another specimen (INV EQU4530). Both sea urchins were juveniles (INV EQU4292, 18.6 mm test diameter; INV EQU4530, 25.4 mm test diameter). The specimens were collected in Piedra de Jairo (PJAI) (INV EQU4292) and La Foca (FOCA) (INV EQU4530) (Figure 1). The gastropods were white in color (Figure 6).

The associations between echinoderms and gastropods have been widely recorded for the family Eulimidae, including 750 species (Warén 1983, Jangoux 1984). For the species *Centrostephanus coronatus* it was reported that they have an ectoparasitic relationship with gastropods of the genus *Echineulima* (Jangoux 1984, Cantera and Neira 1987). These gastropods parasitize the interambulacral zone on the oral side of the test, using a proboscis that penetrates the skeleton to feed on the hemal fluid of its host (Cantera and Neira 1987). This relationship is considered as parasitism because of the damage caused by the gastropod to the sea urchin; among the negative effects that have been recorded for these gastropods, in addition to the fact that they feed on the sea urchin, they produce lesions by their grip, causing dermal swellings and even distor-




Figure 6. **A** *Echineulima* cf. *robusta* **B** *Diadema mexicanum* (INV EQU4292) with three specimens of *Echineulima* cf. *robusta*. Scale bars: 2 mm (**A**), 4 mm (**B**).

tion in the skeleton (Jangoux 1984). In the present work two individuals of *Diadema mexicanum* were collected with several individuals of *Echineulima* cf. *robusta* present on the oral side, close to the mouth (Figure 6). *Echineulima robusta* has already been registered as parasite of the sea urchins of the same family in the Indo-Pacific (Warén 1983); however, for the Colombian Pacific Cantera and Neira (1987) reported *Echineulima* sp. parasitizing *C. coronatus* for Gorgona Island. Other species of *Echineulima* have been reported parasitizing sea urchins in other geographical areas, such as in the Gulf of California (Mexico) and Taboga islands (Panama) where the association between *D. mexicanum* and *Echineulima mittrei* (Petit de la Saussaye, 1851) has been observed (Lützen and Nielsen 1975, Alvarado et al. 2015), and in Hawaii where *Echineulima thanuumi* (Pilsbry, 1921) has been reported parasitizing sea urchins of the genus *Echinotrix* (Cantera and Neira 1987). Olivares-González (1986) found that there exists a preference of the parasite *E. mittrei* for sea urchins with test diameters of approximately 20 mm to 39 mm, although the gastropod was found in sizes ranging from 20 mm to 60 mm. These preferences are related to the energy expenditure; in other words, sea urchins smaller than 20 mm use their energy for body growth while larger animals use it to produce gametic material, which is the presumably the preferred source of food for the gastropod.

Three specimens of *Echineulima* cf. *robusta* were found on one of the sea urchins of *Diadema mexicanum*, and five on the other; however, in this last one the gastropods were found in the oral side of the test separated in what looked like two different groups: the first group had three individuals, one of them bigger than the other two, and the other group had two individuals similar in size. Olivares-González (1986) found that sea urchins with more than one individual have them organized in groups of one female and one or two males; the males sharing a single impression on the test but with different apertures. Although the association between gastropods of the genus *Echineulima* with other sea urchins has already been reported for Gorgona Island (Cantera and Neira 1987), this report constitutes the first report for the Colombian




Figure 7. **A** *Holothuria impatiens* (right) and *Encheliophis vermicularis* (left) **B** Close up of *E. vermicularis*. Scale bars: 8 mm (**A**), 5 mm (**B**).

Pacific of *Echineulima* cf. *robusta* and its relationship with *Diadema mexicanum*. Additionally, in this work we are reporting that *Echineulima* cf. *robusta* can parasitize smaller sea urchins (18.6 mm test diameter) than previously reported.

***Holothuria* (*Thymiosycia*) *impatiens* (Forskål, 1775) – *Encheliophis vermicularis* Müller, 1842**

Figure 7

Material: One specimen of *Encheliophis vermicularis* (Figure 7) was found inside of one specimen of *Holothuria* (*Thymiosycia*) *impatiens* (INV EQU4240), collected in Punta Arusí (PARU) (Figure 1). The fish measured 58.76 mm length.

The family Holothuridae serves as host to several species of pearl fish of the family Carapidae (Luciano et al. 2002). In this family it has been found that the genera *Onuxodon* and *Carapus* act as commensals of the sea cucumbers but feed outside their hosts (Parmentier and Vandewalle 2003, Parmentier et al. 2016), while the genus *Encheliophis* behaves like a parasite, staying and feeding on internal tissues, causing small internal wounds and reducing the gonadal tissues of the host (Parmentier and Das 2004, Parmentier et al. 2006). In this work a specimen of *Encheliophis vermicularis* was found inside of *Holothuria* (*Thymiosycia*) *impatiens*. These fish usually dwell in the respiratory or digestive cavities of sea cucumbers, using them as protection and as a source of food (Trott 1981, Luciano et al. 2002, González-Wangüemert et al. 2014), and in some cases where pearl fish are found in pairs, the sea cucumber also serves as a breeding site (González-Wangüemert et al. 2014). The relationship between the genus *Encheliophis* with sea cucumbers has been reported for species such as *Holothuria* (*Microthele*) *fuscopunctata* Jaeger, 1833, *Holothuria* (*Holothuria*) *tubulosa* Gmelin, 1791, and *Isostichopus fuscus* (Ludwig, 1875), amongst others (Parmentier and Vandewalle 2005, Purcell et al. 2016). Regarding *E. vermicularis*, some authors found that it has a very specific rela-

relationship with *Holothuria* (*Mertensiothuria*) *leucospilota* (Brandt, 1835) (Miyazaki et al. 2014), *Holothuria* (*Mertensiothuria*) *hilla* Lesson, 1830 (James 1995) and with lower incidence with *Holothuria* (*Halodeima*) *atra* Jaeger, 1833 (Smith 1964). Although this fish species has already been reported for Gorgona Island as *Encheliophis hancocki* (Reid, 1940) (Reid 1940, Orrell and Hollowell 2018), a synonymized name of *E. vermicularis* (Froese and Pauly 2018), this work expands its distribution to the northern Chocó.

Conclusions

Despite echinoderms constituting an important group with representatives present in all marine ecosystems, and many different kinds of interaction with other phyla have been described, the information on their associations for the Colombian Pacific was limited. This work has helped to increase the knowledge on echinoderms and their associations with other groups including fishes, mollusks, polychaetes, cnidarians, and sponges from the rocky reefs of the Colombian Pacific and in general from the Tropical Eastern Pacific.

Ethics approval and consent to participate

These species do not appear as evaluated in the IUCN Red List. Riscales project is part of the Biodiversity and Marine Ecosystems research program of the Instituto de Investigaciones Marinas y Costeras of Colombia (INVEMAR), which belongs to the Ministry of Environment and Sustainable Development of Colombia, in accordance with Law 99 of 1993 (Article 18), which does not require Permission of Collection of specimens for Scientific Research according to the decree 1076 of 2015 (Chapter 8, Section 1, Article 2.2.2.8.1.2., Paragraph 1).

Acknowledgements

We would like to thank Katherine Mejia (INVEMAR) for the assistance with octocoral identification and for the photographs used in Figure 3F–H; Adriana Osorno (INVEMAR) and Jaime Cantera (Universidad del Valle) for the identification of the mollusk *Echineulima* cf. *robusta*; Adibe Cardenas (INVEMAR) for the polychaete *Malmgreniella* cf. *variegata* identification; Luis Chasqui and Andrea Polanco (INVEMAR) for the fish *Encheliopsis vermicularis* identification; Sven Zea (Universidad Nacional de Colombia) and Nicoll Lizarazo (Universidad Jorge Tadeo Lozano) for the sponge *Suberites* aff. *ficus* identification; and Bibian Martínez (INVEMAR) and Juan Felipe Lazarus (Universidad del Valle) for the identification of the crustacean *Zenopontonia soror*. We thank Luis Chasqui and David Alonso for the project development; Natalia Rincon, Bibian Martinez and all the researchers of Riscales project for helping during sampling. We thank Lina Vasques (LABSIS-INVEMAR) for making the map used in

Figure 1. The funding came from the “Caracterización biótica de los arrecifes rocosos (riscales y morros) en el Pacífico norte chocoano” (Riscales) project, under the agreement with the Ministry of Environment and Sustainable Development of Colombia, Code: BPIN: 2014011000405. Invemar contribution No. CTRB – 1255.

References

- Aguirre LK, Hooker Y, Willens P, Hajdu E (2011) A new *Clathria* (Demospongiae, Microcionidae) from Peru occurring on rocky substrates as well as epibiontic on *Eucidaris thouarsii* sea urchins. *Zootaxa* 3085: 41–54. <https://www.researchgate.net/publication/287740944>
- Aldous J (1970) An annotated checklist of Hawaiian barnacles (Class Crustacea, subclass Cirripedia) with notes on their nomenclature, habitats and Hawaiian localities. Hawaii Institute of Marine Biology, Technical report No. 19, 133 pp.
- Alvarado JJ, Reyes-Bonilla H, Benitez-Villalobos F (2015) *Diadema mexicanum*, erizo de mar clave en los arrecifes coralinos del Pacífico Tropical Oriental: lo que sabemos y perspectivas futuras (Diadematoidea: Diadematoidea). *Revista de Biología Tropical* 63(2): 135–157. <https://doi.org/10.15517/rbt.v63i2.23140>
- Antokhina TI, Britayev TA (2012) Sea stars and their macrosymbionts in the Bay of Nhatrang, Southern Vietnam. *Paleontological Journal* 46(8): 894–908. <https://doi.org/10.1134/S0031030112080023>
- Antokhina TI, Savinkin OV, Britayev TA (2012) Asteroidea of Vietnam with some notes on their symbionts. In: Britayev TA, Pavlov DS (Eds) Benthic fauna of the bay of Nhatrang southern Vietnam, Vol 2, 406–438.
- Bakus GJ, Green D (1987) The distribution of marine sponges collected from the 1976–1978 Bureau of land management Southern California bight program. *Bulletin of Southern California Academic Science* 86(2): 57–88.
- Bejarano S, Zea S, Díaz JM (2004) Esponjas y otros microhábitats de ofiuros (Ophiuroidea: Echinodermata) en ambientes arrecifales del archipiélago de San Bernardo (Caribe colombiano). *Boletín de Investigaciones Marinas y Costeras* 33: 29–47. <https://doi.org/10.25268/bimc.invemar.2004.33.0.246>
- Britayev TA, Bratova OA, Dgebuadze PY (2013) Symbiotic assemblage associated with the tropical sea urchin, *Salmacis bicolor* (Echinoidea: Temnopleuridae) in the An Thoi archipiélago, Vietnam, *Symbiosis* 61(3): 155–161. <https://doi.org/10.1007/s13199-013-0263-x>
- Bruce AJ (1976) *Periclimenes soror* Nobili, a pontonin shrimp new to the American fauna, with observations on its Indo-West Pacific distribution. *Tethys* 8(4): 299–306.
- Cantera JR, Neira R (1987) Primer registro del género *Echineulima* Luzten y Nilsen, 1975 (Eulimidae) gasterópodo parásito de erizos del mar en la isla Gorgona Colombia. *Anales del Instituto de Investigaciones Marinas Punta Betín* 17: 87–93. <https://doi.org/10.25268/bimc.invemar.1987.17.0.457>
- Cantera JR, von Prael H, Neira R (1987) Moluscos, crustáceos y equinodermos asociados a la gorgonia *Lophogorgia alga* (Duchassaing y Michelotti, 1864), en la isla Gorgona, Colombia. *Boletín Ecotropical* 17: 331–353.

- Caso ME (1979) Ordenes Cidarioidea y Aulodonta. Publ. Especiales Centro de Ciencias del Mar y Limnol. UNAM. <http://biblioweb.tic.unam.mx/cienciasdelmar/especiales/1978-1/articulo453.html>
- Caullery M (1952) Parasitism and symbiosis. Sidgwich and Jackson Ltd., London, 358 pp.
- Clark HL (1948) A report of the echini of the warmer eastern Pacific, based on the collection of the *Velero III*. Allan Hancock Pacific Expeditions 8: ii-xii. 225–352. [pl. 36: fig. 4, pl. 37] <https://www.biodiversitylibrary.org/page/5512986#page/259/mode/1up>
- David B, Stock SR, Carlo F, Hétériér V, de Ridder C (2009) Microstructures of Antarctic cidaroid spines: diversity of shapes and ectosymbiont attachments. *Marine Biology* 156: 1559–1572. <https://doi.org/10.1007/s00227-009-1192-3>
- De Laubenfels MW (1935) Some sponges of lower California (México). The American museum of natural history novitates No. 779: 1–14.
- Díaz JM, Guillot L, Velandia MC (Eds) (2016) La pesca artesanal en la costa norte del Pacífico Colombiano: un horizonte ambivalente. Fundación MarViva, Bogotá, 158 pp. http://marviva.net/sites/default/files/documentos/la_pesca_artesanal_final_web.pdf
- Dickinson MG (1945) Sponges of the Gulf of California. Allan Hancock Pacific Expeditions. 11: 1–251.
- Eeckhaut I, Parmentier E, Becker P, Gomez da Silva S, Jangoux M (2004) Parasites and biotic diseases in field and cultivated sea cucumber. In: Lovatelli A, Conand C, Purcell S, Uthicke S, Hamel JF, Mercier A (Eds) *Advances in sea cucumber aquaculture and management*. FAO Fisheries technical paper 463, 311–326. <http://hdl.handle.net/2268/12699>
- Edmonson C (1935) New and rare Polynesian Crustacea. Occasional Papers Bishop Museum 10(24): 1–38.
- Froese R, Pauly D (Eds) (2018) FishBase. *Encheliophis vermicularis* Müller, World Register of Marine Species. 1842. <http://www.marinespecies.org/aphia.php?p=taxdetails&id=279269> [accessed 13 August 2018]
- GBIF (2019) *Malmgreniella variegata* (Treadwell, 1917). GBIF Backbone Taxonomy. Checklist dataset available online. <https://doi.org/10.15468/39omei> [accessed via GBIF.org on 22 Oct 2019]
- Gómez-Maduro MC, Díaz-Díaz O (2017) Association between the polynoid *Malmgreniella variegata* (Polychaeta: Polynoidae) and *Ophionereis reticulata* (Ophiuroidea: Ophionereididae) first record to the southeastern Caribbean. *Revista de Biología Tropical* 65(1): s85–s91. <https://doi.org/10.15517/rbt.v65i1-1.31669>
- González-Wangüemert M, Maggi C, Valente S, Martínez-Garrido J, Rodrigues NB (2014) *Parastichopus regalis*- The main host of *Carapus acus* in temperate waters of the Mediterranean Sea and northeastern Atlantic Ocean. SPC Beche-de-mer information Bulletin 34: 38–42.
- Gordon I (1939) Redescription of *Periclimenes soror* Nobili (Crustacea, Decapoda). *Annals and Magazine of Natural History* 11(4): 395–400. <https://doi.org/10.1080/00222933908527002>
- Granja-Fernández R, López-Pérez RA (2011) Nuevos registros de ofiuroides (Echinodermata: Ophiuroidea) para localidades de Zihuatanejo (Guerrero) y Puerto Escondido (Oaxaca), Pacífico mexicano. *Revista Mexicana de Biodiversidad* 82: 1320–1323. <https://doi.org/10.22201/ib.20078706e.2011.4.675>

- Granja-Fernández R, Hernández-Moreno P, Bastida-Zavala R (2013) First record of the association between *Malmgreniella* cf. *variegata* (Polychaeta, Polynoidae) and *Ophionereis annulata* (Echinodermata, Ophionereididae) in the Mexican Pacific. *Symbiosis*. 60: 85–90. <https://doi.org/10.1007/s13199-013-0239-x>
- Hendler G, Miller JE, Pawson DL, Kier PM (1995) Sea Stars, Sea Urchins and Allies: Echinoderms of Florida and the Caribbean. Smithsonian Institution, Washington, 195 pp.
- Hendler G, Migotto AE, Ventura CRR, Wilk L (2012) Epizoic *Ophiothela* brittle stars have invaded the Atlantic. *Coral Reefs* 31: 1005. <https://doi.org/10.1007/s00338-012-0936-6>
- Henkel TP, Pawlik JR (2005) Habitat use by sponge-dwelling brittlestars. *Marine Biology* 146: 301–313. <https://doi.org/10.1007/s00227-004-1448-x>
- Hétérier V, De Ridder C (2004) Comparative biodiversity of ectosymbionts in two Antarctic cidaroid echinoids, *Ctenocidaris spinosa* and *Rhynchocidaris triplopora*. In: Heinzeller T, Nebelsick JH (Eds) Proceedings of the 11th I.E.C., Echinoderms, 201–205. <https://doi.org/10.1201/9780203970881.ch35>
- Hyman LH (1955) The invertebrates: Echinodermata. The coelomate. Bilateria, vol IV. Mc Graw Hill, New York, 763 pp.
- James DB (1995) Animal associations in echinoderms. *Journal of the Marine Biological Associations of India* 37(1–2): 272–276.
- Jangoux M (1984) Diseases of echinoderms. *Helgolander Meeresunters* 37: 207–216. <https://doi.org/10.1007/BF01989305>
- Jangoux M (1990) Diseases of Echinodermata. In: Kinne O (Ed.) Diseases of Marine Animals. Biologische. Anstalt Helgoland. 3:439–567.
- Lawley JW, Carvahal A, Faria E, Lindner A (2018) Occurrence of the non-indigenous brittle star *Ophiothela* cf. *mirabilis* Verrill, 1867 (Echinodermata, Ophiuroidea) in natural and anthropogenic habitats off Santa Catarina, Brazil. *Check list* 14(2): 453–459. <https://doi.org/10.15560/14.2.453>
- Luciano B, Lyman A, McMillan S, Nickels A (2002) The symbiotic relationship between sea cucumbers (Holothuriidae) and pearlfish (Carapidae). UCSC BioE 159 Moorea class papers. 13 pp. <http://bio.classes.ucsc.edu/bioe159/Moorea%202006/Old%20Student%20Papers/Pearl%20Fishes%20-%202002.pdf>
- Lützen J, Nielsen K (1975) Contribution to the anatomy and biology of *Echineulima* n. g. (Prosobranchia: Eulimidae), parasitic on sea urchins. *Videnska-belige Meddlelser Dansk Naturhistorisk Forening*. 138: 171–199.
- Macía S, Robinson MP (2012) Reproductive Pattern in the Caridean Shrimp *Gnathophyllodes mineri* Schmitt (Gnathophyllidae), a Symbiont of Sea Urchins. *Journal of Crustacean Biology* 32(5): 727–732. <https://doi.org/10.1163/193724012X638491>
- Mantelatto MC, Vidon LF, Silveira RB, Menegola C, Da Rocha RM, Creed JC (2016) Host species of the non-indigenous brittle star *Ophiothela mirabilis* (Echinodermata: Ophiuroidea): an invasive generalist in Brazil? *Marine Biodiversity records*. 9:8. <https://doi.org/10.1186/s41200-016-0013-x>
- Marin IN, Anker A, Britayev TA, Palmer AR (2005) Symbiosis between the Alpheid Shrimp, *Athanas ornithorhynchus* Banner and Banner, 1973 (Crustacea: Decapoda), and the Brittle Star *Macrophiothrix longipeda* (Lamarck, 1816) (Echinodermata: Ophiuroidea). *Zoological*

- Studies. 44(2): 234–241. Available online at: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.652.8869&rep=rep1&type=pdf>
- Millott N (1953) A remarkable association between *Ophionereis reticulata* (Say) and *Harmothoe lunulata* (Delle Chiaje). Bulletin of Marine Science 3(2): 96–99.
- Miyazaki S, Ichiba T, Reimer JD, Tanaka J (2014) Chemoattraction of the pearfish *Encheliophis vermicularis* to the sea cucumber *Holothuria leucospilota*. Chemoecology, 24(3): 121–126. <https://doi.org/10.1007/s00049-014-0152-7>
- Morton B (1988) Partnerships in the sea: Hong Kong's marine symbioses. Hong Kong University Press, Hong Kong, 121 pp.
- Neira R, Cantera JR (2005) Composición taxonómica y distribución de las asociaciones de equinodermos en los ecosistemas litorales del Pacífico Colombiano. Revista de Biología Tropical 53(3): 195–206.
- Nobili G (1904) Diagnoses préliminaires de vingt-huit espèces nouvelles de Stomatopodes et Décapodes Macroures de la Mer Rouge. Bulletin du Muséum d'Histoire naturelle 10: 228–238.
- OBIS (2018) Distribution records of *Malmgreniella variegata* (Treadwell, 1917). Ocean biogeographic information system. Intergovernmental oceanographic commission of UNESCO. www.Iobis.org [accessed 18 July 2018]
- Olivares-González E (1986) Algunos aspectos sobre la biología del erizo de espina larga *Diadema mexicanum* (Echinoidea: Echinodermata). I – Periodos reproductivos y II – Parasitismo. Dissertation. Universidad autónoma de Baja California sur, México, 68 pp.
- Orrell T, Hollowell T (2018) NMNH Extant Specimen Records. Version 1.19. National Museum of Natural History, Smithsonian Institution. Occurrence dataset. <https://www.gbif.org/occurrence/1319593860> [Accessed on: 13 August 2018]
- Padilla-Pérez MS, Rodríguez-Troncoso AP, Sotelo-Casas C, Cupul-Magaña AL (2017) Echinoderms of the Islas Marietas National Park: General aspects, importance and visual identification as a tool for their protection. Áreas Naturales Protegidas Scripta 3(2): 51–92. <https://doi.org/10.18242/anpscripta.2017.03.03.02.0003>
- Pardo R, Neira R, Cantera JR (1988) Taxonomía y ecología de equinodermos litorales del Pacífico Colombiano. Memorias VI Seminario Nacional de Ciencias y Tecnologías del mar, 115–120.
- Parmentier E, Das K (2004) Commensal vs. Parasitic relationship between Carapini fish and their host: some further insight through $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements. Journal of Experimental Marine Biology Ecology 310(1): 47–58. <https://doi.org/10.1016/j.jembe.2004.03.019>
- Parmentier E, Vandewalle P (2003) Morphological adaptations of pearlfish (Carapidae) to their various habitats. In: Val AL, Kapoor BG (Eds) Fish adaptations. Enfield: Science Publishers 261–276. <http://darwin.biology.utah.edu/PiqueArticles/Morphological%20Adaptations%20of%20Pearlfish.pdf>
- Parmentier E, Vandewalle P (2005) Further insight on carapid-holothuroid relationships. Marine Biology 146: 455–465. <https://doi.org/10.1007/s00227-004-1467-7>
- Parmentier E, Mercier A, Hamel JF (2006) New host and geographical distribution for the pearlfish *Carapus mourlani* (Carapidae) with a discussion on its biology. Copeia, 122–128. [https://doi.org/10.1643/0045-8511\(2006\)006\[0122:NHAGDF\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2006)006[0122:NHAGDF]2.0.CO;2)

- Parmentier E, Lanterbecq D, Eeckhaut I (2016) From commensalism to parasitism in Carapidae (Ophiidiiformes): heterochronic modes of development. *PeerJ*, 4: e1786. <https://doi.org/10.7717/peerj.1786>
- Pettibone MH (1993) Scaled polychaetes (Polynoidea) associated with ophiuroids and other invertebrates and review species referred to *Malmgrenia* McIntosh and replaced by *Malmgreniella* Hartman, with descriptions of new taxa. *Smithsonian Contributions to Zoology* 583: 1–92. <https://doi.org/10.5479/si.00810282.538>
- Purcell SW, Conand C, Uthicke S, Byrne M (2016) Ecological roles of exploited sea cucumbers. *Oceanography and marine biology: an Annual Revision*, 54: 367–386. <https://doi.org/10.1201/9781315368597-8>
- Reid ED (1940) A new genus and species of pearl fish, family Carapidae, from off Gorgona Island, Colombia 398. In: Markle DF, Olney J E (1990) Systematics of the pearlfishes (Pisces: Carapidae). *Bulletin of Marine Science* 47(2): 269–410.
- Ross DM (1983) Symbiotic relations. 163–214. In: Vernberg FJ, Vernberg WB (Eds) *Behavior and ecology Vol 7*. Academic Press, New York, 336 pp.
- Rubio EA, Angulo JA (2003) Peces coralinos del Pacífico coralino. Unidad de artes gráficas, Facultad de ciencias, Universidad del Valle, Cali, 317 pp.
- Rupert EE, Barnes RD (1996) *Zoología de los invertebrados*. 6ta edición. McGraw-Hill Interamericana, México, 1114 pp.
- Salazar SI, López IG (1983) Estudio preliminar sobre la epifauna de *Herperocidaris asteriscus* (Echinodermata: Echinoidea). *Ciencias Marinas* 9(2): 109–119. <https://doi.org/10.7773/cm.v9i2.421>
- Salazar-Silva P (2009) Polinoide Kinberg, 1856 39: 449–471. In: de León-González JA, Bastida-Zavala JR, Carrera-Parra LF, García-Garza ME, Peña-Rivera A, Salazar-Vallejo SI, Solís Weiss V (Eds) *Poliquetos (Annelida: Polychaeta) de México y América Tropical*. Universidad Autónoma de Nuevo León, Monterrey, Mexico, 737 pp.
- Santa-Isabel LM, Pinto WR, Souza OF (1996) Associação entre *Ophionereis reticulata* (say) (Ophiuroidea, Ophionereididae) e *Malmgreniella variegata* (Treadwell) (Polychaeta, Polynoidea) das praias de Salvador, Bahia, Brasil. *Revista Brasileira de zoologia* 13(1): 137–142. <https://doi.org/10.1590/S0101-81751996000100013>
- Schoppe S (1991) *Echinometra lucunter* (Linnaeus) (Echinoidea, Echinometridae) as host complex association in the Caribbean Sea. *Helgol Meeresunters* 45(3): 373–379. <https://doi.org/10.1007/BF02365526>
- Schoppe S (1996) *Ophiothrix synoecina*, new species (Echinodermata: Ophiuroidea) from the coast of Colombia. *Bulletin of Marine Science* 58(2): 429–437.
- Schoppe S, Werding B (1996) The boreholes of the sea urchin genus *Echinometra* (Echinodermata: Echinoidea: Echinometridae) as microhabitat in tropical South America. *Marine Ecology* 17:181–186. <https://doi.org/10.1111/j.1439-0485.1996.tb00499.x>
- Smith CL (1964) Some Pearlfishes from Guam, with notes on their ecology. *Pacific science* 18: 34–30.
- Sotelo-Casas RC, Rodríguez-Troncoso AP (2014) Los equinodermos: invertebrados esenciales dentro de la estructura de la comunidad coralina. In: Cifuentes JL, Cupul FG (Eds) *Temas sobre investigaciones costeras*. Universidad de Guadalajara, 56–81.

- Summers MM, Rouse GW (2014) Phylogeny of Myzostomida (Annelida) and their relationships with echinoderm hosts. *Evolutionary Biology* 14: 170. <https://doi.org/10.1186/s12862-014-0170-7>
- Thé de Araújo J, Oliveira M, Matthews-Cascon H, Correia FA (2018) The invasive brittle star *Ophiothela mirabilis* Verrill, 1867 (Echinodermata, Ophiuroidea) in the southwestern Atlantic: filling gaps of distribution, with comments on an octocoral host. *Latin American Journal of Aquatic Research* 46(5): 1123–1127. <https://doi.org/10.3856/vol46-issue5-fulltext-25>
- Trott L (1981) A general review of the pearlfish (Pisces: Carapidae). *Bulletin of marine science* 31(3): 623–629.
- Warén A (1983) A generic revision of the family Eulimidae. *Journal of Molluscan Studies* 13: 1–96. https://doi.org/10.1093/mollus/49.Supplement_13.1
- Werner WE (1967) The distribution and ecology of the barnacle *Balanus trigonus*. *Bulletin of Marine Science* 17(1): 64–84.