

numbers surviving. Overall, under climate change conditions juveniles have increased growth rates leading to larger size at the end of the first growth season yet reduced survival.

Work related to ICES via WGIPEM (the Working Group on Integrative, Physical-biological and Ecosystem Modelling).

## (20) Marine plastics: aligning national research and monitoring with international guidelines

**Bavo De Witte<sup>1</sup>, Karien De Cauwer<sup>2</sup>, Lisa Devriese<sup>3</sup>, Kris Hostens<sup>1</sup>**

<sup>1</sup> Flanders Research Institute for Agriculture, Fisheries and Food, Ankerstraat 1, 8400 Oostende, Belgium

<sup>2</sup> Royal Belgian Institute of Natural Sciences, Vautierstraat 29, 1000 Brussel, Belgium

<sup>3</sup> Flanders Marine Institute, InnovOcean site, Wandelaarkaai 7, 8400 Oostende, Belgium

The project Marine Plastics, funded by the European Maritime and Fisheries Fund (EFMZV), focuses on the presence of litter at the Belgian fisheries areas and the Belgian Part of the North Sea (BPNS), comprising both macro- and microlitter. Within the part on macrolitter, the main goals are to assess the presence of seabed litter at the Belgian fisheries areas, to identify impact of specific anthropogenic sources and to detect time trends in litter pollution. The part on microlitter concerns the identification of microplastics in seafood and the initiation of a Belgian monitoring approach for microplastics in seawater and sediment of the BPNS, in accordance with MSFD descriptor 10.

Many standardization issues remain. Macrolitter data collection and reporting differs between EU countries, the use of different gears for macrolitter sampling highly affects the extraction efficiency, there is a lack of standardization of macrolitter data assessment. The same range of issues, and even more, have to be clarified before a harmonized approach for microlitter monitoring and evaluation is reached. Which methods should be applied for microplastics sampling, separation, detection, identification and reporting? What size range and classes should be considered? What quality measures should be taken and how can we align international monitoring laboratories?

The interaction with the ICES working group on marine litter and microplastics (WGML) is of primary importance for the Marine Plastics project. WGML data assessments on international macrolitter datasets, collected within the International Bottom Trawl Surveys (IBTS) and Beam Trawl Surveys (BTS), influence the Marine Plastics project assessment approach, as similar statistical methods can be used. Vice versa, Belgian data has a unique value since marine litter is collected with the same fishing gear on a large area within the Belgian BTS. Besides the BTS, a second unique dataset is obtained within Belgian environmental monitoring at the BPNS. The latter has a large sampling density in an area with many human activities.

For microplastic research, the interaction with ICES WGML is also essential, as method alignment, standardization and quality assurance for monitoring purposes are key issues of this working group. Currently, methods for microlitter monitoring are not yet fixed and it will be a big challenge to align protocols between different EU member states in order to reach comparable results for MSFD descriptor 10. The active role of Belgian researchers at ICES WGML ensures that the Marine Plastic project is in accordance with newest guidelines and state-of-the-art protocols. Finally, the Marine Plastics project will provide the first integrated environmental monitoring assessment for microplastics in Belgium. This will be the necessary knowledge base for follow-up processes and actions by policy makers, the blue economy and the marine experts.

Work related to ICES via WGML (the Working Group on Marine Litter).

## (21) Long-term changes in demersal fish abundance and distribution in the Belgian part of the North Sea

**Jolien Buyse<sup>1</sup>, Steven Degraer<sup>2</sup>, Annelies De Backer<sup>1</sup> and Kris Hostens<sup>1</sup>**

<sup>1</sup> Flanders Research Institute for Agriculture, Fisheries and Food, Ankerstraat 1, 8400 Oostende, Belgium

<sup>2</sup> Operational Directorate Natural Environment, Royal Belgian Institute of Natural Sciences, Brussels, Belgium

Email: jolien.buyse@ilvo.vlaanderen.be

Climate change and other anthropological influences such as fisheries have long-term effects on fish abundance and distribution. To study the combined effects of these stressors on demersal fish species in the Belgian part of the North Sea,

autumn beam trawl survey data from 1985-2018 were analysed. Common trends in fish density over time were identified for the most important species using complementary multivariate techniques and linked to environmental variables. Further, changes in abundances of singular species were analysed using univariate linear models, which can explain patterns over time by the addition of explanatory variables to the models. Lastly, changes in length over time were also modelled for commercially interesting species. Using such a combination of different methods and data gives a good general overview of the most important drivers of local fish abundances linked to climatic and anthropological stressors. Information about such drivers are key for better understanding the marine environment and thereby influencing policy in terms of fisheries management and climate change mitigation.

Work related to ICES via WGNSSK (Working Group on the Assessment of Demersal Stocks in the North Sea and Skagerrak).

## (22) Tributyltin: an aggressive bottom-up stressor in a marine multistressor environment. A quality status report

**Koen Parmentier**,<sup>1,2</sup> Yves Verhaegen,<sup>1,3</sup> Bavo De Witte,<sup>1</sup> Daan Delbare,<sup>1</sup> Stefan Hoffman,<sup>1</sup> Patrick Roose,<sup>2</sup> Ketil Hylland,<sup>4</sup> Thierry Burgeot,<sup>5</sup> Guy Smaghe,<sup>3</sup> Kris Cooreman<sup>1</sup>

<sup>1</sup> Institute of Agriculture, Fisheries and Food Research, Animal Sciences, Ostend, Belgium

<sup>2</sup> Operational Directorate Natural Environment, Royal Belgian Institute of Natural Sciences, Ostend, Belgium

<sup>3</sup> Ghent University, Laboratory of Agrozoology, Faculty of Bioscience Engineering, Ghent, Belgium

<sup>4</sup> University of Oslo, Department of Biosciences, Oslo, Norway.

<sup>5</sup> Institut Français de Recherche pour l'Exploitation de la Mer, Unit of Biogeochemistry and Ecotoxicology, Nantes, France

The restrictions and the concerted action of the global ban on the use and presence of tributyltin (TBT) in marine applications to protect ecosystems in the marine environment in 2008 was mainly based on the economic impact on shellfish industries and the dramatic extinction of local mollusc populations in the past. In contrast to the vast datasets on effects on molluscs, the knowledge on impacts on species from other taxa remained in the uncertain until almost two decades ago. The assumption on a long-term TBT-mediated pernicious metabolic bottom-up regulation of the crustacean *Crangon crangon* population was provoked by the outcome of an EU-project 'Sources, Consumer Exposure and Risks of Organotin Contamination in Seafood.' This work reports high TBT body burdens in *C. crangon* in 2003, at the start of the transition period to the global ban. Experimental research on the TBT impact in *C. crangon* focused on agonistic interference with natural ecdysteroid hormones at the metabolic pathways regulating growth and reproduction and the biogeochemical distribution of the chemical. Metabolic, topical and population-relevant biological endpoints in *C. crangon* and other crustaceans are evaluated in relation to the temporal and spatial trends on TBT's occurrence and distribution in the field during and after the introduction of the tributyltin restrictions and endocrine-related incidents. Arguments are forwarded to relate the German Bight incident on growth and reproduction failure in the *C. crangon* population, despite the lack of direct evidence, to the pernicious impact of tributyltin in 1990/91 and previous years. The extreme occurrence of TBT in *C. crangon* from other parts of the southern North Sea and evidence on the high body burdens as dose metrics of exposure also feeds the suspicion on detrimental impacts in those areas. We further demonstrate the complexity of distinguishing and assessing the individual roles of unrelated stressors on a population in an integrated evaluation at the ecosystem level.

The Marine Chemistry Working Group (MCWG) is chaired by Koen Parmentier (RBINS). This group got considerable input from the Working Group on Biological Effects of Contaminants (WGBEC) and to a lesser extent from Working Group on Crangon Fisheries and Life History (WGCAN).

## (23) Towards open science products for ecosystem science

Lennert Schepers<sup>1</sup>, Lennert Tyberghein<sup>1</sup>

<sup>1</sup> Flanders Marine Institute (VLIZ), Wandelaarkaai 7, 8400 Oostende, Belgium

E-mail: lennert.schepers@vliz.be, lennert.tyberghein@vliz.be

Ecosystem science needs to integrate a variety of (biological) data sources and to use state-of-the-art methods to improve the knowledge of complex marine ecosystems. In this talk, we demonstrate how the Data Centre of Flanders Marine Institute (VLIZ) is organizing an open science data flow from collecting data to the development of biological products on Essential Ocean Variables (EOVs) that serves ecosystem assessments.