



Royal Netherlands Institute for Sea Research

This is a pre-copyedited, author-produced version of an article accepted for publication, following peer review.

Sorokin, D.Y.; Elcheninov, A.G.; Khizhniak, T.V.; Koenen, M.; Bale, N.J.; Sinninghe Damsté, J.S.; Kublanov, I.V. (2022). *Natronocalculus amylovorans* gen. nov., sp. nov., and *Natranaeroarchaeum aerophilus* sp. nov., dominant culturable amylolytic natronoarchaea from hypersaline soda lakes in southwestern Siberia. *Syst. Appl. Microbiol.* 45(4): 126336. DOI : 10.1016/j.syapm.2022.126336

Published version: <https://dx.doi.org/10.1016/j.syapm.2022.126336>

NIOZ Repository: <http://imis.nioz.nl/imis.php?module=ref&refid=353088>

[Article begins on next page]

The NIOZ Repository gives free access to the digital collection of the work of the Royal Netherlands Institute for Sea Research. This archive is managed according to the principles of the [Open Access Movement](#), and the [Open Archive Initiative](#). Each publication should be cited to its original source - please use the reference as presented.

When using parts of, or whole publications in your own work, permission from the author(s) or copyright holder(s) is always needed.

2 ***Natronocalculus amylovorans* gen. nov., sp. nov., and *Natranaeroarchaeum***
3 ***aerophilus* sp. nov., dominant culturable amyolytic natronoarchaea from**
4 **hypersaline soda lakes in southwestern Siberia.**

5
6
7 Dimitry Y. Sorokin^{a,b*}, Alexander G. Elcheninov^a, Tatjana V. Khizhniak^a, Michel Koenen^c, Nicole
8 J. Bale^c, Jaap S. Sinninghe Damsté^c and Ilya V. Kublanov^a

9
10 ^a*Winogradsky Institute of Microbiology, Research Centre of Biotechnology, Russian Academy of Sciences,*
11 *Moscow, Russia*

12 ^b*Department of Biotechnology, Delft University of Technology, The Netherlands*

13 ^c*NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and*
14 *Biogeochemistry, Texel, The Netherlands*

15
16 *Author for correspondence:

17 D.Y. Sorokin; e-mail: soroc@inmi.ru; d.sorokin@tudelft.nl

18
19
20
21
22
23
24
25
26
27 Running title: *Natronocalculus amylovorans* gen. nov., sp. nov. and *Natranaeroarchaeum*
28 *aerophilus* sp. nov.

29
30 The draft genome sequences of strains AArc-St1-1^T and AArc-St2^T are deposited in the GenBank
31 under the numbers JAKRVY000000000 and JAKRVX000000000, respectively

32

33 **Abstract**

34 Several pure cultures of alkaliphilic haloarchaea were enriched and isolated from hypersaline
35 soda lakes in southwestern Siberia using amylopectin and fructans as substrates. Phylogenomic
36 analysis placed the isolates into two distinct groups within the class *Halobacteria*. Four isolates
37 forming group 1 were closely related to a recently described *Natronaeroarchaeum sulfidigenes* and
38 the other three strains forming group 2 represent a novel genus-level phylogenetic lineage. All
39 isolates are saccharolytic archaea growing with various starch-like alpha-glucans including soluble
40 starch, amylopectin, dextrin, glycogen, pullulane and cyclodextrin. In addition, group 1 can also
41 use levan while group 2 - inulin (plant storage beta-fructans). Group 1 strains can also grow
42 anaerobically with either glucose or maltose using elemental sulfur as the electron acceptor. Both
43 groups are moderately alkaliphilic with a pH range for growth from 7.2 to 9.3 (optimum between
44 8.0-8.8) and low Mg-demanding extreme halophiles growing optimally at 4 M total Na⁺. The major
45 respiratory menaquinone is MK-8:8 and the core biphytanyl lipids are dominated by archaeol (C₂₀-
46 C₂₀) and a less abundant extended archaeol (C₂₀-C₂₅) with PG and PGP-Me as polar groups. The
47 four isolates of group 1 are suggested to be classified into a new species as *Natronaeroarchaeum*
48 *aerophilus* sp. nov. (type strain AArc-St1-1^T=JCM 32519^T=UQM 41458^T). The three isolates of
49 group 2 are proposed to form a new genus and species for which the name *Natronocalculus*
50 *amylovorans* gen. nov., sp. nov. is suggested (type strain AArc-St2^T=JCM 32475^T=UQM 41459^T).

51
52 Key words: hypersaline soda lakes, natronoarchaea, amylolytic, starch, fructans,

53 *Natronaeroarchaeum*, *Halobacteria*

54

55

56

57 **Introduction**

58 Most of the known species of aerobic extremely halophilic and haloalkaliphilic euryarchaea
59 (natronoarchaea) were enriched and isolated from hypersaline alkaline lakes on unspecific media
60 containing rich soluble organic substrates, such as peptone, yeast extract or simple sugars [1-5].

61 Yet, these extremophilic, organoheterotrophic archaea are definitely more important in organic
62 matter mineralization, in particular organic polymers, in hypersaline systems than is widely
63 recognized. For example, a test for amylolytic activity with soluble starch is included into the
64 minimal standards for taxonomy characterization of haloarchaea [6] but starch is rarely used for
65 targeted isolation of amylolytic haloarchaea which would not only hydrolyze the polymer but also
66 able to utilize it as growth substrate. And this situation is also true for other polysaccharides. So
67 far, only a few examples of haloarchaea specialized on utilization of recalcitrant polysaccharides
68 have been described in pure culture, which is particularly true for natronoarchaea living in alkaline
69 hypersaline (soda) lakes. Our recent targeted enrichments from such lakes using insoluble forms of
70 cellulose and chitin resulted in isolation of several groups of natronoarchaea highly specialized on
71 utilization of either various forms of cellulose and xylan (genera *Natronobiforma* and
72 *Natronolimnobius*) or chitin (genus *Natrarchaeobius*) [7-9]. Following further in this direction, we
73 used insoluble starch (amylopectin) or beta-fructans as substrates to enrich for amylolytic or
74 fructanolytic natronoarchaea from hypersaline soda lakes. So far, we are aware only of a single
75 amylolytic natronarchaeon, *Natronococcus amylolyticus*, specifically isolated from the hypersaline
76 soda lake Magadi in Kenya using starch as a growth substrate [10-11], while no natronoarchaea,
77 growing on fructans are currently known.

78
79 Here we describe phenotypic, phylogenetic and genomic properties of two novel taxa of amylolytic
80 natronoarchaea enriched from hypersaline soda lakes in southwestern Siberia, which specialized on
81 utilizing various alpha-glucan and beta-fructan polysaccharides as growth substrates.

82 **Materials and Methods**

83

84 *Enrichment and cultivation conditions*

85 The sources of inocula were mixed surface (0-3 cm deep) aerated sediments and the near bottom
86 brines obtained from four hypersaline soda lakes in Kulunda Steppe (Tanatar and Bitter lake
87 systems, Altai region). The lakes brines total salt concentration varied from 20 to 40%, the total
88 soluble carbonate alkalinity - from 2.5 to 5 M and the pH - from 10.2 to 11.0. The top flocculant
89 sediment layer together with the near bottom brines was sucked into 50 ml syringe through silicon
90 tubing and placed into a sterile 50 ml Falcon tube, resulting in an approximate volumetric ratio
91 between the solid and liquid fractions of 1:5. After transportation to the laboratory the samples
92 were separated onto clean brine top layer and concentrated sediment, all of which then were kept at
93 4°C. Before inoculation, the two fractions from each of four lakes were combined in equal
94 proportions to make two master mixes (brine and sediments) used as inocula at 1% final (v/v).

95 For the enrichment and further cultivation of pure cultures two basic mineral media were
96 used, both containing 4 M total Na⁺. The neutral 4 M NaCl base medium included (g l⁻¹): NaCl,
97 240; K₂HPO₄ 2.5 g l⁻¹; NH₄Cl 0.5 g l⁻¹, KCl 5 g l⁻¹, 20 mg l⁻¹ yeast extract and was adjusted to pH 7
98 with 10% KH₂PO₄. The alkaline sodium carbonate base contained (g l⁻¹): Na₂CO₃ 190, NaHCO₃
99 45, NaCl 16, KCl 5, K₂HPO₄ 1 and 20 mg l⁻¹ yeast extract (pH 10). After sterilization, the basic
100 media were supplemented with 1 ml l⁻¹ of trace metal and vitamin solutions [12] and 1 mM
101 MgSO₄. NH₄Cl (4 mM) was also added from 1 M sterile stock solution to the carbonate base after
102 sterilization. To prepare final medium with a certain pH/alkalinity, the two ready to use bases were
103 mixed in different proportions resulting in pH range from 8 to 10. For the enrichments from salt
104 lakes, the neutral base medium was used as it is, while for the soda lake enrichments the neutral
105 and alkali base media were mixed 3:1 with the final pH 9.5. For the pH range from 7 to 8, the NaCl
106 base was supplied with various amounts of 1 M filter-sterilized NaHCO₃, while for the pH below 7

107 it was titrated with sterile 1 M KH_2PO_4 . Carbon and energy substrates were added from sterile
108 10% stock solution.

109 For isolation of pure cultures, several rounds of enrichments were repeated at dilutions
110 1:100 and finally the sediment-free enrichments were plated onto solid medium obtained by
111 mixing the liquid alkaline medium and 4% washed agar at 50°C in ratio 3:2. To compensate for the
112 decreased salinity, solid NaCl was added to the liquid medium before mixing with melted agar to
113 bring the final salt concentration back to 4 M total Na^+ . The isolation of pure cultures was achieved
114 from separate colonies which grew back in liquid medium with the target polysaccharide and the
115 purity was confirmed by 16S rRNA gene and genome sequencing.

116

117 *Pure culture characterization*

118 Cell morphology was examined by using phase contrast microscopy (Zeiss Axioplane Imaging 2,
119 Germany). Substrate utilization profiles were performed in medium containing 1 part of the
120 alkaline base and 7 parts of the NaCl base (final pH 9.0). For the pH profiling, the two media were
121 mixed in various proportions as described above and soluble starch served as the substrate. The
122 growth was measured by increase of OD_{600} with pH monitoring at each point. Anaerobic
123 cultivation was performed as described previously [13]. Catalase and oxidase activity were tested
124 with 3% (v/v) H_2O_2 and 0.1% *N,N,N,N* tetramethyl-p-phenylenediamine hydrochloride,
125 respectively, using cell-free extract (obtained by sonication) from cells of type strains. The
126 protease, esterase/lipase activities were tested on plates spotted with fully grown liquid cultures:
127 using casein/gelatin (hydrolysis zones after flooding with 10% TCA) and emulsified
128 tributyrin/olive oil (turbidity clearance), respectively. Antibiotic sensitivity of type strains [AArc-](#)
129 [St1-1 and AArc-St2](#) was tested in liquid medium at pH 9 with starch as substrate.

130 The intact polar lipids (IPLs) and respiratory quinones were analyzed as described
131 previously [14]. Briefly, the lipid fraction was extracted from freeze-dried cells with sonication in

132 methanol:dichloromethane:phosphate buffer (2:1:0.8, v:v), followed by phase separation by
133 adjusting the solvent mix to a ratio 1:1:0.9. The lipids and quinones were analyzed by normal
134 phase, high performance liquid chromatography-ion trap mass spectrometry (HPLC-ITMS) and
135 identified by masses and mass spectral fragmentation according to literature [14-15].

136

137 *Genome sequencing*

138 Genomic DNA from the type strains AArc-St1-1 and AArc-St2 was extracted using DNeasy
139 PowerLyzer Microbial Kit (Qiagen) according to manufacturer instructions. Quality and quantity
140 of the DNA samples were measured with Trinean Xpose spectrophotometer (PLT Scientific
141 Instruments) and Qubit 2.0 fluorometer (Thermo Fisher Scientific) DNA libraries were prepared
142 using KAPA HyperPlus kit (KAPAbiosystems) according to manufacturer recommendations.
143 Paired-end sequencing (2x100bp) was performed using Illumina NextSeq. Obtained reads were
144 filtered (quality and length) with CLC Genomics Workbench v.10. Genomes of the strains were
145 assembled using SPAdes v.3.15.2 [16] --isolate mode with --trusted-contigs option (contigs
146 obtained from Unicycler v.0.4.9 [17] were used as trusted contigs). Contigs with length \leq 500bp or
147 with low coverage were eliminated. Genome assemblies statistics were checked with Quast v.5.0.2
148 [18-19]. Completeness and contamination levels were detected using CheckM v.1.1.2 [20] with
149 archaea-specific marker set.

150

151 *Phylogenetic and genomic analyses*

152 For 16S rRNA gene sequence-based phylogenetic analysis 16S rRNA gene sequences of the seven
153 isolates were aligned with the sequences of type species of all genera within *Halobacteria* (as well
154 as *Archaeoglobus fulgidus* VC-16, *Methanocella paludicola* SANAE, *Methanothermobacter*
155 *thermautotrophicus* Delta H used as the outgroup). Multiple sequence alignment and phylogenetic
156 tree construction were performed as described earlier [9]. For phylogenomic analysis based on the

157 “ar122” set of conserved single copy archaeal proteins [21] the protein sequences were identified
158 and aligned in *in silico* proteomes of type species of all genera within *Halobacteria* (nontype
159 species were taken for *Halalkalicoccus*, *Halorbellus*, *Natronoarchaeum* and *Halohasta* genera
160 because the genomes of type species are not available) using the GTDB-tk v.1.7.0 with reference
161 data v.202 [22]. The phylogenomic tree was constructed in the RAxML v.8.2.12 [23] with the
162 PROTGAMMAILG model of amino acid substitution; local support values were 1000 rapid
163 bootstrap replications. Phylogenetic trees were visualized using iTOL v.6.5.2 [24]. The whole
164 genome-based comparisons were done as described by Sorokin et al. [25].

165 For functional genome analysis, genes encoding carbohydrate-active enzymes (CAZymes)
166 were searched in genomes of AArc-St2 and AArc-St1-1 strains using dbCAN v.3.0.2 [26]. Further
167 manual checking of the specificity of discovered glycosidases and other CAZymes were performed
168 using BLAST against Swiss-Prot/PDB database.

169

170 **Results and discussion**

171

172 *Enrichment and isolation of pure cultures*

173 The primary enrichments for amylolytic and fructan-utilizing natronoarchaea were performed with
174 amylopectin (insoluble starch) or levan/inulin, respectively, in the presence of 200 mg l⁻¹
175 streptomycin to suppress growth of bacteria. The enrichments from sediment fraction showed
176 visible growth (after removal of the sediment particles by low speed centrifugation) after one week
177 of incubation, while the brine enrichments became turbid and pinkish after 10-14 days of
178 incubation. After several 1:100 transfers, the liquid cultures were plated and individual pink-
179 colored colonies were transferred back to the corresponding liquid media. This procedure was
180 repeated 2 more times to ensure the homogeneity of colony morphology. The purity and identity of

181 obtained cultures were verified by the 16S rRNA gene sequencing. The list of isolates is shown in
182 **Table 1.**

183 Cell morphology of the isolates grown at pH 9 and 4 M total Na⁺ with soluble starch is
184 shown on **Fig. 1.** Cultures of all strains were dominated by nonmotile (with only a few
185 occasionally showing slow motility) flattish cocci with irregular contour, typical for haloarchaea,
186 with a small fraction of flat rods ("boards"). The colonies of all isolates were colored from orange
187 to red and concentrated cell pellets were bright red, what is also typical for aerobic haloarchaea.

188
189 *Chemotaxonomy*

190 Membrane polar lipids and respiratory lipoquinones were analyzed in two strains, one of which
191 represents a group 1 (AArc-St1-1^T), while another a group 2 (AArc-St2^T). In both, the core lipids
192 were dominated by archaeol (AR; C₂₀-C₂₀) with a smaller proportion of extended archaeol (Ext-
193 AR; C₂₀-C₂₅). The polar head groups of the intact polar lipids were phosphatidylglycerophosphate
194 methylether (PGP-Me) and phosphatidylglycerol (PG). Both the core lipids and the polar head
195 groups are very similar to the closest phylogenetic relatives of the amylolytic natronoarchaea (see
196 below in comparative tables). The major difference is the absence of glycolipids and sulfolipids
197 (such as phosphatidylglycerosulfate and sulfated glycosyl diethers) in natronoarchaea, which are
198 more common in neutrophilic haloarchaea. The only respiratory lipoquinone species detected in
199 both strains was the fully saturated MK-8:8, one of the most common in haloarchaea [15].

200
201 *Phylogenetic and genomic analyses*

202 The genome of AArc-St2 was assembled to 20 scaffolds including one circular plasmid, while the
203 genome of AArc-St1-1 was assembled into 32 scaffolds with no plasmids in it (Suppl. **Table S1**).
204 Genome size was 3.26 Mbp (GC content was 51.5%) for strain AArc-St2 and 3.29 Mbp (GC
205 content was 61%) for strain AArc-St1-1. Completeness and contamination levels for AArc-St2

206 genome were 100% / 0% and for AArc-St1-1 – 99.07% / 1.87%. Genomic sequences are available
207 in NCBI GenBank database with accession numbers JAKRVX000000000 (AArc-St2) and
208 JAKRVY000000000 (AArc-St1-1).

209 Seven natronoarchaeal isolates formed two clusters on 16S rRNA gene sequence-based
210 phylogenetic tree. Four strains of the group 1 clustered with the recently described facultatively
211 anaerobic sulfur-reducing natronarchaeon *Natranaeroarchaeum sulfidigenes* which can also use
212 starch as a substrate for growth either aerobically or anaerobically [27-28]. The three closely
213 related isolates from the group 2 formed a novel genus-level lineage with the genera
214 "*Halalkalirubrum*" and *Halohasta* as the nearest neighbors (around 93 and 92 % sequence identity
215 to its type species, respectively). This potentially new genus lineage also includes multiple
216 uncharacterized isolates from various hypersaline habitats whose 16S rRNA gene sequences were
217 deposited recently in the GenBank. However, the 16S rRNA gene sequence-based phylogenetic
218 tree (**Fig. 2a**) had relatively low bootstrap support of its inner nodes within the class *Halobacteria*.
219 Phylogenomic tree based on 122 archaeal conserved single-copy protein markers of the strains
220 AArc-St1-1 and AArc-St2 and other haloarchaea supported the branching, obtained 16S rRNA
221 gene sequence-based tree but showed a better resolution of the inner nodes (**Fig. 2b**). Strain AArc-
222 St2 formed a novel-genus branch in a distinct cluster containing genera "*Halalkalirubrum*",
223 *Halohasta* and *Halonotius*. The latter three genera are currently classified in the order
224 *Haloferacales*, family *Halorubraceae* [29]. Strain AArc-St1-1 was closely related to
225 *Natranaeroarchaeum sulfidigenum* AArc-S and formed a potential new species in the genus which
226 is currently classified within the family *Natranarchaeaceae* [28]. Strains AArc-St2 and AArc-St1-
227 1 were proposed to be type species within their lineages.

228 For additional support of phylogenetic and phylogenomic analyses, ANI and AAI values
229 were calculated for the genomes of AArc-St1-1 and AArc-St2 and the nearest relatives (Suppl.
230 **Table S2 and S3**). The AAI values between strain AArc-St2 and the species of three related

231 haloarchaeal genera with available genome sequences ranged from 62.7% to 70.1%. These values
232 are below the level of AAI for the representatives of the majority of genera, for which AAI results
233 were compared [30], but it is similar to the intergenera level within the whole cluster. The ANI
234 values varied from 70.7% to 73%. Same calculations for strain AArc-St1-1 and
235 *Natranaeroarchaeum sulfidigenum* AArc-S showed AAI and ANI values of 90.1% and 88.8%,
236 respectively, confirming the separate species status of AArc-St1-1 within the genus
237 *Natranaeroarchaeum*. Although, according to the 16S rRNA gene sequence phylogeny, strain
238 AArc-St1-2 might be sufficiently distant from the other three members of the group 1 and the type
239 strain AArc-S^T of *Natranaeroarchaeum sulfidigenes* (98.6-98.8 and 98.5%, respectively), the
240 proposed type species of the genus *Natranaeroarchaeum*, its phenotypic properties were quite
241 similar to the other group 1 strains implying it would be more practical to classify all four isolates
242 of this group in a single species.

243
244 *Metabolic properties*

245 All isolates were capable of utilizing alpha-1,4/1,6 glucans as growth substrates, including soluble
246 starch, amylopectin (insoluble starch), glycogen, dextrin, cyclodextrin and pullulan. Furthermore,
247 the group 1 isolates can also grow with levan (polyfructose with beta-2,6 backbond) while the
248 group 2 strains utilized another fructan - inulin (polyfructose with beta-2,1 backbond). Two other
249 alpha-bonded polysaccharides tested, including dextran from *Leuconostoc* and arabinan were not
250 utilized by any of the seven isolates, as well as various beta-glucans (amorphous cellulose, xylan,
251 xyloglucan, chitin, mannan, glucomannan, galactomannan, lichenan, laminarin and galactan). All
252 strains can also grow on three sugar dimers including maltose, α,α -trehalose and cellobiose. In
253 addition, the group 2 strains were able to grow with glycerol. None of the other tested substrates,
254 except for a weak growth with mannose, gave positive results (glucose, sucrose, galactose,
255 arabinose, rhamnose, raffinose, aminosugars, uronic acids, xylose and arabinose, sugar alcohols,

256 C2-C6 organic acids, pepton). Such limited substrate profile characterize the isolates as narrow-
257 specialized saccharolytics. Anaerobic fermentative growth with either starch, maltose or arginine
258 was not observed. None of the isolates were capable of anaerobic respiratory growth with soluble
259 starch or maltose as substrates, using thiosulfate, DMSO, fumarate or nitrate as the electron
260 acceptors. However, all four group 1 isolates grew anaerobically with maltose as the electron donor
261 and carbon source and elemental sulfur as the electron acceptor, similar to the closely related type
262 species of the genus *Natranaeroarchaeum*. In 15 d incubation (4 M Na⁺, pH 9, 30°C) the following
263 amount of sulfide was produced: 15.6 mM by AArc-St1-1, 5.8 mM by AArc-St1-2; 19 mM by
264 AArc-St1-3 and 10.5 mM by AArc-St-lev1. In comparison, *Natranaeroarchaeum sulfidigenes*
265 formed 30 mM sulfide in 6 days of cultivation at pH 9.5.

266 Two type strains tested positive for catalase reaction and TMPD-oxidase. The protease,
267 esterase and lipase activities were negative in all strains in the spot-plate tests. Ammonium and
268 yeast extract (but not nitrate) can serve as the N-source in cultures grown with soluble starch for
269 both strains. Urea was only utilized by strain AArc-St1-1 consistent with the presence of the
270 *ureABCDEFGF* urease operon in the genome. Indole formation from tryptophan (Kovac's reagent
271 test) showed a weak positive result only for strain AArc-St2. The type strains grown in liquid
272 culture at pH 9 with soluble starch were insensitive to streptomycin, penicillin G, ampicillin,
273 kanamycin, vancomycin and gentamicin up to 200 mg l⁻¹. Rifampicin and chloramphenicol
274 inhibited growth at 50 mg l⁻¹, and tetracyclin - at 100 mg l⁻¹.

275 All isolates grew well at as low Mg concentration as 1 mM, while in their sodium
276 requirement they are typical extreme halophiles, growing optimally at 4 M total Na⁺ and within the
277 range from 3 to 5 M (tested at pH 8.8). The cells of isolates in both groups lyzed at salinity
278 downshift below 2 M total Na⁺. The pH profiling of four cultures at 4 M total Na⁺ showed that they
279 are moderate alkaliphiles with optimal growth within a pH range from 8.0 to 8.8 (**Fig. 3**). The

280 maximum growth temperature of type strains grown with soluble starch at pH 8.5 and 4 M total
281 Na⁺ was 50°C for the group 1 strains and 48°C for the group 2 strains.

282 Comparative properties of the group 1 and group 2 isolates with their nearest phylogenetic
283 relatives are shown in **Tables 2** and **3**.

284 The main difference of the two groups from each other was in utilization of two different
285 fructans and the phylogeny. The key difference of the group 1 isolates from the type species of the
286 genus *Natrananaeroaechaeum* was their inability for anaerobic growth and inability to grow at
287 extremely high pH values above 9.3. The main difference of the group 2 isolates from the nearest
288 related genera is that they are the only ones isolated from soda lakes. The two out of three related
289 genera (*Halohasta* and *Halonotius*) are definitely neutrophiles, while, despite the reported ability
290 of "*Halalkalirubrum salinum*" to grow up to pH 10.5, there is a doubt about it. First, the organism
291 is isolated from a salt lake with pH 8.5, thus being only a slightly alkaline salt (but not soda) lake.
292 Even natronoarchaea isolated from hypersaline soda lakes with permanent pH above 10, seldom
293 grow above pH 10.2. Secondly, the final pH values were apparently not measured during the pH
294 profiling, which makes the reported values for the maximum pH unverified. Hence, the newly
295 isolated amyolytic strains from soda lakes can still be considered as first obligate alkaliphilic
296 (albeit only moderate) representatives of this group of related genera. They also differs from the
297 other three genera in their alpha-glucan/fructan substrate specialization and the absence of glyco-
298 and sulfo-lipids in their membranes.

299

300 *Genomic analysis*

301 The genome search the two type strains (dbCAN) identified a set of genes typically encoding
302 alpha-amylases and alpha-glucosidases (GH13 and 15 families) in both representatives of two
303 groups, although in AArc-St1-1 the total number and the fraction of putative extracellular
304 amylases are much more abundant (Suppl. **Table S4**, consensus results from HHMER/DIAMOND

305 tools). This is also in agreement with the growth and amylase activity results (Suppl. **Fig. 1**). These
306 enzymes would allow utilization of a spectrum of alpha-linked glucans such as amylopectin,
307 soluble starch, dextrin, glycogen, pullulan, maltose and trehalose as sole carbon and energy
308 sources. In this respect strain AArc-St1-1 is highly similar to the type species of the genus
309 *Natranaeroarchaeum* [16-17]. Furthermore genomes of both type strains contain genes coding for
310 beta-fructosidases of the GH families 32 and 68, which is in agreement with their ability to use
311 fructans as growth substrate. As for the other glucanases encoded in two genomes, such as the
312 beta-endo-1,3/1,4-glucanases of GH16 and 81 families (in AArc-St1-1), and pectin lyase (PL
313 family) in AArc-St2, none of the tested potential polysaccharide substrates for these hydrolases
314 supported growth (laminarin, lichenan, xylan, beta-glucan, glucomannan, mannan, curdlan,
315 pachyman or pectin).

316 In respect to the osmoprotection and pH homeostasis, both genomes encode a range of
317 typical potassium import complexes (but variable in copy numbers) and a multisubunit Na^+/H^+
318 antiporter of the Mrp family. Both genomes lacks genes for organic osmolyte import and synthesis,
319 indicating that the organisms rely solely on the potassium accumulation strategy. Both strains
320 produce catalase/peroxidase and have a haem-copper family cytochrome *c* terminal oxidase of the
321 *aa₃* type. In addition, strain AArc-St2 has another terminal oxidase of the *ba₃* type (Supplementary
322 **Table S5**). These also agree with the positive tests for catalase and oxidase in both strains.

323 A major difference between the two type strains was found in the presence of two types of
324 other respiratory complexes. Strain AArc-St2 genome contains a locus apparently coding for the
325 aerobic type of CO-dehydrogenase (Cox, most probably of the type II) lacking in AArc-St1-1.
326 Although the capacity to oxidize CO at low concentration has been demonstrated for several
327 haloarchaeal species [31-32], the physiological role of such potential is still unclear. One of the
328 possibilities is CO detoxification.

329 The genome of AArc-St1-1 contains loci apparently encoding enzymes for sulfur-
330 dependent anaerobic respiration, which are lacking in AArc-St2: two of them are highly
331 homologous to the PsrABCD/SseA and PhsABCD complexes encoded in the genome of type
332 species of genus *Natranaeroarchaeum* responsible for sulfur- and thiosulfate-dependent anaerobic
333 respiration, respectively [27]. In addition, there is a second encoded PsrABCD lacking sulfur
334 transferase more homologous to the one present in sulfur-respiring *Halalkaliarchaeum*
335 *desulfuricum* (Supplementary **Table S5**). In our experience, such genomic potential must enable
336 the anaerobic sulfur respiration in strain AArc-St1-1, similar to the type species
337 *Natranaeroarchaeum sulfidigenes*. And this, indeed, is directly confirmed by the growth
338 experiments, although the activity of sulfur reduction, in general, was lower in the novel isolates
339 than in the type species of the genus *Natranaeroarchaeum*. Furthermore, none of the novel group 1
340 isolates were capable of thiosulfate-dependent anaerobic respiration, which was a prominent trait
341 in the type species. On the other hand, the amylolytic isolates grew much better at fully aerobic
342 conditions, while *N. sulfidigenes* needed transition via microaerophilic conditions before it started
343 to grow at aerobic conditions. This reflects the difference in enrichment conditions used to isolate
344 these closely related but still differentially specialized species of the same genus.

345 Overall, on the basis of distinct phenotypic, phylogenetic and genomic features, the group 1
346 isolates from hypersaline soda lakes are proposed to be classified in a novel species within the
347 genus *Natranaeroarchaeum* as *Natranaeroarchaeum aerophilus* sp. nov. (type strain AArc-St1-1),
348 while the group 2 isolates are forming a new species in a new genus for which the name
349 *Natronocalculus amylovorans* gen. nov., sp. nov. (type strain AArc-St2). The protologues for the
350 new taxa are presented in **Tables 4** and **5**.

351

352 **Funding information**

353 This work was supported by the Russian Science Foundation (20-14-00250) and by the Russian Ministry of
354 Higher Education and Science (genomes sequencing and annotation) for the DYS, AGE, TVK and IVK. NB
355 and JSSD acknowledge support from the ERC Horizon 2020 Program (grant agreement № 694569 –
356 MICROLIPIDS).

357 **Conflict of interests**

358 The authors declare that there is no conflict of interests.

359

360

361 **References**

362

- 363 1. Oren, A. (2011) Diversity of halophiles. In: Horikoshi, K. (Ed.) Extremophiles Handbook. Springer,
364 Tokyo, pp. 309-325.
- 365 2. Andrei, A.Ş., Banciu, H.L., Oren, A. (2012) Living with salt: Metabolic and phylogenetic diversity
366 of archaea inhabiting saline ecosystems. FEMS Microbiol. Lett. 330, 1-9.
- 367 3. Oren, A. (2013) Life at high salt concentrations. In: Rosenberg, E., DeLong, E.F., Lory, S.,
368 Stackebrandt, E., Thompson, F., (Eds.), The Prokaryotes: Prokaryotic Communities and
369 Ecophysiology, Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 421–40.
- 370 4. Oren, A. (2015) Halophilic microbial communities and their environments. Curr Opin Biotechnol
371 33, 119-24.
- 372 5. Grant, W.D., Jones, B.E. (2016) Bacteria, archaea and viruses of soda lakes. In: Schagerl, M., (Ed.),
373 Soda Lakes of East Africa, Springer International Publishing, Cham, pp. 97–147.
- 374 6. Oren, A., Ventosa, A., Grant, W.D. (1997) Proposed minimal standards for description of new taxa
375 in the order *Halobacteriales*. Int J Syst Bacteriol 47, 233–238.
- 376 7. Sorokin, D.Y., Toshchakov, S.V., Kolganova, T.V., Kublanov, I.V. (2015) Halo(natrono)archaea
377 isolated from hypersaline lakes utilize cellulose and chitin as growth substrates. Frontiers Microbiol
378 6, 942.
- 379 8. Sorokin, D.Y., Khijniak, T.V., Kostrikina, N.A., Elcheninov, A.G., Toshchakov, S.V., Bale, N.J.,
380 Sinnighe Damsté, J.S., Kubalnov, I.V. (2018) *Natronobiforma cellulositropha* gen. nov., sp. nov., a
381 novel haloalkaliphilic member of the family *Natrialbaceae* (class *Halobacteria*) from hypersaline
382 alkaline lakes. Syst Appl Microbiol 41, 355–362.
- 383 9. Sorokin, D.Y., Elcheninov, A.G., Toshchakov, S.V., Bale, N.J., Sinnighe Damsté, J.S., Khijniak,
384 T.V., Kublanov, I.V. (2019) *Natrarchaeobius chitinivorans* gen. nov., sp. nov., and
385 *Natrarchaeobius halalkaliphilus* sp. nov., alkaliphilic, chitin-utilizing haloarchaea from hypersaline
386 alkaline lakes. Syst Appl Microbiol 42, 309-318.
- 387 10. Kobayashi, T., Kanai H., Hayashi T., Akiba T., Akaboshi R., Horikoshi K. (1992) Haloalkaliphilic
388 maltotriose-forming α -amylase from the archaebacterium *Natronococcus* sp. strain Ah-36. J
389 Bacteriol 174, 3439-3444.
- 390 11. Kanai, H., Yashi, K., T, Aono, R., Kudo, T. (1995) *Natronococcus amylolyticus* sp. nov., a
391 haloalkaliphilic archaeon. Int J Syst Bacteriol 45, 762-766.
- 392 12. Pfennig, N., Lippert, K.D. (1966) Über das Vitamin B12-Bedürfnis phototropher
393 Schwefelbakterien. Arch Mikrobiol 55, 245-256.

- 394 13. Sorokin, D.Y., Yakimov, M.M., Messina, E., Merkel, A.Y., Koenen, M., Bale, N.J., Sinninghe
395 Damsté, J.S. (2021) *Halapricum desulfuricans* sp. nov., carbohydrate-utilizing sulfur-reducing
396 haloarchaea from hypersaline lakes. Syst Appl Microbiol 44, 126249.
- 397 14. Bale, N.J., Sorokin, D.Y., Hopmans, E.C., Koenen, M., Rijpstra, W.I.C., Villanueva, L., Wienk, H.,
398 Sinninghe Damsté, J.S. (2019) New insights into the polar lipid composition of halophilic
399 euryarchaea from hyper saline lakes. Front Microbiol 10, 377.
- 400 15. Elling, F.J., Becker, K.W., Könneke, M., Schröder J.M., Kellermann, M.Y., Thomm, M., Hinrichs,
401 K.-U. (2016) Respiratory quinones in Archaea: phylogenetic distribution and application as
402 biomarkers in the marine environment. Environ. Microbiol 18, 692–707.
- 403 16. Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
404 Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G.,
405 Alekseyev, M.A., Pevzner, P.A. (2012) SPAdes: A New Genome Assembly Algorithm and Its
406 Applications to Single-Cell Sequencing. J Comput Biol 19, 455–477.
- 407 17. Wick, R.R., Judd, L.M., Gorrie, C.L., Holt, K.E. (2017) Unicycler: Resolving bacterial genome
408 assemblies from short and long sequencing reads. PLoS Comput Biol 13, 1-22.
- 409 18. Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G. (2013) QUASt: Quality assessment tool for
410 genome assemblies. Bioinformatics 29, 1072–1075.
- 411 19. Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., Gurevich, A. (2018) Versatile genome
412 assembly evaluation with QUASt-LG. Bioinformatics 34, i142–50.
- 413 20. Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W. (2015) CheckM:
414 Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes.
415 Genome Res 25, 1043–55.
- 416 21. Rinke C, Chuvochina M, Mussig AJ, P-A Chaumeil, Davín AA, Waite DW, Whitman WB, Parks
417 DH, Hugenholtz P. (2021) A standardized archaeal taxonomy for the Genome Taxonomy Database.
418 Nat Microbiol 6, 946-959.
- 419 22. Parks, D.H., Chuvochina, M., Waite, D.W., Rinke, C., Skarszewski, A., Chaumeil, P.-A.,
420 Hugenholtz, P. (2018) A standardized bacterial taxonomy based on genome phylogeny substantially
421 revises the tree of life. Nat Biotechnol 36(10), 996–1004.
- 422 23. Stamatakis, A. (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of
423 large phylogenies. Bioinformatics 30, 1312–1313.
- 424 24. Letunic, I., Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
425 developments. Nucleic Acids Res 47, W256-W259.
- 426 25. Sorokin, D.Y., Elcheninov, A.G., Khijniak, T. V., Zaharycheva, A.P., Boueva, O. V., Ariskina, E.
427 V., Bunk, B., Spröer, C., Evtushenko, L.I., Kublanov, I. V., Hahnke, R.L. (2022)
428 *Natronosporangium hydrolyticum* gen. nov., sp. nov., a haloalkaliphilic polyhydrolytic
429 actinobacterium from a soda solonchak soil in Central Asia. Syst Appl Microbiol 45, 126307.

- 430 26. Zhang, H., Yohe, T., Huang, L., Entwistle, S., Wu, P., Yang, Z., Busk, P.K., Xu, Y., Yin, Y. (2018)
431 dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res*
432 46(W1), W95–101.
- 433 27. Sorokin, D.Y., Messina, E., Smedile, F., La Cono, V., Hallsworth, J.E. (2021) Carbohydrate-
434 dependent sulfur respiration in halo(alkali)philic euryarchaea from hypersaline lakes. *Environ*
435 *Microbiol* 23, 3779-3808.
- 436 28. Sorokin, D.Y., Yakimov, M.M., Messina, E., Merkel, A.Y., Koenen, M., Bale N.J., Sinninghe
437 Damsté J.S. (2022) *Natronaerarchaeum desulfuricum* gen. nov., sp. nov., carbohydrate-utilizing
438 sulfur-respiring haloarchaeon from hypersaline lakes and proposal of a new family
439 *Natronoarchaeaceae* fam. nov. in the order *Halobacteriales*. *Int J Syst Evol Microbiol* 72,
440 10.1099/ijsem.0.005332.
- 441 29. Gupta, R.S., Naushad, S., Fabros, R., Adeolu, M. (2016) A phylogenomic reappraisal of family-
442 level divisions within the class *Halobacteria*: proposal to divide the order *Halobacteriales* into the
443 families *Halobacteriaceae*, *Haloarculaceae* fam. nov., and *Halococcaceae* fam. nov., and the order
444 *Haloferacales* into the families, *Haloferacaceae* and *Halorubraceae* fam. nov. *Antonie van*
445 *Leeuwenhoek* 109, 565-587.
- 446 30. Luo, C., Rodriguez-R, L.M., Konstantinidis, K.T. (2014) MyTaxa: An advanced taxonomic
447 classifier for genomic and metagenomic sequences. *Nucleic Acids Res* 42, e73.
- 448 31. King, G.M. (2015) Carbon monoxide as a metabolic energy source for extremely halophilic
449 microbes: Implications for microbial activity in Mars regolith. *Proc Nat Ac Sci USA* 112, 4465-
450 4470.
- 451 32. Myers, M.R., King, G.M. (2020) *Halobacterium bonnevillei* sp. nov., *Halobaculum saliterrae* sp.
452 nov. and *Halovenus carboxidivorans* sp. nov., three novel carbon monoxide-oxidizing Halobacteria
453 from saline crusts and soils. *Int J Syst Evol Microbiol* 70, 4261-4268.
- 454 33. Shimane, Y., Hatada, Y., Minegishi, H., Mizuki, T., Echigo, A. (2010) *Natronoarchaeum*
455 *mannanilyticum* gen. nov., sp. nov., an aerobic, extremely halophilic archaeon isolated from
456 commercial salt. *Int J Syst Evol Microbiol* 60, 2529–2534.
- 457 34. Zhenqiang Zuo . Dahe Zhao . Jian Zhou . Jing Han . Hua Xiang (2021) *Halalkalirubrum salinum*
458 gen. nov., sp. nov., a halophilic archaeon isolated from a saline lake. *Ant van Leeuwenhoek* 114,
459 83–94.
- 460 35. Mou, Y.-Z., Qiu, X.-X., Zhao, M.-L., Cui, H.-L., Oh, D., Dyall-Smith, M.L. (2012) *Halohasta*
461 *litorea* gen. nov. sp. nov., and *Halohasta litchfieldiae* sp. nov., isolated from the Daliang
462 aquaculture farm, China and from Deep Lake, Antarctica, respectively. *Extremophiles* 16, 895-901.
- 463 36. Durán-Viseras, A., Andrei A.-S., Ghai, R., Sánchez-Porro, C., Ventosa, A. (2019) New *Halonotius*
464 species provide genomics-based insights into cobalamin synthesis in haloarchaea. *Front Microbiol*
465 10, 1928.
- 466

467

468 **Table 1.** List of pure cultures of natronoarchaea enriched and isolated from
 469 hypersaline soda lakes in southwestern Siberia with amylopectin and fructans

Strain	Source	Enrichment substrate	Phylogenetic group	Closest relative
AArc-St1-1 ^T	Sediments	Amylopectin	Group 1	<i>Natronaeroarchaeum sulfidigenes</i>
AArc-St1-2	Brines			
AArc-St1-3	Brines			
AArc-lev1	Sediments	Levan		
AArc-St2 ^T	Sediments	Amylopectin	Group 2	<i>"Halalkalirubrum halophilum"</i>
AArc-St3	Brines			
AArc-in2	Sediments			

470

471

472

473 **Table 2.** Comparative properties of amylolytic natronoarchaea of group 1 with the type species
 474 of the nearest phylogenetically related genera [28, 33]

Property	<i>"Natranaeroarchaeum aerophilus"</i> (4 strains)	<i>Natranaeroarchaeum sulfidigenes</i> JCM 34033 ^T	<i>Natronoarchaeum mannanilyticum</i> JCM 16328 ^T
Cell morphology	flat pleomorphic, motility not observed	flat pleomorphic, motile	pleomorphic, nonmotile
Pigmentation	red	red (aerobic); pink (anaerobic)	red
PHA accumulation	-	-	-
Aerobic growth	+	+	+
Anaerobic growth by: sugar fermentation	-	-	-
sulfur/thiosulfate respiration	+	+	-
sulfoxide respiration	-	-	-
Number of Psr/Phs operons in genomes	3	2	0*
<i>e</i> -donors for anaerobic growth	glucose, maltose	sugars, starch, glycerol	-
Substrates for aerobic growth	starch-like alpha-glucans, levan, maltose, cellobiose, trehalose	sugars, starch, yeast extract	lactose, raffinose, sucrose, maltose, cellobiose, starch, galactomannan, pyruvate, lactate, glutamate, yeast extract, peptone
Amylase	+	+	+
Esterase/lipase	-	- (tributylin/olive oil)	- (Tween 80)
Protease	-	- (gelatin, casein)	- (gelatin)
Catalase/oxidase	+/+	+/+(w)	-/+(w)
Indole from tryptophane	-	-	+
Salinity range (opt.) (M Na ⁺)	3.0-5.0 (4.0)	2.5-4.5 (3.5)	1.6-4.2 (2.5-3.2)
pH range (opt.)	7.2-9.3 (8.0-8.8)	8.5-10.2 (9.5-9.7)	6.0-9.5 (8.5-9.0)
Temperature max (°C)	50 (at pH 8.5)	45 (at pH 9)	55
Core lipids	C ₂₀ -C ₂₀ , C ₂₀ -C ₂₅ DGE	C ₂₀ -C ₂₀ , C ₂₀ -C ₂₅ DGE	NR
Intact membrane polar lipids: phospholipids	PG, PGP-Me	PG, PGP-Me	PG, PGP-Me, PGP, S ₂ -DGDE
glycolipids/sulfolipids	-	-	NR
Respiratory lipoquinones	MK-8:8	MK-8:8	NR
DNA G+C	61.0 (genome)	60.8% (genome)	63.0 (mol%)
Type of hypersaline habitat	Hypersaline soda lakes		Marine solar saltern

475 NR, not reported; (v) - variable property in different species of the same genus; w (weak); Psr/Phs - polysulfide/thiosulfate
 476 reductase; *genome of *N. phillipinensis*. Lipids: (PG) phosphatidylglycerol, phosphatidylglycero-phosphate (PGP), (PGP-Me)
 477 phosphatidylglycerophosphate methyl ester, disulfated diglycosyl diether (S₂-DGDE), (DGE) - dialkyl glycerol ether.
 478

479 **Table 3.** Comparative properties of group 2 isolates with nearest phylogenetic relatives [34-36].

Property	" <i>Natranocalculus amylovorans</i> " (3 isolates)	" <i>Halalkalirubrum halophilum</i> "	<i>Halohasta</i> (2 species)	<i>Halonotius</i> (4 species)
Cell morphology	flat pleomorphic nonmotile	pleomorphic, nonmotile	rods, motile	polymorphic rods, motility (V)
Pigmentation	red	red	red	red
PHA accumulation	-	NR	NR	
Anaerobic growth by:				
sugar fermentation	-	NR	NR	NR
sulfur respiration	-	NR	NR	NR
DMSO respiration	-	-	-	+(1 species)
Growth substrates				
carbohydrates:	starch-like alpha-glucans, inulin, maltose, cellobiose, trehalose, glycerol	glucose, maltose, fructose, sorbose, lactose, xylose, mannitol, sorbitol	glucose, sucrose; mannose, galactose, lactose, maltose (all V)	glucose, arabinose, fructose, galactose, sucrose, maltose, raffinose, xylose, mannitol, sorbitol, glycerol, (all V)
organic acids:	none	acetate, pyruvate, lactate, fumarate, succinate, citrate	pyruvate, lactate, succinate, malate, fumarate, citrate (V);	pyruvate, citrate, tartrate (all V)
Amylase	+ (soluble starch)	- (soluble starch)	- (soluble starch)	- (soluble starch)
Esterase/lipase	- (tributylin/olive oil)	- (Tweens)	- (Tween 80)	- (Tween 80)
Protease	- (gelatin, casein)	- (gelatin, casein)	- (gelatin, casein)	- (gelatin, casein)
Catalase/oxidase	+/+	+/+	+/V	V/V
Indole from tryptophane	+(w)	+	-	NR
Salinity range (opt.) M Na ⁺	3-5 (4.0)	1.9-4.2 (2.5)	2.0-4.7 (2.5-3.0)	2.5-6.0** (3.0-4.0)
Mg ²⁺ demand	low	low	high	high
pH range (opt.)	7.2-9.3 (8.5-8.8)	7.0-10.5* (8.5-9.5)	5.5-9.0* (7.0-7.5)	5.0-9.0 (7.0-7.5)
Temperature max (°C)	48 (at pH 8.5)	42	45-50	45-50
Core lipids	C ₂₀ -C ₂₀ , C ₂₀ -C ₂₅ DGE	NR	NR	NR
Intact membrane polar lipids:				
phospholipids:	PG, PGP-Me	PG, PGP-Me	PG, PGP-Me, PA	PG, PGP-Me
glycolipids:	-	1, unidentified	S-DGD-1	S-DGD-1
sulfolipids:	-	PGS		PGS (V)
Respiratory lipoquinones	MK-8:8	NR	NR	NR
DNA G+C (% genomic)	51.5 (type strain)	58.4 (type strain)	58.8 (type species)	59.7-62.7 (4 species)
Type of hypersaline habitat	soda lakes	salt lake	solar saltern	solar saltern, saline soils

480 NR, not reported; (V), variable property in different species of the same genus; * actual final pH values were not measured;
 481 ** reported for the type species, but not verified in any further research; PA, phosphatidic acid; PGS, phosphatidylglycerol
 482 sulfate; S-DGD-1, sulfated mannosyl glucosyl diether; other abbreviations (see **Table 2**).

483
 484
 485
 486
 487
 488
 489
 490
 491

Table 4. *Natronaeroarchaeum aerophilus*: protologue

Parameter	Species: <i>Natronaeroarchaeum aerophilus</i> sp. nov.
Author	Dimitry Y. Sorokin
Species name	<i>aerophilus</i>
Genus name	<i>Natronaeroarchaeum</i>
Specific epithet	<i>aerophilus</i>
Species status (SPST)	sp. nov.
Etymology	a.e.ro'phi.lus Gr. masc. n. <i>aer</i> , air; N.L. masc. adj. <i>philus</i> (from Gr. masc. adj. <i>philos</i>), friend, loving; N.L. masc. adj. <i>aerophilus</i> , air-loving.
Description of the new taxon	The cells are angular, flat, polymorphic coccoids or rods, mostly nonmotile, variable in size from 1 to 3 µm. The cells lyse in hypotonic solutions below 1 M NaCl, produces red carotenoids. The core membrane diether lipids are composed of C ₂₀ -C ₂₀ DGE (archaeol) and C ₂₀ -C ₂₅ DGE (extended archaeol). The polar lipid head groups include phosphatidylglycerolphosphate methyl ester (PGP-Me) and phosphatidylglycerol (PG). The dominant respiratory quinone is MK-8:8. Saccharolytic with limited substrate profile including several starch-like alpha-glucans, levan, maltose, trehalose and cellobiose. Capable of anaerobic sulfogenic growth with glucose and maltose as electron donors and sulfur as electron acceptor. Ammonium, urea and yeast extract serve as the N-source. Oxidase and catalase positive. Maximum growth temperature is 50°C. Extremely halophilic with a range of total Na ⁺ for growth from 3 to 5 M (optimum at 4 M) and moderately alkaliphilic, with a pH range for growth from 7.2 to 9.3 (optimum at 8.0-8.8). The G + C content of the DNA is 61.0% (genome of the type strain). Habitat - aerobic sediments and brines of hypersaline soda lakes. The type strain (AArc-St1-1 ^T =JCM 32519 ^T =UQM 41458 ^T) was isolated aerobic sediments of hypersaline soda lakes in Kulunda Steppe (Altai, Russia). The species also includes other three closely related strains isolated from the same area. The draft genome of type strain is deposited in the GenBank under accession number JAKRVY000000000.
Authors	Dimitry Y. Sorokin, Alexander G. Elcheninov, Tatjana V. Khizhniak, Michel Koenen, Nicole J. Bale, Jaap S. Sinninghe Damsté, Ilya V. Kublanov
Title	<i>Natronocalculus amylovorans</i> gen. nov., sp. nov., and <i>Natronaeroarchaeum aerophilus</i> sp. nov., dominant culturable amylolytic natronoarchaea from hypersaline soda lakes in southwestern Siberia.
Journal	Systematic and Applied Microbiology
Corresponding author	Dimitry Y. Sorokin
E-mail of corresponding author	soroc@inmi.ru; d.sorokin@tudelft
Designation of the type strain	AArc-St1-1
Strain collection numbers	JCM 32519 ^T =UQM 41458 ^T
16S rRNA gene accession numbers	MG584707- MG584709; ON003450
Genome accession numbers	JAKRVY000000000 (type strain)
Genome status	Draft
G+C, %	61.0 (genome of type strain)
Country of origin	Russian Federation
Region of origin	Altai region
Date of isolation	2016
Source of isolation	Surface aerobic sediments from hypersaline soda lakes
Sampling dates	2015-07-07
Geographic location	S-W Siberia, Kulunda Steppe; southern Russia
Latitude	51°39' N; 49°10' N; 48°14' N
Longitude	79°48' E; 46°39' E; 46°35' E
Depth	0-2 cm
Temperature of the sample	20°C
pH of the sample	10-11
Salinity of the sample	18-36%
Number of strains in study	4
Source of isolation of non-type strains	Surface aerobic sediments and brines from hypersaline soda lakes, S-W Siberia, Kulunda Steppe; southern Russia
Growth medium, incubation conditions	4 M total Na ⁺ , pH 9; incubation - 37°C; starch as substrates; aerobic
Conditions of preservation	Deep freezing in 15% glycerol (v/v)
Gram stain	Negative
Cell shape	Pleomorphic flat coccoids
Cell size	0.8-2 µm in diameter
Motility (MOTY)	Mostly nonmotile
Sporulation	None
Colony morphology	Flat, compact, max. 2 mm, red
Temperature range for growth	nd
Lowest temperature for growth	nd
Highest temperature for growth	50
Optimal temperature for growth	37-40
Lowest pH for growth	7.2
Highest pH for growth	9.3
Optimum pH for growth	8-8.8
pH category	Moderately alkaliphilic
Lowest Na ⁺ concentration for growth	3.0 M
Highest Na ⁺ concentration for growth	5.0 M
Optimum salt concentration for growth	4.0 M total Na ⁺
Other salts important for growth	KCl; Na-carbonates
Salinity category	Extremely halophilic
Relation to oxygen	Facultatively anaerobic
O ₂ conditions for strain testing	Fully aerobic
Carbon source used (class)	Carbohydrates

Specific compounds	Starch-like alpha-glucans, levan
Nitrogen source	Ammonium, urea, yeast extract
Terminal electron acceptor	O ₂ and S ₈
Energy metabolism	Chemoorganotrophic
Phospholipids	Core membrane lipids are C ₂₀ -C ₂₀ DGE (archaeol) and C ₂₀ -C ₂₅ DGE (extended archaeol). Polar head groups are phosphatidylglycerophosphate methylester (PGP-Me) and phosphatidylglycerol (PG)
Respiratory lipoquinones	MK-8:8
Glycolipids (GLYC)	-
Habitat (HABT)	Hypersaline soda lakes
Extraordinary features (EXTR)	Narrowly specialized amylolytics

493

494

495 **Table 5.** *Natronocalculus amylovorans*: protologue

Parameter	Genus: <i>Natronocalculus</i> gen. nov.	Species: <i>Natronocalculus amylovorans</i> sp. nov.
Author (AUTE)	Dimitry Y. Sorokin	
Species name (SPNA)		<i>amylovorans</i>
Genus name (GENA)	<i>Natronocalculus</i>	
Specific epithet (SPEP)	-	<i>amylovorans</i>
Species status (SPST)	-	sp. nov.
Etymology (GETY/SPTY)	Na.tro.no.cal'cu.lus N.L. neut. n. <i>natron</i> , arbitrarily derived from the Arabic n. natrun or natron soda; L. masc. n. <i>calculus</i> , pebble, gravel; N.L. masc. n. <i>Natronocalculus</i> , soda loving pebble-shaped cells	a.my.lo.vo'rans. Gr. neut. n. <i>amylon</i> , starch; L. inf. v. <i>vorare</i> , to devour; N.L. part. adj. <i>amylovorans</i> , <u>eating</u> starch
Type species of the genus (TYPE)	<i>Natronocalculus amylovorans</i>	yes
Description of new taxon	Obligately aerobic and organotrophic member of <i>Halobacteriales</i> narrowly specialized on utilization of starch-like polymers. Core lipids are dominated by archaeol and extended archaeol with PGP-Me and PG polar groups. Glycolipids are not present. MK-8:8 is the dominant lipoquinone. Extremely halophilic and moderately alkaliphilic inhabitants of hypersaline soda lakes. The three-letter abbreviation is Ncl.	The cells are angular, flat, polymorphic coccoids or rods, nonmotile, from 0.8 to 2.0 µm. The cells lyse in hypotonic solutions below 1 M NaCl. Colonies are orange-red. The core membrane diether lipids include of C ₂₀ -C ₂₀ DGE (archaeol) and C ₂₀ -C ₂₅ DGE (extended archaeol). The polar lipid head groups consists of _osphatidylglycerolphosphate methyl ester (PGP-Me) and phosphatidylglycerol (PG). The dominant respiratory menaquinone is MK-8:8. Obligately aerobic saccharolytic with limited substrate profile including several starch-like alpha-glucans, inulin, maltose, trehalose, cellobiose and glycerol. Ammonium and yeast extract serve as the N-source. Oxidase and catalase positive. Maximum growth temperature is 48°C. Extremely halophilic with a range of total Na ⁺ for growth from 3 to 5 M (optimum at 4 M) and moderately alkaliphilic, with a pH range for growth from 7.2 to 9.3 (optimum at 8.5-8.8). The G + C content of the DNA is 51.5% (genome of the type strain). Habitat - aerobic sediments and brines of hypersaline soda lakes. The type strain (AArc-St2 ^T =JCM 32475 ^T =UQM 41459 ^T) was isolated <u>from</u> aerobic sediments of hypersaline soda lakes in Kulunda Steppe (Altai, Russia). The species also includes 2 other closely related strains isolated from the same area. The draft genome of type strain is deposited in the GenBank under accession number JAKRVX000000000.
Authors (AUT)	Dimitry Y. Sorokin, Alexander G. Elcheninov, Tatjana V. Khizhniak, Michel Koenen, Nicole J. Bale, Jaap S. Sinninghe Damsté, Ilya V. Kublanov	
Title (TITL)	<i>Natronocalculus amylovorans</i> gen. nov., sp. nov., and <i>Natraneroarchaeum aerophilus</i> sp. nov., dominant culturable amyolytic natronoarchaea from hypersaline soda lakes in southwestern Siberia.	
Journal (JOUR)	Systematic and Applied Microbiology	
Corresponding author (COAU)	Dimitry Y. Sorokin	
E-mail of corresponding author (EMAU)	d.sorokin@tudelft; soroc@inmi.ru	
Strain collection numbers (COLN)	-	JCM 32475; UQM 41459
16S rRNA gene accession number (16 SR)	-	MG584710; ON000203; ON000205
Genome accession numbers		JAKRVX000000000
Genome status (GSTA)		Draft
GC mol % (GGCM)	-	51.5 (genome type strain)
Country of origin (COUN)	Russian Federation	Russian Federation
Region of origin (REGI)	-	Altai region
Date of isolation (DATI)	-	2016
Source of isolation (SOUR)	Hypersaline soda lakes	Surface sediments and brines of hypersaline soda lakes in southwestern Siberia
Sampling dates (DATS)	2015	2015
Geographic location (GEOL)	S-W Siberia	S-W Siberia
Latitude (LATI)	-	51°39' N; 49°10' N; 48°14' N
Longitude (LONG)	-	79°48' E; 46°39' E; 46°35' E
Depth (DEPT)		0-2 cm
Temperature of the sample (TEMS)		20°C
pH of the sample (PHSA)		10-11
Salinity of the sample (SALS)		18-36%
Number of strains in study (NSTR)	3	3
Source of isolation of non-type strains (SAMP)	-	Surface sediments and brines of hypersaline soda lakes in southwestern Siberia
Growth medium, incubation conditions (CULT)		4 M total Na ⁺ , pH 9; incubation - 37°C; starch as substrates; aerobic
Conditions of preservation (PRES)	Deep freezing in 15% glycerol (v/v)	
Gram stain (GRAM)	negative	
Cell shape (CSHA)	Pleomorphic, from flat irregular coccoids	
Cell size (CSZI)	-	0.8-2 µm in diameter

Motility (MOTY)	-	nonmotile
Sporulation (SPOR)	none	
Colony morphology (COLM)		Pink-orange, up to 2 mm, flat
Temperature range for growth (TEMR)		
Lowest temperature for growth (TEML)		
Highest temperature for growth (TEMH)		48 (at pH 8.5)
Optimal temperature for growth (TEMO)		37-40°C
Lowest pH for growth (PHLO)		7.2
Highest pH for growth (PHHI)		9.3
Optimum pH for growth (PHOP)		8.5-8.8
pH category (PHCA)	alkaliphile (optimum > 8.5)	
Lowest NaCl concentration for growth (SALL)	3.0 M total Na ⁺	
Highest NaCl concentration for growth (SALH)	5 M total Na ⁺	
Optimum salt concentration for growth (SALO)	4.0 M total Na ⁺	
Other salts important for growth	Sodium carbonates	
Salinity category (SALC)	extremely halophilic	
Relation to oxygen (OREL)	aerobe	
O ₂ conditions for strain testing (OCON)	aerobic	
Carbon source used (class) (CSUC)	carbohydrates	
Specific compounds (CSUC)	Starch-like alpha glucans and inulin	
Nitrogen source (NSOU)	Ammonium, yeast extract	
Terminal electron acceptor (ELAC)	O ₂	
Energy metabolism (EMET)	chemoorganotrophic	
Phospholipids (PHOS)	Core membrane lipids are archaeol (C ₂₀ -C ₂₀ DGE) and extended archaeol (C ₂₀ -C ₂₅ DGE) Polar lipids are phosphatidylglycerophosphate methyl ester (PGP-Me) and phosphatidylglycerol (PG)	
Glycolipids (GLYC)	-	
Respiratory <u>lipoquinones</u>	MK8:8	
Habitat (HABT)	Hypersaline soda lakes	

496

497

498

499 **Legends to the figures**

500

501 **Fig. 1.** Cell morphology (phase contrast microphotographs) of starch-utilizing natronoarchaea growing
 502 aerobically at 4 M total Na⁺, pH 9 and 37°C. **(a-d)**, group 1, including strains AArc-St1-1^T, AArc-St1-
 503 2, AArc-St1-3 and AArc-lev11, respectively. **(e-f)**, group 2, including strains AArc-St2^T and AArc-
 504 in2.

505 **Figure 2.** Phylogeny of amylolytic natronoarchaea.

506 **(a)** Figure X. (a) 16S rRNA gene sequence-based maximum-likelihood phylogenetic tree, showing the position
 507 of AArc-St2^T and AArc-St1-1^T (in bold) within the *Halobacteria* class. The black circles at nodes indicate that
 508 the percentage of corresponding support values was above 50. *Archaeoglobus fulgidus* VC-16^T, *Methanocella*
 509 *paludicola* SANAET, *Methanothermobacter thermautotrophicus* Delta H^T were used as an outgroup (not shown).
 510 Species in clusters: I (*Halodesulfurarchaeum*, *Halanaeroarchaeum*, *Halarchaeum*, *Halobacterium*,
 511 *Salarchaeum*, *Halocalculus*); II (*Halomicroarcula*, *Haloarcula*, *Halorientalis*, *Halorhabdus*, *Halococcoides*,
 512 *Halapricum*, *Salinirussus*, *Halovenus*); III (*Haloglomerus*, *Natronomonas*, *Halosegnis*, *Salinirubellus*,
 513 *Halomarina*, *Halocatena*); IV (*Halopelagius*, *Haloferax*, *Halogeometricum*, *Haloquadratum*, *Halobellus*); V
 514 (*Halococcus*, *Halalkalicoccus*, *Haloarchaeobius*, *Halorubellus*); VI (*Halorussus*, *Halomicrococcus*,
 515 *Haladaptatus*); VII (*Saliphagus*, *Natribaculum*, *Halovarius*, *Natronococcus*, *Halovivax*, *Natronobiforma*,
 516 *Halostagnicola*, *Natronobacterium*, *Halopiger*, *Halobiforma*, *Natrarchaeobaculum*, *Natronolimnohabitans*,
 517 *Natronolimnobius*, *Natronorubrum*, *Natrinema*, *Haloterrigena*, *Natrialba*, *Natrarchaeobius*, *Salinadaptatus*);
 518 VIII (*Halosimplex*, *Salinibaculum*, *Halosiccatus*, *Halomicrobium*); IX (*Salinigranum*, *Haloplanus*, *Halobium*,
 519 *Halegenticoccus*, *Halogranum*, *Haloprofundus*, *Halolamina*); X (*Halalkaliarchaeum*, *Halopenitus*, *Halorubrum*,
 520 *Haloparvum*).

521 **(b)** Maximum likelihood phylogenetic tree based on concatenated alignment of 122 conserved archaeal proteins
 522 and showing position of strains AArc-St2^T and AArc-St1-1^T (in bold) within the class *Halobacteria*. The branch
 523 lengths correspond to the number of substitutions per site with corrections associated with the models. The black
 524 circles at nodes indicate that the percentage of corresponding support values was above 50. *Archaeoglobus*
 525 *fulgidus* VC-16^T, *Methanocella paludicola* SANAET and *Methanothermobacter thermautotrophicus* Delta H^T
 526 were used as an outgroup (not shown). Species in clusters: I (*Halanaeroarchaeum*, *Halodesulfurarchaeum*,
 527 *Halarchaeum*, *Halobacterium*, *Salarchaeum*, *Halocalculus*); II (*Halalkalicoccus*, *Halorussus*, *Halomicrococcus*,
 528 *Haladaptatus*); III (*Haloarchaeobius*, *Halorubellus*, *Halovivax*, *Saliphagus*, *Natronobiforma*, *Halostagnicola*,
 529 *Natrarchaeobius*, *Natrarchaeobaculum*, *Salinadaptatus*, *Halopiger*, *Natronolimnobius*, *Natronobacterium*,
 530 *Halobiforma*, *Natrialba*, *Natronococcus*, *Natrinema*, *Haloterrigena*, *Natronorubrum*, *Natronolimnohabitans*);
 531 IV (*Halococcus*, *Halocatena*, *Halomarina*, *Natronomonas*, *Haloglomerus*, *Halosegnis*, *Halorientalis*, *Halapricum*,
 532 *Salinirussus*, *Salinibaculum*, *Halovenus*, *Halosimplex*, *Halococcoides*, *Halorhabdus*, *Halomicrobium*,
 533 *Halosiccatus*, *Halomicroarcula*, *Haloarcula*); V (*Haloplanus*, *Salinirubrum*, *Haloprofundus*, *Halegenticoccus*,
 534 *Halogranum*, *Salinigranum*, *Haloferax*, *Halopelagius*, *Halogeometricum*, *Halobellus*, *Haloquadratum*).

535

536

537 **Fig. 3.** pH profiles for growth with soluble starch in amylolytic natronoarchaea at 4 M total Na⁺ and
 538 37°C. Actual final pH are shown. The results are average from two parallel incubations.

2 ***Natronocalculus amylovorans* gen. nov., sp. nov., and *Natranaeroarchaeum***
3 ***aerophilus* sp. nov., dominant culturable amyolytic natronoarchaea from**
4 **hypersaline soda lakes in southwestern Siberia.**

5
6
7 Dimitry Y. Sorokin^{a,b*}, Alexander G. Elcheninov^a, Tatjana V. Khizhniak^a, Michel Koenen^c, Nicole
8 J. Bale^c, Jaap S. Sinninghe Damsté^c and Ilya V. Kublanov^a

9
10 ^a*Winogradsky Institute of Microbiology, Research Centre of Biotechnology, Russian Academy of Sciences,*
11 *Moscow, Russia*

12 ^b*Department of Biotechnology, Delft University of Technology, The Netherlands*

13 ^c*NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Microbiology and*
14 *Biogeochemistry, Texel, The Netherlands*

15
16 *Author for correspondence:

17 D.Y. Sorokin; e-mail: soroc@inmi.ru; d.sorokin@tudelft.nl

18
19
20
21
22
23
24
25
26
27 Running title: *Natronocalculus amylovorans* gen. nov., sp. nov. and *Natranaeroarchaeum*
28 *aerophilus* sp. nov.
29

30 The draft genome sequences of strains AArc-St1-1^T and AArc-St2^T are deposited in the GenBank
31 under the numbers JAKRVY000000000 and JAKRVX000000000, respectively
32

33 **Abstract**

34 Several pure cultures of alkaliphilic haloarchaea were enriched and isolated from hypersaline
35 soda lakes in southwestern Siberia using amylopectin and fructans as substrates. Phylogenomic
36 analysis placed the isolates into two distinct groups within the class *Halobacteria*. Four isolates
37 forming group 1 were closely related to a recently described *Natronaeroarchaeum sulfidigenes* and
38 the other three strains forming group 2 represent a novel genus-level phylogenetic lineage. All
39 isolates are saccharolytic archaea growing with various starch-like alpha-glucans including soluble
40 starch, amylopectin, dextrin, glycogen, pullulane and cyclodextrin. In addition, group 1 can also
41 use levan while group 2 - inulin (plant storage beta-fructans). Group 1 strains can also grow
42 anaerobically with either glucose or maltose using elemental sulfur as the electron acceptor. Both
43 groups are moderately alkaliphilic with a pH range for growth from 7.2 to 9.3 (optimum between
44 8.0-8.8) and low Mg-demanding extreme halophiles growing optimally at 4 M total Na⁺. The major
45 respiratory menaquinone is MK-8:8 and the core biphytanyl lipids are dominated by archaeol (C₂₀-
46 C₂₀) and a less abundant extended archaeol (C₂₀-C₂₅) with PG and PGP-Me as polar groups. The
47 four isolates of group 1 are suggested to be classified into a new species as *Natronaeroarchaeum*
48 *aerophilus* sp. nov. (type strain AArc-St1-1^T=JCM 32519^T=UQM 41458^T). The three isolates of
49 group 2 are proposed to form a new genus and species for which the name *Natronocalculus*
50 *amylovorans* gen. nov., sp. nov. is suggested (type strain AArc-St2^T=JCM 32475^T=UQM 41459^T).

51
52 Key words: hypersaline soda lakes, natronoarchaea, amylolytic, starch, fructans,

53 *Natronaeroarchaeum*, *Halobacteria*

54

55

56

57 **Introduction**

58 Most of the known species of aerobic extremely halophilic and haloalkaliphilic euryarchaea
59 (natronoarchaea) were enriched and isolated from hypersaline alkaline lakes on unspecific media
60 containing rich soluble organic substrates, such as peptone, yeast extract or simple sugars [1-5].
61 Yet, these extremophilic, organoheterotrophic archaea are definitely more important in organic
62 matter mineralization, in particular organic polymers, in hypersaline systems than is widely
63 recognized. For example, a test for amylolytic activity with soluble starch is included into the
64 minimal standards for taxonomy characterization of haloarchaea [6] but starch is rarely used for
65 targeted isolation of amylolytic haloarchaea which would not only hydrolyze the polymer but also
66 able to utilize it as growth substrate. And this situation is also true for other polysaccharides. So
67 far, only a few examples of haloarchaea specialized on utilization of recalcitrant polysaccharides
68 have been described in pure culture, which is particularly true for natronoarchaea living in alkaline
69 hypersaline (soda) lakes. Our recent targeted enrichments from such lakes using insoluble forms of
70 cellulose and chitin resulted in isolation of several groups of natronoarchaea highly specialized on
71 utilization of either various forms of cellulose and xylan (genera *Natronobiforma* and
72 *Natronolimnobius*) or chitin (genus *Natrarchaeobius*) [7-9]. Following further in this direction, we
73 used insoluble starch (amylopectin) or beta-fructans as substrates to enrich for amylolytic or
74 fructanolytic natronoarchaea from hypersaline soda lakes. So far, we are aware only of a single
75 amylolytic natronarchaeon, *Natronococcus amylolyticus*, specifically isolated from the hypersaline
76 soda lake Magadi in Kenya using starch as a growth substrate [10-11], while no natronoarchaea,
77 growing on fructans are currently known.

78
79 Here we describe phenotypic, phylogenetic and genomic properties of two novel taxa of amylolytic
80 natronoarchaea enriched from hypersaline soda lakes in southwestern Siberia, which specialized on
81 utilizing various alpha-glucan and beta-fructan polysaccharides as growth substrates.

82 **Materials and Methods**

83

84 *Enrichment and cultivation conditions*

85 The sources of inocula were mixed surface (0-3 cm deep) aerated sediments and the near bottom
86 brines obtained from four hypersaline soda lakes in Kulunda Steppe (Tanatar and Bitter lake
87 systems, Altai region). The lakes brines total salt concentration varied from 20 to 40%, the total
88 soluble carbonate alkalinity - from 2.5 to 5 M and the pH - from 10.2 to 11.0. The top flocculant
89 sediment layer together with the near bottom brines was sucked into 50 ml syringe through silicon
90 tubing and placed into a sterile 50 ml Falcon tube, resulting in an approximate volumetric ratio
91 between the solid and liquid fractions of 1:5. After transportation to the laboratory the samples
92 were separated onto clean brine top layer and concentrated sediment, all of which then were kept at
93 4°C. Before inoculation, the two fractions from each of four lakes were combined in equal
94 proportions to make two master mixes (brine and sediments) used as inocula at 1% final (v/v).

95 For the enrichment and further cultivation of pure cultures two basic mineral media were
96 used, both containing 4 M total Na⁺. The neutral 4 M NaCl base medium included (g l⁻¹): NaCl,
97 240; K₂HPO₄ 2.5 g l⁻¹; NH₄Cl 0.5 g l⁻¹, KCl 5 g l⁻¹, 20 mg l⁻¹ yeast extract and was adjusted to pH 7
98 with 10% KH₂PO₄. The alkaline sodium carbonate base contained (g l⁻¹): Na₂CO₃ 190, NaHCO₃
99 45, NaCl 16, KCl 5, K₂HPO₄ 1 and 20 mg l⁻¹ yeast extract (pH 10). After sterilization, the basic
100 media were supplemented with 1 ml l⁻¹ of trace metal and vitamin solutions [12] and 1 mM
101 MgSO₄. NH₄Cl (4 mM) was also added from 1 M sterile stock solution to the carbonate base after
102 sterilization. To prepare final medium with a certain pH/alkalinity, the two ready to use bases were
103 mixed in different proportions resulting in pH range from 8 to 10. For the enrichments from salt
104 lakes, the neutral base medium was used as it is, while for the soda lake enrichments the neutral
105 and alkali base media were mixed 3:1 with the final pH 9.5. For the pH range from 7 to 8, the NaCl
106 base was supplied with various amounts of 1 M filter-sterilized NaHCO₃, while for the pH below 7

107 it was titrated with sterile 1 M KH_2PO_4 . Carbon and energy substrates were added from sterile
108 10% stock solution.

109 For isolation of pure cultures, several rounds of enrichments were repeated at dilutions
110 1:100 and finally the sediment-free enrichments were plated onto solid medium obtained by
111 mixing the liquid alkaline medium and 4% washed agar at 50°C in ratio 3:2. To compensate for the
112 decreased salinity, solid NaCl was added to the liquid medium before mixing with melted agar to
113 bring the final salt concentration back to 4 M total Na^+ . The isolation of pure cultures was achieved
114 from separate colonies which grew back in liquid medium with the target polysaccharide and the
115 purity was confirmed by 16S rRNA gene and genome sequencing.

116

117 *Pure culture characterization*

118 Cell morphology was examined by using phase contrast microscopy (Zeiss Axioplane Imaging 2,
119 Germany). Substrate utilization profiles were performed in medium containing 1 part of the
120 alkaline base and 7 parts of the NaCl base (final pH 9.0). For the pH profiling, the two media were
121 mixed in various proportions as described above and soluble starch served as the substrate. The
122 growth was measured by increase of OD_{600} with pH monitoring at each point. Anaerobic
123 cultivation was performed as described previously [13]. Catalase and oxidase activity were tested
124 with 3% (v/v) H_2O_2 and 0.1% *N,N,N,N* tetramethyl-p-phenylenediamine hydrochloride,
125 respectively, using cell-free extract (obtained by sonication) from cells of type strains. The
126 protease, esterase/lipase activities were tested on plates spotted with fully grown liquid cultures:
127 using casein/gelatin (hydrolysis zones after flooding with 10% TCA) and emulsified
128 tributyrin/olive oil (turbidity clearance), respectively. Antibiotic sensitivity of type strains AArc-
129 St1-1 and AArc-St2 was tested in liquid medium at pH 9 with starch as substrate.

130 The intact polar lipids (IPLs) and respiratory quinones were analyzed as described
131 previously [14]. Briefly, the lipid fraction was extracted from freeze-dried cells with sonication in

132 methanol:dichloromethane:phosphate buffer (2:1:0.8, v:v), followed by phase separation by
133 adjusting the solvent mix to a ratio 1:1:0.9. The lipids and quinones were analyzed by normal
134 phase, high performance liquid chromatography-ion trap mass spectrometry (HPLC-ITMS) and
135 identified by masses and mass spectral fragmentation according to literature [14-15].

136

137 *Genome sequencing*

138 Genomic DNA from the type strains AArc-St1-1 and AArc-St2 was extracted using DNeasy
139 PowerLyzer Microbial Kit (Qiagen) according to manufacturer instructions. Quality and quantity
140 of the DNA samples were measured with Trinean Xpose spectrophotometer (PLT Scientific
141 Instruments) and Qubit 2.0 fluorometer (Thermo Fisher Scientific) DNA libraries were prepared
142 using KAPA HyperPlus kit (KAPAbiosystems) according to manufacturer recommendations.
143 Paired-end sequencing (2x100bp) was performed using Illumina NextSeq. Obtained reads were
144 filtered (quality and length) with CLC Genomics Workbench v.10. Genomes of the strains were
145 assembled using SPAdes v.3.15.2 [16] --isolate mode with --trusted-contigs option (contigs
146 obtained from Unicycler v.0.4.9 [17] were used as trusted contigs). Contigs with length \leq 500bp or
147 with low coverage were eliminated. Genome assemblies statistics were checked with Quast v.5.0.2
148 [18-19]. Completeness and contamination levels were detected using CheckM v.1.1.2 [20] with
149 archaea-specific marker set.

150

151 *Phylogenetic and genomic analyses*

152 For 16S rRNA gene sequence-based phylogenetic analysis 16S rRNA gene sequences of the seven
153 isolates were aligned with the sequences of type species of all genera within *Halobacteria* (as well
154 as *Archaeoglobus fulgidus* VC-16, *Methanocella paludicola* SANAE, *Methanothermobacter*
155 *thermautotrophicus* Delta H used as the outgroup). Multiple sequence alignment and phylogenetic
156 tree construction were performed as described earlier [9]. For phylogenomic analysis based on the

157 “ar122” set of conserved single copy archaeal proteins [21] the protein sequences were identified
158 and aligned in *in silico* proteomes of type species of all genera within *Halobacteria* (nontype
159 species were taken for *Halalkalicoccus*, *Halorbellus*, *Natronoarchaeum* and *Halohasta* genera
160 because the genomes of type species are not available) using the GTDB-tk v.1.7.0 with reference
161 data v.202 [22]. The phylogenomic tree was constructed in the RAxML v.8.2.12 [23] with the
162 PROTGAMMAILG model of amino acid substitution; local support values were 1000 rapid
163 bootstrap replications. Phylogenetic trees were visualized using iTOL v.6.5.2 [24]. The whole
164 genome-based comparisons were done as described by Sorokin et al. [25].

165 For functional genome analysis, genes encoding carbohydrate-active enzymes (CAZymes)
166 were searched in genomes of AArc-St2 and AArc-St1-1 strains using dbCAN v.3.0.2 [26]. Further
167 manual checking of the specificity of discovered glycosidases and other CAZymes were performed
168 using BLAST against Swiss-Prot/PDB database.

169

170 **Results and discussion**

171

172 *Enrichment and isolation of pure cultures*

173 The primary enrichments for amylolytic and fructan-utilizing natronoarchaea were performed with
174 amylopectin (insoluble starch) or levan/inulin, respectively, in the presence of 200 mg l⁻¹
175 streptomycin to suppress growth of bacteria. The enrichments from sediment fraction showed
176 visible growth (after removal of the sediment particles by low speed centrifugation) after one week
177 of incubation, while the brine enrichments became turbid and pinkish after 10-14 days of
178 incubation. After several 1:100 transfers, the liquid cultures were plated and individual pink-
179 colored colonies were transferred back to the corresponding liquid media. This procedure was
180 repeated 2 more times to ensure the homogeneity of colony morphology. The purity and identity of

181 obtained cultures were verified by the 16S rRNA gene sequencing. The list of isolates is shown in
182 **Table 1.**

183 Cell morphology of the isolates grown at pH 9 and 4 M total Na⁺ with soluble starch is
184 shown on **Fig. 1.** Cultures of all strains were dominated by nonmotile (with only a few
185 occasionally showing slow motility) flattish cocci with irregular contour, typical for haloarchaea,
186 with a small fraction of flat rods ("boards"). The colonies of all isolates were colored from orange
187 to red and concentrated cell pellets were bright red, what is also typical for aerobic haloarchaea.

188

189 *Chemotaxonomy*

190 Membrane polar lipids and respiratory lipoquinones were analyzed in two strains, one of which
191 represents a group 1 (AArc-St1-1^T), while another a group 2 (AArc-St2^T). In both, the core lipids
192 were dominated by archaeol (AR; C₂₀-C₂₀) with a smaller proportion of extended archaeol (Ext-
193 AR; C₂₀-C₂₅). The polar head groups of the intact polar lipids were phosphatidylglycerophosphate
194 methylether (PGP-Me) and phosphatidylglycerol (PG). Both the core lipids and the polar head
195 groups are very similar to the closest phylogenetic relatives of the amylolytic natronoarchaea (see
196 below in comparative tables). The major difference is the absence of glycolipids and sulfolipids
197 (such as phosphatidylglycerosulfate and sulfated glycosyl diethers) in natronoarchaea, which are
198 more common in neutrophilic haloarchaea. The only respiratory lipoquinone species detected in
199 both strains was the fully saturated MK-8:8, one of the most common in haloarchaea [15].

200

201 *Phylogenetic and genomic analyses*

202 The genome of AArc-St2 was assembled to 20 scaffolds including one circular plasmid, while the
203 genome of AArc-St1-1 was assembled into 32 scaffolds with no plasmids in it (Suppl. **Table S1**).
204 Genome size was 3.26 Mbp (GC content was 51.5%) for strain AArc-St2 and 3.29 Mbp (GC
205 content was 61%) for strain AArc-St1-1. Completeness and contamination levels for AArc-St2

206 genome were 100% / 0% and for AArc-St1-1 – 99.07% / 1.87%. Genomic sequences are available
207 in NCBI GenBank database with accession numbers JAKRVX000000000 (AArc-St2) and
208 JAKRVY000000000 (AArc-St1-1).

209 Seven natronoarchaeal isolates formed two clusters on 16S rRNA gene sequence-based
210 phylogenetic tree. Four strains of the group 1 clustered with the recently described facultatively
211 anaerobic sulfur-reducing natronarchaeon *Natranaeroarchaeum sulfidigenes* which can also use
212 starch as a substrate for growth either aerobically or anaerobically [27-28]. The three closely
213 related isolates from the group 2 formed a novel genus-level lineage with the genera
214 "*Halalkalirubrum*" and *Halohasta* as the nearest neighbors (around 93 and 92 % sequence identity
215 to its type species, respectively). This potentially new genus lineage also includes multiple
216 uncharacterized isolates from various hypersaline habitats whose 16S rRNA gene sequences were
217 deposited recently in the GenBank. However, the 16S rRNA gene sequence-based phylogenetic
218 tree (**Fig. 2a**) had relatively low bootstrap support of its inner nodes within the class *Halobacteria*.
219 Phylogenomic tree based on 122 archaeal conserved single-copy protein markers of the strains
220 AArc-St1-1 and AArc-St2 and other haloarchaea supported the branching, obtained 16S rRNA
221 gene sequence-based tree but showed a better resolution of the inner nodes (**Fig. 2b**). Strain AArc-
222 St2 formed a novel-genus branch in a distinct cluster containing genera "*Halalkalirubrum*",
223 *Halohasta* and *Halonotius*. The latter three genera are currently classified in the order
224 *Haloferacales*, family *Halorubraceae* [29]. Strain AArc-St1-1 was closely related to
225 *Natranaeroarchaeum sulfidigenum* AArc-S and formed a potential new species in the genus which
226 is currently classified within the family *Natranarchaeaceae* [28]. Strains AArc-St2 and AArc-St1-
227 1 were proposed to be type species within their lineages.

228 For additional support of phylogenetic and phylogenomic analyses, ANI and AAI values
229 were calculated for the genomes of AArc-St1-1 and AArc-St2 and the nearest relatives (Suppl.
230 **Table S2 and S3**). The AAI values between strain AArc-St2 and the species of three related

231 haloarchaeal genera with available genome sequences ranged from 62.7% to 70.1%. These values
232 are below the level of AAI for the representatives of the majority of genera, for which AAI results
233 were compared [30], but it is similar to the intergenera level within the whole cluster. The ANI
234 values varied from 70.7% to 73%. Same calculations for strain AArc-St1-1 and
235 *Natranaeroarchaeum sulfidigenum* AArc-S showed AAI and ANI values of 90.1% and 88.8%,
236 respectively, confirming the separate species status of AArc-St1-1 within the genus
237 *Natranaeroarchaeum*. Although, according to the 16S rRNA gene sequence phylogeny, strain
238 AArc-St1-2 might be sufficiently distant from the other three members of the group 1 and the type
239 strain AArc-S^T of *Natranaeroarchaeum sulfidigenes* (98.6-98.8 and 98.5%, respectively), the
240 proposed type species of the genus *Natranaeroarchaeum*, its phenotypic properties were quite
241 similar to the other group 1 strains implying it would be more practical to classify all four isolates
242 of this group in a single species.

243
244 *Metabolic properties*

245 All isolates were capable of utilizing alpha-1,4/1,6 glucans as growth substrates, including soluble
246 starch, amylopectin (insoluble starch), glycogen, dextrin, cyclodextrin and pullulan. Furthermore,
247 the group 1 isolates can also grow with levan (polyfructose with beta-2,6 backbond) while the
248 group 2 strains utilized another fructan - inulin (polyfructose with beta-2,1 backbond). Two other
249 alpha-bonded polysaccharides tested, including dextran from *Leuconostoc* and arabinan were not
250 utilized by any of the seven isolates, as well as various beta-glucans (amorphous cellulose, xylan,
251 xyloglucan, chitin, mannan, glucomannan, galactomannan, lichenan, laminarin and galactan). All
252 strains can also grow on three sugar dimers including maltose, α,α -trehalose and cellobiose. In
253 addition, the group 2 strains were able to grow with glycerol. None of the other tested substrates,
254 except for a weak growth with mannose, gave positive results (glucose, sucrose, galactose,
255 arabinose, rhamnose, raffinose, aminosugars, uronic acids, xylose and arabinose, sugar alcohols,

256 C2-C6 organic acids, pepton). Such limited substrate profile characterize the isolates as narrow-
257 specialized saccharolytics. Anaerobic fermentative growth with either starch, maltose or arginine
258 was not observed. None of the isolates were capable of anaerobic respiratory growth with soluble
259 starch or maltose as substrates, using thiosulfate, DMSO, fumarate or nitrate as the electron
260 acceptors. However, all four group 1 isolates grew anaerobically with maltose as the electron donor
261 and carbon source and elemental sulfur as the electron acceptor, similar to the closely related type
262 species of the genus *Natranaeroarchaeum*. In 15 d incubation (4 M Na⁺, pH 9, 30°C) the following
263 amount of sulfide was produced: 15.6 mM by AArc-St1-1, 5.8 mM by AArc-St1-2; 19 mM by
264 AArc-St1-3 and 10.5 mM by AArc-St-lev1. In comparison, *Natranaeroarchaeum sulfidigenes*
265 formed 30 mM sulfide in 6 days of cultivation at pH 9.5.

266 Two type strains tested positive for catalase reaction and TMPD-oxidase. The protease,
267 esterase and lipase activities were negative in all strains in the spot-plate tests. Ammonium and
268 yeast extract (but not nitrate) can serve as the N-source in cultures grown with soluble starch for
269 both strains. Urea was only utilized by strain AArc-St1-1 consistent with the presence of the
270 *ureABCDEFG* urease operon in the genome. Indole formation from tryptophan (Kovac's reagent
271 test) showed a weak positive result only for strain AArc-St2. The type strains grown in liquid
272 culture at pH 9 with soluble starch were insensitive to streptomycin, penicillin G, ampicillin,
273 kanamycin, vancomycin and gentamicin up to 200 mg l⁻¹. Rifampicin and chloramphenicol
274 inhibited growth at 50 mg l⁻¹, and tetracyclin - at 100 mg l⁻¹.

275 All isolates grew well at as low Mg concentration as 1 mM, while in their sodium
276 requirement they are typical extreme halophiles, growing optimally at 4 M total Na⁺ and within the
277 range from 3 to 5 M (tested at pH 8.8). The cells of isolates in both groups lysed at salinity
278 downshift below 2 M total Na⁺. The pH profiling of four cultures at 4 M total Na⁺ showed that they
279 are moderate alkaliphiles with optimal growth within a pH range from 8.0 to 8.8 (**Fig. 3**). The

280 maximum growth temperature of type strains grown with soluble starch at pH 8.5 and 4 M total
281 Na⁺ was 50°C for the group 1 strains and 48°C for the group 2 strains.

282 Comparative properties of the group 1 and group 2 isolates with their nearest phylogenetic
283 relatives are shown in **Tables 2** and **3**.

284 The main difference of the two groups from each other was in utilization of two different
285 fructans and the phylogeny. The key difference of the group 1 isolates from the type species of the
286 genus *Natrananaeroaechaeum* was their inability for anaerobic growth and inability to grow at
287 extremely high pH values above 9.3. The main difference of the group 2 isolates from the nearest
288 related genera is that they are the only ones isolated from soda lakes. The two out of three related
289 genera (*Halohasta* and *Halonotius*) are definitely neutrophiles, while, despite the reported ability
290 of "*Halalkalirubrum salinum*" to grow up to pH 10.5, there is a doubt about it. First, the organism
291 is isolated from a salt lake with pH 8.5, thus being only a slightly alkaline salt (but not soda) lake.
292 Even natronoarchaea isolated from hypersaline soda lakes with permanent pH above 10, seldom
293 grow above pH 10.2. Secondly, the final pH values were apparently not measured during the pH
294 profiling, which makes the reported values for the maximum pH unverified. Hence, the newly
295 isolated amyolytic strains from soda lakes can still be considered as first obligate alkaliphilic
296 (albeit only moderate) representatives of this group of related genera. They also differs from the
297 other three genera in their alpha-glucan/fructan substrate specialization and the absence of glyco-
298 and sulfo-lipids in their membranes.

299

300 *Genomic analysis*

301 The genome search the two type strains (dbCAN) identified a set of genes typically encoding
302 alpha-amylases and alpha-glucosidases (GH13 and 15 families) in both representatives of two
303 groups, although in AArc-St1-1 the total number and the fraction of putative extracellular
304 amylases are much more abundant (Suppl. **Table S4**, consensus results from HHMER/DIAMOND

305 tools). This is also in agreement with the growth and amylase activity results (Suppl. **Fig. 1**). These
306 enzymes would allow utilization of a spectrum of alpha-linked glucans such as amylopectin,
307 soluble starch, dextrin, glycogen, pullulan, maltose and trehalose as sole carbon and energy
308 sources. In this respect strain AArc-St1-1 is highly similar to the type species of the genus
309 *Natranaeroarchaeum* [16-17]. Furthermore genomes of both type strains contain genes coding for
310 beta-fructosidases of the GH families 32 and 68, which is in agreement with their ability to use
311 fructans as growth substrate. As for the other glucanases encoded in two genomes, such as the
312 beta-endo-1,3/1,4-glucanases of GH16 and 81 families (in AArc-St1-1), and pectin lyase (PL
313 family) in AArc-St2, none of the tested potential polysaccharide substrates for these hydrolases
314 supported growth (laminarin, lichenan, xylan, beta-glucan, glucomannan, mannan, curdlan,
315 pachyman or pectin).

316 In respect to the osmoprotection and pH homeostasis, both genomes encode a range of
317 typical potassium import complexes (but variable in copy numbers) and a multisubunit Na^+/H^+
318 antiporter of the Mrp family. Both genomes lacks genes for organic osmolyte import and synthesis,
319 indicating that the organisms rely solely on the potassium accumulation strategy. Both strains
320 produce catalase/peroxidase and have a haem-copper family cytochrome *c* terminal oxidase of the
321 *aa₃* type. In addition, strain AArc-St2 has another terminal oxidase of the *ba₃* type (Supplementary
322 **Table S5**). These also agree with the positive tests for catalase and oxidase in both strains.

323 A major difference between the two type strains was found in the presence of two types of
324 other respiratory complexes. Strain AArc-St2 genome contains a locus apparently coding for the
325 aerobic type of CO-dehydrogenase (Cox, most probably of the type II) lacking in AArc-St1-1.
326 Although the capacity to oxidize CO at low concentration has been demonstrated for several
327 haloarchaeal species [31-32], the physiological role of such potential is still unclear. One of the
328 possibilities is CO detoxification.

329 The genome of AArc-St1-1 contains loci apparently encoding enzymes for sulfur-
330 dependent anaerobic respiration, which are lacking in AArc-St2: two of them are highly
331 homologous to the PsrABCD/SseA and PhsABCD complexes encoded in the genome of type
332 species of genus *Natranaeroarchaeum* responsible for sulfur- and thiosulfate-dependent anaerobic
333 respiration, respectively [27]. In addition, there is a second encoded PsrABCD lacking sulfur
334 transferase more homologous to the one present in sulfur-respiring *Halalkaliarchaeum*
335 *desulfuricum* (Supplementary **Table S5**). In our experience, such genomic potential must enable
336 the anaerobic sulfur respiration in strain AArc-St1-1, similar to the type species
337 *Natranaeroarchaeum sulfidigenes*. And this, indeed, is directly confirmed by the growth
338 experiments, although the activity of sulfur reduction, in general, was lower in the novel isolates
339 than in the type species of the genus *Natranaeroarchaeum*. Furthermore, none of the novel group 1
340 isolates were capable of thiosulfate-dependent anaerobic respiration, which was a prominent trait
341 in the type species. On the other hand, the amylolytic isolates grew much better at fully aerobic
342 conditions, while *N. sulfidigenes* needed transition via microaerophilic conditions before it started
343 to grow at aerobic conditions. This reflects the difference in enrichment conditions used to isolate
344 these closely related but still differentially specialized species of the same genus.

345 Overall, on the basis of distinct phenotypic, phylogenetic and genomic features, the group 1
346 isolates from hypersaline soda lakes are proposed to be classified in a novel species within the
347 genus *Natranaeroarchaeum* as *Natranaeroarchaeum aerophilus* sp. nov. (type strain AArc-St1-1),
348 while the group 2 isolates are forming a new species in a new genus for which the name
349 *Natronocalculus amylovorans* gen. nov., sp. nov. (type strain AArc-St2). The protologues for the
350 new taxa are presented in **Tables 4** and **5**.

351

352 **Funding information**

353 This work was supported by the Russian Science Foundation (20-14-00250) and by the Russian Ministry of
354 Higher Education and Science (genomes sequencing and annotation) for the DYS, AGE, TVK and IVK. NB
355 and JSSD acknowledge support from the ERC Horizon 2020 Program (grant agreement № 694569 –
356 MICROLIPIDS).

357 **Conflict of interests**

358 The authors declare that there is no conflict of interests.

359

360

361 **References**

362

- 363 1. Oren, A. (2011) Diversity of halophiles. In: Horikoshi, K. (Ed.) Extremophiles Handbook. Springer,
364 Tokyo, pp. 309-325.
- 365 2. Andrei, A.Ş., Banciu, H.L., Oren, A. (2012) Living with salt: Metabolic and phylogenetic diversity
366 of archaea inhabiting saline ecosystems. FEMS Microbiol. Lett. 330, 1-9.
- 367 3. Oren, A. (2013) Life at high salt concentrations. In: Rosenberg, E., DeLong, E.F., Lory, S.,
368 Stackebrandt, E., Thompson, F., (Eds.), The Prokaryotes: Prokaryotic Communities and
369 Ecophysiology, Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 421–40.
- 370 4. Oren, A. (2015) Halophilic microbial communities and their environments. Curr Opin Biotechnol
371 33, 119-24.
- 372 5. Grant, W.D., Jones, B.E. (2016) Bacteria, archaea and viruses of soda lakes. In: Schagerl, M., (Ed.),
373 Soda Lakes of East Africa, Springer International Publishing, Cham, pp. 97–147.
- 374 6. Oren, A., Ventosa, A., Grant, W.D. (1997) Proposed minimal standards for description of new taxa
375 in the order *Halobacteriales*. Int J Syst Bacteriol 47, 233–238.
- 376 7. Sorokin, D.Y., Toshchakov, S.V., Kolganova, T.V., Kublanov, I.V. (2015) Halo(natrono)archaea
377 isolated from hypersaline lakes utilize cellulose and chitin as growth substrates. Frontiers Microbiol
378 6, 942.
- 379 8. Sorokin, D.Y., Khijniak, T.V., Kostrikina, N.A., Elcheninov, A.G., Toshchakov, S.V., Bale, N.J.,
380 Sinnighe Damsté, J.S., Kubalnov, I.V. (2018) *Natronobiforma cellulositropha* gen. nov., sp. nov., a
381 novel haloalkaliphilic member of the family *Natrialbaceae* (class *Halobacteria*) from hypersaline
382 alkaline lakes. Syst Appl Microbiol 41, 355–362.
- 383 9. Sorokin, D.Y., Elcheninov, A.G., Toshchakov, S.V., Bale, N.J., Sinnighe Damsté, J.S., Khijniak,
384 T.V., Kublanov, I.V. (2019) *Natrarchaeobius chitinivorans* gen. nov., sp. nov., and
385 *Natrarchaeobius halalkaliphilus* sp. nov., alkaliphilic, chitin-utilizing haloarchaea from hypersaline
386 alkaline lakes. Syst Appl Microbiol 42, 309-318.
- 387 10. Kobayashi, T., Kanai H., Hayashi T., Akiba T., Akaboshi R., Horikoshi K. (1992) Haloalkaliphilic
388 maltotriose-forming α -amylase from the archaeobacterium *Natronococcus* sp. strain Ah-36. J
389 Bacteriol 174, 3439-3444.
- 390 11. Kanai, H., Yashi, K., T, Aono, R., Kudo, T. (1995) *Natronococcus amylolyticus* sp. nov., a
391 haloalkaliphilic archaeon. Int J Syst Bacteriol 45, 762-766.
- 392 12. Pfennig, N., Lippert, K.D. (1966) Über das Vitamin B12-Bedürfnis phototropher
393 Schwefelbakterien. Arch Mikrobiol 55, 245-256.

- 394 13. Sorokin, D.Y., Yakimov, M.M., Messina, E., Merkel, A.Y., Koenen, M., Bale, N.J., Sinninghe
395 Damsté, J.S. (2021) *Halapricum desulfuricans* sp. nov., carbohydrate-utilizing sulfur-reducing
396 haloarchaea from hypersaline lakes. Syst Appl Microbiol 44, 126249.
- 397 14. Bale, N.J., Sorokin, D.Y., Hopmans, E.C., Koenen, M., Rijpstra, W.I.C., Villanueva, L., Wienk, H.,
398 Sinninghe Damsté, J.S. (2019) New insights into the polar lipid composition of halophilic
399 euryarchaea from hyper saline lakes. Front Microbiol 10, 377.
- 400 15. Elling, F.J., Becker, K.W., Könneke, M., Schröder J.M., Kellermann, M.Y., Thomm, M., Hinrichs,
401 K.-U. (2016) Respiratory quinones in Archaea: phylogenetic distribution and application as
402 biomarkers in the marine environment. Environ. Microbiol 18, 692–707.
- 403 16. Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
404 Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G.,
405 Alekseyev, M.A., Pevzner, P.A. (2012) SPAdes: A New Genome Assembly Algorithm and Its
406 Applications to Single-Cell Sequencing. J Comput Biol 19, 455–477.
- 407 17. Wick, R.R., Judd, L.M., Gorrie, C.L., Holt, K.E. (2017) Unicycler: Resolving bacterial genome
408 assemblies from short and long sequencing reads. PLoS Comput Biol 13, 1-22.
- 409 18. Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G. (2013) QUASt: Quality assessment tool for
410 genome assemblies. Bioinformatics 29, 1072–1075.
- 411 19. Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., Gurevich, A. (2018) Versatile genome
412 assembly evaluation with QUASt-LG. Bioinformatics 34, i142–50.
- 413 20. Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W. (2015) CheckM:
414 Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes.
415 Genome Res 25, 1043–55.
- 416 21. Rinke C, Chuvochina M, Mussig AJ, P-A Chaumeil, Davín AA, Waite DW, Whitman WB, Parks
417 DH, Hugenholtz P. (2021) A standardized archaeal taxonomy for the Genome Taxonomy Database.
418 Nat Microbiol 6, 946-959.
- 419 22. Parks, D.H., Chuvochina, M., Waite, D.W., Rinke, C., Skarszewski, A., Chaumeil, P.-A.,
420 Hugenholtz, P. (2018) A standardized bacterial taxonomy based on genome phylogeny substantially
421 revises the tree of life. Nat Biotechnol 36(10), 996–1004.
- 422 23. Stamatakis, A. (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of
423 large phylogenies. Bioinformatics 30, 1312–1313.
- 424 24. Letunic, I., Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
425 developments. Nucleic Acids Res 47, W256-W259.
- 426 25. Sorokin, D.Y., Elcheninov, A.G., Khijniak, T. V., Zaharycheva, A.P., Boueva, O. V., Ariskina, E.
427 V., Bunk, B., Spröer, C., Evtushenko, L.I., Kublanov, I. V., Hahnke, R.L. (2022)
428 *Natronosporangium hydrolyticum* gen. nov., sp. nov., a haloalkaliphilic polyhydrolytic
429 actinobacterium from a soda solonchak soil in Central Asia. Syst Appl Microbiol 45, 126307.

- 430 26. Zhang, H., Yohe, T., Huang, L., Entwistle, S., Wu, P., Yang, Z., Busk, P.K., Xu, Y., Yin, Y. (2018)
431 dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res*
432 46(W1), W95–101.
- 433 27. Sorokin, D.Y., Messina, E., Smedile, F., La Cono, V., Hallsworth, J.E. (2021) Carbohydrate-
434 dependent sulfur respiration in halo(alkali)philic euryarchaea from hypersaline lakes. *Environ*
435 *Microbiol* 23, 3779-3808.
- 436 28. Sorokin, D.Y., Yakimov, M.M., Messina, E., Merkel, A.Y., Koenen, M., Bale N.J., Sinninghe
437 Damsté J.S. (2022) *Natronaerarchaeum desulfuricum* gen. nov., sp. nov., carbohydrate-utilizing
438 sulfur-respiring haloarchaeon from hypersaline lakes and proposal of a new family
439 *Natronoarchaeaceae* fam. nov. in the order *Halobacteriales*. *Int J Syst Evol Microbiol* 72,
440 10.1099/ijsem.0.005332.
- 441 29. Gupta, R.S., Naushad, S., Fabros, R., Adeolu, M. (2016) A phylogenomic reappraisal of family-
442 level divisions within the class *Halobacteria*: proposal to divide the order *Halobacteriales* into the
443 families *Halobacteriaceae*, *Haloarculaceae* fam. nov., and *Halococcaceae* fam. nov., and the order
444 *Haloferacales* into the families, *Haloferacaceae* and *Halorubraceae* fam. nov. *Antonie van*
445 *Leeuwenhoek* 109, 565-587.
- 446 30. Luo, C., Rodriguez-R, L.M., Konstantinidis, K.T. (2014) MyTaxa: An advanced taxonomic
447 classifier for genomic and metagenomic sequences. *Nucleic Acids Res* 42, e73.
- 448 31. King, G.M. (2015) Carbon monoxide as a metabolic energy source for extremely halophilic
449 microbes: Implications for microbial activity in Mars regolith. *Proc Nat Ac Sci USA* 112, 4465-
450 4470.
- 451 32. Myers, M.R., King, G.M. (2020) *Halobacterium bonnevillei* sp. nov., *Halobaculum saliterrae* sp.
452 nov. and *Halovenus carboxidivorans* sp. nov., three novel carbon monoxide-oxidizing Halobacteria
453 from saline crusts and soils. *Int J Syst Evol Microbiol* 70, 4261-4268.
- 454 33. Shimane, Y., Hatada, Y., Minegishi, H., Mizuki, T., Echigo, A. (2010) *Natronoarchaeum*
455 *mannanilyticum* gen. nov., sp. nov., an aerobic, extremely halophilic archaeon isolated from
456 commercial salt. *Int J Syst Evol Microbiol* 60, 2529–2534.
- 457 34. Zhenqiang Zuo . Dahe Zhao . Jian Zhou . Jing Han . Hua Xiang (2021) *Halalkalirubrum salinum*
458 gen. nov., sp. nov., a halophilic archaeon isolated from a saline lake. *Ant van Leeuwenhoek* 114,
459 83–94.
- 460 35. Mou, Y.-Z., Qiu, X.-X., Zhao, M.-L., Cui, H.-L., Oh, D., Dyall-Smith, M.L. (2012) *Halohasta*
461 *litorea* gen. nov. sp. nov., and *Halohasta litchfieldiae* sp. nov., isolated from the Daliang
462 aquaculture farm, China and from Deep Lake, Antarctica, respectively. *Extremophiles* 16, 895-901.
- 463 36. Durán-Viseras, A., Andrei A.-S., Ghai, R., Sánchez-Porro, C., Ventosa, A. (2019) New *Halonotius*
464 species provide genomics-based insights into cobalamin synthesis in haloarchaea. *Front Microbiol*
465 10, 1928.
- 466

467

468 **Table 1.** List of pure cultures of natronoarchaea enriched and isolated from
 469 hypersaline soda lakes in southwestern Siberia with amylopectin and fructans

Strain	Source	Enrichment substrate	Phylogenetic group	Closest relative
AArc-St1-1 ^T	Sediments	Amylopectin	Group 1	<i>Natronaeroarchaeum sulfidigenes</i>
AArc-St1-2	Brines			
AArc-St1-3	Brines			
AArc-lev1	Sediments	Levan		
AArc-St2 ^T	Sediments	Amylopectin	Group 2	<i>"Halalkalirubrum halophilum"</i>
AArc-St3	Brines			
AArc-in2	Sediments			

470

471

472

473 **Table 2.** Comparative properties of amylolytic natronoarchaea of group 1 with the type species
 474 of the nearest phylogenetically related genera [28, 33]

Property	<i>"Natranaeroarchaeum aerophilus"</i> (4 strains)	<i>Natranaeroarchaeum sulfidigenes</i> JCM 34033 ^T	<i>Natronoarchaeum mannanilyticum</i> JCM 16328 ^T
Cell morphology	flat pleomorphic, motility not observed	flat pleomorphic, motile	pleomorphic, nonmotile
Pigmentation	red	red (aerobic); pink (anaerobic)	red
PHA accumulation	-	-	-
Aerobic growth	+	+	+
Anaerobic growth by: sugar fermentation	-	-	-
sulfur/thiosulfate respiration	+ (with sulfur)	+ (with sulfur and S ₂ O ₃ ²⁻)	-
sulfoxide respiration	-	-	-
Number of Psr/Phs operons in genomes	3	2	0*
<i>e</i> -donors for anaerobic growth	glucose, maltose	sugars, starch, glycerol	-
Substrates for aerobic growth	starch-like alpha-glucans, levan, maltose, cellobiose, trehalose	sugars, starch, yeast extract	lactose, raffinose, sucrose, maltose, cellobiose, starch, galactomannan, pyruvate, lactate, glutamate, yeast extract, peptone
Amylase	+	+	+
Esterase/lipase	-	- (tributylin/olive oil)	- (Tween 80)
Protease	-	- (gelatin, casein)	- (gelatin)
Catalase/oxidase	+/+	+/+(w)	-/+(w)
Indole from tryptophane	-	-	+
Salinity range (opt.) (M Na ⁺)	3.0-5.0 (4.0)	2.5-4.5 (3.5)	1.6-4.2 (2.5-3.2)
pH range (opt.)	7.2-9.3 (8.0-8.8)	8.5-10.2 (9.5-9.7)	6.0-9.5 (8.5-9.0)
Temperature max (°C)	50 (at pH 8.5)	45 (at pH 9)	55
Core lipids	C ₂₀ -C ₂₀ , C ₂₀ -C ₂₅ DGE	C ₂₀ -C ₂₀ , C ₂₀ -C ₂₅ DGE	NR
Intact membrane polar lipids: phospholipids	PG, PGP-Me	PG, PGP-Me	PG, PGP-Me, PGP, S ₂ -DGDE
glycolipids/sulfolipids	-	-	
Respiratory lipoquinones	MK-8:8	MK-8:8	NR
DNA G+C	61.0 (genome)	60.8% (genome)	63.0 (mol%)
Type of hypersaline habitat	Hypersaline soda lakes		Marine solar saltern

475 NR, not reported; (v) - variable property in different species of the same genus; w (weak); Psr/Phs - polysulfide/thiosulfate
 476 reductase; *genome of *N. phillipinensis*. Lipids: (PG) phosphatidylglycerol, phosphatidylglycero-phosphate (PGP), (PGP-Me)
 477 phosphatidylglycerophosphate methyl ester, disulfated diglycosyl diether (S₂-DGDE), (DGE) - dialkyl glycerol ether.
 478

479 **Table 3.** Comparative properties of group 2 isolates with nearest phylogenetic relatives [34-36].

Property	" <i>Natranocalculus amylovorans</i> " (3 isolates)	" <i>Halalkalirubrum halophilum</i> "	<i>Halohasta</i> (2 species)	<i>Halonotius</i> (4 species)
Cell morphology	flat pleomorphic nonmotile	pleomorphic, nonmotile	rods, motile	polymorphic rods, motility (V)
Pigmentation	red	red	red	red
PHA accumulation	-	NR	NR	
Anaerobic growth by:				
sugar fermentation	-	NR	NR	NR
sulfur respiration	-	NR	NR	NR
DMSO respiration	-	-	-	+(1 species)
Growth substrates				
carbohydrates:	starch-like alpha-glucans, inulin, maltose, cellobiose, trehalose, glycerol	glucose, maltose, fructose, sorbose, lactose, xylose, mannitol, sorbitol	glucose, sucrose; mannose, galactose, lactose, maltose (all V)	glucose, arabinose, fructose, galactose, sucrose, maltose, raffinose, xylose, mannitol, sorbitol, glycerol, (all V)
organic acids:	none	acetate, pyruvate, lactate, fumarate, succinate, citrate	pyruvate, lactate, succinate, malate, fumarate, citrate (V);	pyruvate, citrate, tartrate (all V)
Amylase	+ (soluble starch)	- (soluble starch)	- (soluble starch)	- (soluble starch)
Esterase/lipase	- (tributylin/olive oil)	- (Tweens)	- (Tween 80)	- (Tween 80)
Protease	- (gelatin, casein)	- (gelatin, casein)	- (gelatin, casein)	- (gelatin, casein)
Catalase/oxidase	+/+	+/+	+/V	V/V
Indole from tryptophane	+(w)	+	-	NR
Salinity range (opt.) M Na ⁺	3-5 (4.0)	1.9-4.2 (2.5)	2.0-4.7 (2.5-3.0)	2.5-6.0** (3.0-4.0)
Mg ²⁺ demand	low	low	high	high
pH range (opt.)	7.2-9.3 (8.5-8.8)	7.0-10.5* (8.5-9.5)	5.5-9.0* (7.0-7.5)	5.0-9.0 (7.0-7.5)
Temperature max (°C)	48 (at pH 8.5)	42	45-50	45-50
Core lipids	C ₂₀ -C ₂₀ , C ₂₀ -C ₂₅ DGE	NR	NR	NR
Intact membrane polar lipids:				
phospholipids:	PG, PGP-Me	PG, PGP-Me	PG, PGP-Me, PA	PG, PGP-Me
glycolipids:	-	1, unidentified	S-DGD-1	S-DGD-1
sulfolipids:	-	PGS		PGS (V)
Respiratory lipoquinones	MK-8:8	NR	NR	NR
DNA G+C (% genomic)	51.5 (type strain)	58.4 (type strain)	58.8 (type species)	59.7-62.7 (4 species)
Type of hypersaline habitat	soda lakes	salt lake	solar saltern	solar saltern, saline soils

480 NR, not reported; (V), variable property in different species of the same genus; * actual final pH values were not measured;
481 ** reported for the type species, but not verified in any further research; PA, phosphatidic acid; PGS, phosphatidylglycerol
482 sulfate; S-DGD-1, sulfated mannosyl glucosyl diether; other abbreviations (see **Table 2**).

483
484
485
486
487
488
489
490
491

Table 4. *Natronaeroarchaeum aerophilus*: protologue

Parameter	Species: <i>Natronaeroarchaeum aerophilus</i> sp. nov.
Author	Dimitry Y. Sorokin
Species name	<i>aerophilus</i>
Genus name	<i>Natronaeroarchaeum</i>
Specific epithet	<i>aerophilus</i>
Species status (SPST)	sp. nov.
Etymology	a.e.ro'phi.lus Gr. masc. n. <i>aer</i> , air; N.L. masc. adj. <i>philus</i> (from Gr. masc. adj. <i>philos</i>), friend, loving; N.L. masc. adj. <i>aerophilus</i> , air-loving.
Description of the new taxon	The cells are angular, flat, polymorphic coccoids or rods, mostly nonmotile, variable in size from 1 to 3 µm. The cells lyse in hypotonic solutions below 1 M NaCl, produces red carotenoids. The core membrane diether lipids are composed of C ₂₀ -C ₂₀ DGE (archaeol) and C ₂₀ -C ₂₅ DGE (extended archaeol). The polar lipid head groups include phosphatidylglycerolphosphate methyl ester (PGP-Me) and phosphatidylglycerol (PG). The dominant respiratory quinone is MK-8:8. Saccharolytic with limited substrate profile including several starch-like alpha-glucans, levan, maltose, trehalose and cellobiose. Capable of anaerobic sulfogenic growth with glucose and maltose as electron donors and sulfur as electron acceptor. Ammonium, urea and yeast extract serve as the N-source. Oxidase and catalase positive. Maximum growth temperature is 50°C. Extremely halophilic with a range of total Na ⁺ for growth from 3 to 5 M (optimum at 4 M) and moderately alkaliphilic, with a pH range for growth from 7.2 to 9.3 (optimum at 8.0-8.8). The G + C content of the DNA is 61.0% (genome of the type strain). Habitat - aerobic sediments and brines of hypersaline soda lakes. The type strain (AArc-St1-1 ^T =JCM 32519 ^T =UQM 41458 ^T) was isolated aerobic sediments of hypersaline soda lakes in Kulunda Steppe (Altai, Russia). The species also includes other three closely related strains isolated from the same area. The draft genome of type strain is deposited in the GenBank under accession number JAKRVY000000000.
Authors	Dimitry Y. Sorokin, Alexander G. Elcheninov, Tatjana V. Khizhniak, Michel Koenen, Nicole J. Bale, Jaap S. Sinninghe Damsté, Ilya V. Kublanov
Title	<i>Natronocalculus amylovorans</i> gen. nov., sp. nov., and <i>Natronaeroarchaeum aerophilus</i> sp. nov., dominant culturable amylolytic natronoarchaea from hypersaline soda lakes in southwestern Siberia.
Journal	Systematic and Applied Microbiology
Corresponding author	Dimitry Y. Sorokin
E-mail of corresponding author	soroc@inmi.ru; d.sorokin@tudelft
Designation of the type strain	AArc-St1-1
Strain collection numbers	JCM 32519 ^T =UQM 41458 ^T
16S rRNA gene accession numbers	MG584707- MG584709; ON003450
Genome accession numbers	JAKRVY000000000 (type strain)
Genome status	Draft
G+C, %	61.0 (genome of type strain)
Country of origin	Russian Federation
Region of origin	Altai region
Date of isolation	2016
Source of isolation	Surface aerobic sediments from hypersaline soda lakes
Sampling dates	2015-07-07
Geographic location	S-W Siberia, Kulunda Steppe; southern Russia
Latitude	51°39' N; 49°10' N; 48°14' N
Longitude	79°48' E; 46°39' E; 46°35' E
Depth	0-2 cm
Temperature of the sample	20°C
pH of the sample	10-11
Salinity of the sample	18-36%
Number of strains in study	4
Source of isolation of non-type strains	Surface aerobic sediments and brines from hypersaline soda lakes, S-W Siberia, Kulunda Steppe; southern Russia
Growth medium, incubation conditions	4 M total Na ⁺ , pH 9; incubation - 37°C; starch as substrates; aerobic
Conditions of preservation	Deep freezing in 15% glycerol (v/v)
Gram stain	Negative
Cell shape	Pleomorphic flat coccoids
Cell size	0.8-2 µm in diameter
Motility (MOTY)	Mostly nonmotile
Sporulation	None
Colony morphology	Flat, compact, max. 2 mm, red
Temperature range for growth	nd
Lowest temperature for growth	nd
Highest temperature for growth	50
Optimal temperature for growth	37-40
Lowest pH for growth	7.2
Highest pH for growth	9.3
Optimum pH for growth	8-8.8
pH category	Moderately alkaliphilic
Lowest Na ⁺ concentration for growth	3.0 M
Highest Na ⁺ concentration for growth	5.0 M
Optimum salt concentration for growth	4.0 M total Na ⁺
Other salts important for growth	KCl; Na-carbonates
Salinity category	Extremely halophilic
Relation to oxygen	Facultatively anaerobic
O ₂ conditions for strain testing	Fully aerobic
Carbon source used (class)	Carbohydrates

Specific compounds	Starch-like alpha-glucans, levan
Nitrogen source	Ammonium, urea, yeast extract
Terminal electron acceptor	O ₂ and S ₈
Energy metabolism	Chemoorganotrophic
Phospholipids	Core membrane lipids are C ₂₀ -C ₂₀ DGE (archaeol) and C ₂₀ -C ₂₅ DGE (extended archaeol). Polar head groups are phosphatidylglycerophosphate methylester (PGP-Me) and phosphatidylglycerol (PG)
Respiratory lipoquinones	MK-8:8
Glycolipids (GLYC)	-
Habitat (HABT)	Hypersaline soda lakes
Extraordinary features (EXTR)	Narrowly specialized amylolytics

493

494

495 **Table 5.** *Natronocalculus amylovorans*: protologue

Parameter	Genus: <i>Natronocalculus</i> gen. nov.	Species: <i>Natronocalculus amylovorans</i> sp. nov.
Author (AUTE)	Dimitry Y. Sorokin	
Species name (SPNA)		<i>amylovorans</i>
Genus name (GENA)	<i>Natronocalculus</i>	
Specific epithet (SPEP)	-	<i>amylovorans</i>
Species status (SPST)	-	sp. nov.
Etymology (GETY/SPTY)	Na.tro.no.cal'cu.lus N.L. neut. n. <i>natron</i> , arbitrarily derived from the Arabic n. natrun or natron soda; L. masc. n. <i>calculus</i> , pebble, gravel; N.L. masc. n. <i>Natronocalculus</i> , soda loving pebble-shaped cells	a.my.lo.vo'rans. Gr. neut. n. <i>amylon</i> , starch; L. inf. v. <i>vorare</i> , to devour; N.L. part. adj. <i>amylovorans</i> , eating starch
Type species of the genus (TYPE)	<i>Natronocalculus amylovorans</i>	yes
Description of new taxon	Obligately aerobic and organotrophic member of <i>Halobacteriales</i> narrowly specialized on utilization of starch-like polymers. Core lipids are dominated by archaeol and extended archaeol with PGP-Me and PG polar groups. Glycolipids are not present. MK-8:8 is the dominant lipoquinone. Extremely halophilic and moderately alkaliphilic inhabitants of hypersaline soda lakes. The three-letter abbreviation is Ncl.	The cells are angular, flat, polymorphic coccoids or rods, nonmotile, from 0.8 to 2.0 µm. The cells lyse in hypotonic solutions below 1 M NaCl. Colonies are orange-red. The core membrane diether lipids include of C ₂₀ -C ₂₀ DGE (archaeol) and C ₂₀ -C ₂₅ DGE (extended archaeol). The polar lipid head groups consists of phosphatidylglycerolphosphate methyl ester (PGP-Me) and phosphatidylglycerol (PG). The dominant respiratory menaquinone is MK-8:8. Obligately aerobic saccharolytic with limited substrate profile including several starch-like alpha-glucans, inulin, maltose, trehalose, cellobiose and glycerol. Ammonium and yeast extract serve as the N-source. Oxidase and catalase positive. Maximum growth temperature is 48°C. Extremely halophilic with a range of total Na ⁺ for growth from 3 to 5 M (optimum at 4 M) and moderately alkaliphilic, with a pH range for growth from 7.2 to 9.3 (optimum at 8.5-8.8). The G + C content of the DNA is 51.5% (genome of the type strain). Habitat - aerobic sediments and brines of hypersaline soda lakes. The type strain (AArc-St2 ^T =JCM 32475 ^T =UQM 41459 ^T) was isolated from aerobic sediments of hypersaline soda lakes in Kulunda Steppe (Altai, Russia). The species also includes 2 other closely related strains isolated from the same area. The draft genome of type strain is deposited in the GenBank under accession number JAKRVX000000000.
Authors (AUT)	Dimitry Y. Sorokin, Alexander G. Elcheninov, Tatjana V. Khizhniak, Michel Koenen, Nicole J. Bale, Jaap S. Sinninghe Damsté, Ilya V. Kublanov	
Title (TITL)	<i>Natronocalculus amylovorans</i> gen. nov., sp. nov., and <i>Natraneroarchaeum aerophilus</i> sp. nov., dominant culturable amylolytic natronoarchaea from hypersaline soda lakes in southwestern Siberia.	
Journal (JOUR)	Systematic and Applied Microbiology	
Corresponding author (COAU)	Dimitry Y. Sorokin	
E-mail of corresponding author (EMAU)	d.sorokin@tudelft; soroc@inmi.ru	
Strain collection numbers (COLN)	-	JCM 32475; UQM 41459
16S rRNA gene accession number (16 SR)	-	MG584710; ON000203; ON000205
Genome accession numbers		JAKRVX000000000
Genome status (GSTA)		Draft
GC mol % (GGCM)	-	51.5 (genome type strain)
Country of origin (COUN)	Russian Federation	Russian Federation
Region of origin (REGI)	-	Altai region
Date of isolation (DATI)	-	2016
Source of isolation (SOUR)	Hypersaline soda lakes	Surface sediments and brines of hypersaline soda lakes in southwestern Siberia
Sampling dates (DATS)	2015	2015
Geographic location (GEOL)	S-W Siberia	S-W Siberia
Latitude (LATI)	-	51°39' N; 49°10' N; 48°14' N
Longitude (LONG)	-	79°48' E; 46°39' E; 46°35' E
Depth (DEPT)		0-2 cm
Temperature of the sample (TEMS)		20°C
pH of the sample (PHSA)		10-11
Salinity of the sample (SALS)		18-36%
Number of strains in study (NSTR)	3	3
Source of isolation of non-type strains (SAMP)	-	Surface sediments and brines of hypersaline soda lakes in southwestern Siberia
Growth medium, incubation conditions (CULT)		4 M total Na ⁺ , pH 9; incubation - 37°C; starch as substrates; aerobic
Conditions of preservation (PRES)	Deep freezing in 15% glycerol (v/v)	
Gram stain (GRAM)	negative	
Cell shape (CSHA)	Pleomorphic, from flat irregular coccoids	
Cell size (CSZI)	-	0.8-2 µm in diameter

Motility (MOTY)	-	nonmotile
Sporulation (SPOR)	none	
Colony morphology (COLM)		Pink-orange, up to 2 mm, flat
Temperature range for growth (TEMR)		
Lowest temperature for growth (TEML)		
Highest temperature for growth (TEMH)		48 (at pH 8.5)
Optimal temperature for growth (TEMO)		37-40°C
Lowest pH for growth (PHLO)		7.2
Highest pH for growth (PHHI)		9.3
Optimum pH for growth (PHOP)		8.5-8.8
pH category (PHCA)	alkaliphile (optimum > 8.5)	
Lowest NaCl concentration for growth (SALL)	3.0 M total Na ⁺	
Highest NaCl concentration for growth (SALH)	5 M total Na ⁺	
Optimum salt concentration for growth (SALO)	4.0 M total Na ⁺	
Other salts important for growth	Sodium carbonates	
Salinity category (SALC)	extremely halophilic	
Relation to oxygen (OREL)	aerobe	
O ₂ conditions for strain testing (OCON)	aerobic	
Carbon source used (class) (CSUC)	carbohydrates	
Specific compounds (CSUC)	Starch-like alpha glucans and inulin	
Nitrogen source (NSOU)	Ammonium, yeast extract	
Terminal electron acceptor (ELAC)	O ₂	
Energy metabolism (EMET)	chemoorganotrophic	
Phospholipids (PHOS)	Core membrane lipids are archaeol (C ₂₀ -C ₂₀ DGE) and extended archaeol (C ₂₀ -C ₂₅ DGE) Polar lipids are phosphatidylglycerophosphate methyl ester (PGP-Me) and phosphatidylglycerol (PG)	
Glycolipids (GLYC)	-	
Respiratory lipoquinones	MK8:8	
Habitat (HABT)	Hypersaline soda lakes	

496

497

498

499 **Legends to the figures**

500

501 **Fig. 1.** Cell morphology (phase contrast microphotographs) of starch-utilizing natronoarchaea growing
 502 aerobically at 4 M total Na⁺, pH 9 and 37°C. **(a-d)**, group 1, including strains AArc-St1-1^T, AArc-St1-
 503 2, AArc-St1-3 and AArc-lev11, respectively. **(e-f)**, group 2, including strains AArc-St2^T and AArc-
 504 in2.

505 **Figure 2.** Phylogeny of amylolytic natronoarchaea.

506 **(a)** Figure X. **(a)** 16S rRNA gene sequence-based maximum-likelihood phylogenetic tree, showing the position
 507 of AArc-St2^T and AArc-St1-1^T (in bold) within the *Halobacteria* class. The black circles at nodes indicate that
 508 the percentage of corresponding support values was above 50. *Archaeoglobus fulgidus* VC-16^T, *Methanocella*
 509 *paludicola* SANAET, *Methanothermobacter thermautotrophicus* Delta H^T were used as an outgroup (not shown).
 510 Species in clusters: I (*Halodesulfurarchaeum*, *Halanaeroarchaeum*, *Halarchaeum*, *Halobacterium*,
 511 *Salarchaeum*, *Halocalculus*); II (*Halomicroarcula*, *Haloarcula*, *Halorientalis*, *Halorhabdus*, *Halococcoides*,
 512 *Halapricum*, *Salinirussus*, *Halovenus*); III (*Haloglomerus*, *Natronomonas*, *Halosegnis*, *Salinirubellus*,
 513 *Halomarina*, *Halocatena*); IV (*Halopelagius*, *Haloferax*, *Halogeometricum*, *Haloquadratum*, *Halobellus*); V
 514 (*Halococcus*, *Halalkalicoccus*, *Haloarchaeobius*, *Halorubellus*); VI (*Halorussus*, *Halomicrococcus*,
 515 *Haladaptatus*); VII (*Saliphagus*, *Natribaculum*, *Halovarius*, *Natronococcus*, *Halovivax*, *Natronobiforma*,
 516 *Halostagnicola*, *Natronobacterium*, *Halopiger*, *Halobiforma*, *Natrarchaeobaculum*, *Natronolimnohabitans*,
 517 *Natronolimnobiobius*, *Natronorubrum*, *Natrinema*, *Haloterrigena*, *Natricalba*, *Natrarchaeobius*, *Salinadaptatus*);
 518 VIII (*Halosimplex*, *Salinibaculum*, *Halosiccatus*, *Halomicrobium*); IX (*Salinigranum*, *Haloplanus*, *Halobium*,
 519 *Halegenticoccus*, *Halogramum*, *Haloprofundus*, *Halolamina*); X (*Halalkaliarchaeum*, *Halopenitus*, *Halorubrum*,
 520 *Haloparvum*).

521 **(b)** Maximum likelihood phylogenetic tree based on concatenated alignment of 122 conserved archaeal proteins
 522 and showing position of strains AArc-St2^T and AArc-St1-1^T (in bold) within the class *Halobacteria*. The branch
 523 lengths correspond to the number of substitutions per site with corrections associated with the models. The black
 524 circles at nodes indicate that the percentage of corresponding support values was above 50. *Archaeoglobus*
 525 *fulgidus* VC-16^T, *Methanocella paludicola* SANAET and *Methanothermobacter thermautotrophicus* Delta H^T
 526 were used as an outgroup (not shown). Species in clusters: I (*Halanaeroarchaeum*, *Halodesulfurarchaeum*,
 527 *Halarchaeum*, *Halobacterium*, *Salarchaeum*, *Halocalculus*); II (*Halalkalicoccus*, *Halorussus*, *Halomicrococcus*,
 528 *Haladaptatus*); III (*Haloarchaeobius*, *Halorubellus*, *Halovivax*, *Saliphagus*, *Natronobiforma*, *Halostagnicola*,
 529 *Natrarchaeobius*, *Natrarchaeobaculum*, *Salinadaptatus*, *Halopiger*, *Natronolimnobiobius*, *Natronobacterium*,
 530 *Halobiforma*, *Natricalba*, *Natronococcus*, *Natrinema*, *Haloterrigena*, *Natronorubrum*, *Natronolimnohabitans*);
 531 IV (*Halococcus*, *Halocatena*, *Halomarina*, *Natronomonas*, *Haloglomerus*, *Halosegnis*, *Halorientalis*, *Halapricum*,
 532 *Salinirussus*, *Salinibaculum*, *Halovenus*, *Halosimplex*, *Halococcoides*, *Halorhabdus*, *Halomicrobium*,
 533 *Halosiccatus*, *Halomicroarcula*, *Haloarcula*); V (*Haloplanus*, *Salinirubrum*, *Haloprofundus*, *Halegenticoccus*,
 534 *Halogramum*, *Salinigranum*, *Haloferax*, *Halopelagius*, *Halogeometricum*, *Halobellus*, *Haloquadratum*).

535

536

537 **Fig. 3.** pH profiles for growth with soluble starch in amylolytic natronoarchaea at 4 M total Na⁺ and
 538 37°C. Actual final pH are shown. The results are average from two parallel incubations.




Fig.1

Fig. 2a





Fig. 2b






Fig. 3



Click here to access/download
e-Component
supplementary file.pdf