

NEAT (North East Atlantic Taxa):

South Scandinavian marine & maritime Chelicerata & Uniramia

Check-List

compiled at TMBL (Tjärnö Marine Biological Laboratory) by:

Hans G. Hansson 1990-04-11 / small revisions until Nov. 1994, when it for the first time was published on Internet.
Republished as a pdf file February 1996 and again August 1998.

Citation suggested: Hansson, H.G. (Comp.), 1998. NEAT (North East Atlantic Taxa): South Scandinavian marine & maritime Chelicerata & Uniramia. Check-List. Internet pdf Ed., Aug. 1998. [<http://www.tmbi.gu.se>].

Denotations: (TM) = Genotype @ = Associated to * = General note

N.B.: This is one of several preliminary check-lists, covering S. Scandinavian marine animal (and partly marine protocist) taxa. Some financial support from (or via) NKMB (Nordiskt Kollegium för Marin Biologi), during the last years of the existence of this organisation (until 1993), is thankfully acknowledged. The primary purpose of these checklists is to facilitate for everyone, trying to identify organisms from the area, to know which species that earlier have been encountered there, or in neighbouring areas. A secondary purpose is to facilitate for non-experts to find as correct names as possible for organisms, including names of authors and years of description. So far these checklists are very preliminary. Due to restricted access to literature there are (some known, and probably many unknown) omissions in the lists. Certainly also several errors may be found, especially regarding taxa like Plathelminthes and Nematoidea, where the experience of the compiler is very rudimentary. The credibility of some other lists, containing taxa like e.g. Porifera, where, at least in certain subtaxa, taxonomic confusion seems to prevail, is certainly also suboptimal so far.

The compiler must admit his almost definite practical inexpérience of Chelicerata and Uniramia, so I guess that this list is very much more incomplete than the lists dealing with marine animals s.str. The borderline between animals living more or less permanently in the marine littoral and temporary guests in this environment is also more or less artificial, so some of the names in this list should probably not have been included.

Species found in South Scandinavia, as well as from neighbouring areas, chiefly the British Isles, have been considered, as some of them may show to have a slightly more northern distribution, than what is known today. However, species with a typical Lusitanian distribution, with their northern distribution limit around France or Southern British Isles, have as a rule been omitted here, albeit a few species with probable northern limits around the British Isles are listed here until distribution patterns are better known. Arctic Taxa, known only from areas north of N Norway are probably not as well represented in this list as they should be. The compiler would be very grateful for every correction of presumptive lapses and omissions an initiated reader could make.

N.B.: This document is intended to be completely unavailable regarding nomenclature within the meaning of ICZN.

ARTHROPODA Latreille,1829

* Taxa within the remaining sub-phylum, Crustacea, are treated in a separate list.

CHELICERATA Heymons,1901

MEROSTOMATA Woodward,1866

XIPHOSURIDA R. & E. Richter,1924
= LIMULIDA "Dolkvensar"

Xiphosuridae
= Limulidae

Xiphosurinae Pocock,1902

Limulus O.F. Müller,1785
= *Xiphosura* Gronovius,1764 (Is this name rejected for any reason?)

L. polyphemus (Linnaeus,1758)
* American N. Atlantic species, found now and then in Europe
S. Baltic, Skagerrak, North Sea, W. N. Atlantic

ARACHNIDA Cuvier,1812 "Spindeldjur"

* Only "marine families" included

ARANAEAE Clerck,1758 "Spindlar"

* There are no strictly marine spiders in Europe; those in the following list may be found intertidally.

ENTELEGYNAE

Araneoidea

Linyphiidae Blackwall,1859
= Ergonidae Gerhardt,1923
= Micryphantidae Bertkau,1872 "Tackvävare"

Cnephalocotes Simon,1884

C. curtis Simon,1884?
England

Erigone Audouin, in Savigny,1826

E. arctica (White,1852) *maritima* Kulczynski,1902
Böhlsän (e.g. Saltö)-Uppland, Norway, Finland, Denmark, Germany, Belgium, Br. Isles, Arctic shores etc.

E. longipalpis (Sundevall,1830)
Skåne, Norway, Finland, Denmark, Germany, Br. Isles, Arcachon, etc.

Hilarotarsus Hull,1911

H. reprobus (O. Pickard-Cambridge,1879)
* Littoral & in wrack.
Skåne, Norway, Belgium, England, France

Acanthophyma ?,19???

* Probably not a recognized genus today?

A. gowerensis (Lockett,19???)
Ireland, Britain (Gower peninsula)

Oedothorax Bertkau,1883

O. fuscus (Blackwall,1834)
Norway, Belgium, Britain

O. apicatus (Blackwall,1850)
*Littoral
Finland, Norway, Belgium

O. retosus (Westring,1851)
*Littoral & in wrack.
Uppland, Åland, Finland, Norway, Belgium

Baryphyma Simon,1884
=? Praestigia Millidge,1954

B. duffeyi (Millidge,1954)
= *Praestigia duffeyi* Millidge,1954
SE England, Belgium

B. maritimum (Crocker & Parker,1970)
Belgium

Saritatas Keyserling,1886
=? *Perimonea* Jackson,1932

P. britteni (Jackson,1913)
Britain, Belgium

Trichoncus Simon,1884

T. hackmanni Millidge,1955

S England

Lycosoidea Sundevall,1833 "Jaktspindlar"

Lycosidae Sundevall,1833 "Vargspindlar"

Arctosa Koch,1847

A. perita (Latreille,1799)
Europe, e.g. Sweden, Norway, Germany, ?Denmark

A. cinerea (J.C. Fabricius,1777) "Strandjaktspindeln"
=? *Trochosa cinerea* (J.C. Fabricius,1777)

Sweden, Norway, Denmark

Pardosa C.L. Koch,1847

= *Lycosa* Latreille,1804 (p.p.)

P. arenicola (F.O. Pickard-Cambridge,18???) *fucicola* (Dahl,1???)

= *Lycosa arenicola* F.O. Pickard-Cambridge,18?? *fucicola* Dahl,1???

* Exclusively littoral.

Uppland, Norway, England, Germany

P. pullata (Clerck,1758)

Sweden, Norway, England, Germany, S. Finland

P. purbeckensis F.O. Pickard-Cambridge,1895

Br. Isles, Belgium

P. ? fluvialis (Blackwall,1???)

= *Lycosa fluvialis* : Auctt.

Småland-Uppland, Finland

Trochosa Koch,1847

T. ruricola (De Geer,1778)

* Littoral.
Sörmland-Uppland, Norway, Britain, Germany

Dictynoidea F.O. Pickard-Cambridge,1871

Dictynidae F.O. Pickard-Cambridge,1871

Dictyna Sundevall,1833

D. major Menge,1869

N England, Scotland, Belgium

Clubionoidea (Simon,1895) Petrunkevitch,1955

Gnaphosidae Pocock,1898 "Plattbukspindlar"

Haplodrassus Chamberlin,1922

H. minor (F.O. Pickard-Cambridge,1879)

Norway, S England, Germany

Thomisoidae Sundevall,1833

Thomisidae Sundevall,1833 "Krabbspindlar"

Misumena Latreille,1804

M. varia (Clerck,1758)

= *Araneus vatus* Martyn,1793 [ex Clerck,1757]

Sweden, Norway, Arcachon

Salticoidea Blackwall,1841 "Hoppspindlar"

Salticidae Blackwall,1841 "Egentliga hoppspindlar"

Euophrays Koch, in Panzer,1834

E. browningi Millidge & Lockett,1955

SE England

Salticus Latreille,1804

S. mutabilis H. Lucas,1846

Arcachon

S. scenicus (Clerck,1758)

= *Aranea scenica* Linnaeus,1758 ex Clerck

Norway, Belgium, Arcachon

Heliophanus C.L. Koch, in Panzer,1835

H. auratus C.L. Koch, in Panzer,1835

Belgium, Arcachon

H. tribulosus Simon,1868

Belgium, Arcachon

PSEUDOSCORPIONIDA Latreille,1825 "Klokrypare"

* With the exception of gen. *Chthonius*, only species restricted to sea shores have been considered

CHTHONIINA Beier,1932

Chthonioidae Beier,1932

Chthoniidae Daday,1889

Kewochthonius Chamberlin,1929

- K. halberti* (Kew,1916)
England, Ireland, France
- Chthonius* C.L. Koch,1843
* Found also in certain inland localities.
- C. tetrachelatus* (Preyssler,1790)
Bohuslän-Ostergötland, Öland, Gotland, Åland, C. Europe, Br. Isles, Denmark, Norway
- C. kewi* Gabbott,1966
Br. Isles
- NEOBISIINA Beier,1932
- Neobisioidae Chamberlin,1931
- Neobisidae Chamberlin,1931
- Neobisium Chamberlin,1931
- N. (Neobisium) maritimum* (Leach,1812)
W. & SW. Br. Isles, Ireland, W. France
- CHELIFERINA
- Cheliferoidea Chamberlin,1931
- Cheliferidae Hagen,1879
- Cheliferinae Simon,1879
- Dactylochelifer* Beier,1932
- D. latreilli* (Leach,1817)
@ *Ammophila arenaria* & *Elymus pycnanthus*
Br. Isles, E. & S. Sweden (e.g. Uppland), Denmark, Finland, W. & Central Europe; e.g. N. Germany
- ACARI Leach,1817 "Kvalster"
- ACTINOTRICHIDA
- ASTIGMATIDA
- Hemisarcopoidea
- Hyadesiidae* Halbert,1915
= *Glyphaeidae*
- Hyadesia* Mégnin,1889 (TM) *Hyadesia uncinifer* Mégnin,1889 - Tierra del Fuego
= *Lentungula* Michael,1893
- H. fusca* (Lohmann,1894)
= *Lentungula fusca* Lohmann,1894
Baltic, Bohuslän, W Ireland, North Sea shores
- H. algivorus* (Michael,1893)
= *Lentungula algivorus* Michael,1893
S England
- Algophagidae
- Algophagus* Hughes,1955
- Neohyadesia ??,19??
- Acaroidea
- Acandiae Oudemans,1904 "Skabbkvalster"
- Pontoppidania Oudemans,1923
- P. littoralis* (Halbert,1920)
= *Tyrophilus littoralis* Halbert,1920
@ in moist decaying seaweeds
W Ireland
- Rhizoglyphus* Claparède,1869
- R. sp.*
Britain
- Tyrophagus* Oudemans,1924
- T. sp.*
Britain
- Suidasiidae
- Suidasia
- PSOROPTINA
@ with few exceptions parasites of birds or mammals
- Psoroptoidea
- Sarcoptidae
- Sarcoptes* Latreille,1802
- ?S. sp. : Jacobsen,1966
@ skin of *Phoca vitulina*
German North Sea, Netherlands
- PROSTIGMATIDA Kramer,1887
= TROMBIDIIFORMES
= ACTINEDIDA
- Nautarachnidiae
* What infraorder & superfamily?
- Nautarachna* Moniez,1888
- Thysidae
* What infraorder & superfamily?
- Thyopsis* Piersig,1899
- ENDEOSTIGMATINA Grandjean,1937
- Nanorchestidae Grandjean,1937
- Nanorchestes Topsent & Trouessart,1890
- N. amphibius* Topsent & Trouessart,1890
Ireland, Isle of Whight
- Bimichaelidae
= Pachyanathidae
= Alvidae G. Canestrini,1891
- Pachygnathidae Dugès,1834
- P. sp.*
Britain
- Alicus* Berlese,1890
= *Alycus* Koch,1842, non Rafinesque,1815 (Coleoptera)
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized?
- A. (Leptalicus) oblongus* Halbert,1920
W Ireland
- A. latus* Halbert,1920
W Ireland
- EUPODINA
- Bdelloidea
- Bdellidae Dugès,1834
- Bdella* Latreille,1795
- B. interrupta* G.O. Evans,1952
* In crevices of the rocky littoral.
Br. Isles
- B. capillata* Kramer,1881
Scandinavia, Germany, Britain, Clare Island, Italy
- B. decipiens* Thorell,1871
= *Bdella vulgaris* (Hermann,1???) var. *littoralis* Moniez,1890
Sweden, W Ireland, Arcachon, Siberia
- B. sanguinea* Trouessart,1???
?Germany
- B. littoralis* (Linnaeus,1758)
= *Molgus littoralis* (Linnaeus,1758) (TM of *Molgus* Dujardin,1842)
= *Molgus acutus* (Thorell,1871)
= *M. villosus* (Kramer,1883)
= *M. basteri* (Michael,1896)
* An up to 3.5 mm long, shining red species, on decaying seaweed.
Scotland, Norway, Ireland
- Cyta* von Heyden,1826
- C. latirostris* (Hermann,18??)
W Ireland
- Hoplomolgus Berlese,1923
- H. sp.*
Britain
- Neomolgus* Oudemans,1938
- N. sp.*
Britain
- Eupodoidea
- Eupodidae C.L. Koch,1842
- Eupodes* Koch, in Panzer,1835
- E. halophilus* Halbert,1920
England, W Ireland
- Penthaleidae Oudemans,1931
- Hylotydeus* Berlese,1891
- H. hydrodromus* (Berlese & Trouessart,1889)
= *Notophallus hydrodromus* Berlese & Trouessart,1889
Br. Isles
- H. hydrodromus* (Berlese & Trouessart,1889) var. *albolineatus* Halbert,1915
W Ireland
- Chromatydae Berlese,1903
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
- C. ovatus* (C.L. Koch,1838)
= *Penthalodus ovatus* C.L. Koch,1838
@ among *Sembalanus*, *Fucus spiralis* & *F. vesiculosus*
W Norway, Germany, Ireland, Britain, Italy
- Penthalodiidae Sig Thor,1933
- Penthalodes Murray,1877
- P. sp.*
Britain
- Rhagidiidae Oudemans,1922
- Foveachetidae Zacharda,1???
= *Rhagidia* Thorell,1871
- F. halophila* (Laboulbène,1851)
= *Gammarus halophilus* Laboulbène,1851
= *Nornera halophila* Moniez,1889
W Ireland, ?French Atlantic coast
- Robustochetes* Zacharda,1???
- R. sp.*
Britain
- Tydeoidea

<i>Tydeidae</i> Kramer,1877	
<i>Lasiotydaeus</i> Berlese,1908 (TM <i>L. glycophagimus</i> Berlese,1908?)	
<i>L. (Melanotydaeus</i> Berlese,1910) <i>brevistylus</i> Halbert,1920	
W Ireland	
<i>Retetydeus</i> Thor,1931	
<i>R. sp.</i>	
Britain	
<u>Nematalycoidae</u>	
<u>Nematalycidae</u>	
Halacaroidea	
Halacaridae Murray,1877 "Havskvalster"	
Rhomhognathinae Viets,1927	
<i>Isohactrus</i> Newell,1947	
<i>I. levis</i> (Viets,1927)	
= <i>Rhombognathus levis</i> Viets,1927	
Baltic, Bohuslän, North Sea	
<i>I. setosus</i> (Lohmann,1889)	
= <i>Aletes setosus</i> Lohmann,1889	
Baltic, Halland, North Sea	
<i>I. uniscutatus</i> (Viets,1939)	
= <i>Rhombognathus uniscutatus</i> Viets,1939	
Baltic, North Sea	
<i>I. hartmanni</i> Bartsch,1972	
* Not included in Green & Macquitty, 1987	
Öresund, Helsingør	
<i>I. ungulatus</i> Bartsch,1975	
Roscoff, Falmouth, Isle of Man, Pembrokeshire	
<i>Metarhombognathus</i> Newell,1947	
<i>M. armatus</i> (Lohmann,1893)	
= <i>Rhombognathus mollis</i> Viets	
Baltic, North Sea	
<i>M. nudus</i> (Viets,1928)	
= <i>Rhombognathus conectus</i> Viets	
Cullercoats, Scotland, Murmansk	
<i>Rhombognathides</i> Viets,1927	
<i>R. mucronatus</i> (Viets,1927)	
Bornholm, North Sea, Norway	
<i>R. pascens</i> (Lohmann,1889)	
Baltic, Bohuslän, North Sea, Norway	
<i>R. seahami</i> (Hodge,1860)	
= <i>Pachygrathus seahami</i> Hodge,1860	
Baltic, Bohuslän, Norway, North Sea, Kola peninsula, Novaja Zemlja, Iceland, Shetlands, Ireland, E & S Britain, E USA	
<i>R. spinipes</i> (Viets,1933)	
Baltic, North Sea	
<i>R. merrimani</i> Newell,1947	
Bohuslän, Norway, England, Wales, Ireland	
<i>R. exoplus</i> (Trouessart,1901)	
* Not included in Green & Macquitty, 1987	
Bretagne, North Sea	
<i>R. trionyx</i> (Trouessart,1899)	
* Not included in Green & Macquitty, 1987	
Bretagne	
<i>Rhombognathus</i> Trouessart,1888	
<i>R. notops</i> (Gosse,1855)	
Baltic, Bohuslän, North Sea	
<i>R. subtilis</i> Bartsch,1975	
Norway, Spitzbergen, Cullercoats, Northumberland	
<i>R. lioyng</i> Trouessart,1900	
Baltic, Bergen	
<i>R. magnirostris</i> Trouessart,1889	
Kullen, Bergen, French Mediterranean	
<i>R. cetratus</i> Bartsch,1975	
Elbe-estuary	
<i>R. cryptorhynchus</i> Trouessart,1901	
* Not included in Green & Macquitty, 1987	
Bretagne	
<i>R. plumifer</i> ?,???	
?	
<i>R. intermedius</i> Schulz,1933	
Kieler Bucht, Helgoland	
<i>R. procerus</i> ?,???	
?	
<u>Actacarinae</u> Viets,1939	
<i>Actacarus</i> Schulz,1936	
<i>A. obductus</i> Bartsch,1977	
* Not included in Green & Macquitty, 1987	
Bohuslän, North Carolina	
<i>A. pygmaeus</i> Schulz,1936	
* Not included in Green & Macquitty, 1987	
Kieler Bucht, North Sea, Mediterranean, Black Sea	
<i>A. bacescui</i> Kennerth-Ionescu,1970	
* Not included in Green & Macquitty, 1987	
N Spain-Mediterranean-Black Sea	
<i>Halacarinae</i> Viets,1927	
<i>Agauopsis</i> Viets,1927	
<i>A. brevipalpus</i> (Trouessart,1889)	
= <i>Agae brevipalpus</i> Trouessart,1889	
England, Wales, Ireland, France	
<i>A. tricuspid</i> Benard,1962	
= <i>A. vidiae</i> Krantz,1970	
England, Bretagne	
<i>A. aculeata</i> (Trouessart,1896)	
* Not included in Green & Macquitty, 1987	
Biscay in 1410m among corals	
<i>A. microrhyncha</i> (Trouessart,1889)	
* Not included in Green & Macquitty, 1987	
Calais	
<i>A. hirsuta</i> (Trouessart,1889)	
* Not included in Green & Macquitty, 1987	
Mediterranean	
<i>Agae</i> Lohmann,1889	
<i>A. chevreuxi</i> (Trouessart,1889)	
* Not included in Green & Macquitty, 1987	
Roscoff	
<i>A. panopae</i> (Lohmann,1893)	
* Not included in Green & Macquitty, 1987	
Monaco	
<i>Anomalohalacarus</i> Newell,1949	
<i>A. anomalus</i> (Trouessart,1894)	
* In sublittoral sediments.	
Bohuslän, Skagerrak, Engl. channel, Bretagne	
<i>A. intermedius</i> Bartsch,1976	
* Shallow water species	
North Sea, Bretagne, Ireland	
<i>A. marcandrei</i> Monniot,1967	
* In sublittoral sediments.	
Bohuslän, Skagerrak, Bretagne, Yorkshire	
<i>A. minutus</i> Bartsch,1976	
* Shallow water species.	
Baltic, North Sea, Bretagne, Yorkshire	
<i>A. poizatii</i> Bartsch,1985	
* In sublittoral sediments.	
Bohuslän, Yorkshire	
<i>A. acenomus</i> Bartsch,1985	
* In sublittoral sediments.	
Bohuslän, Yorkshire	
<i>A. arenarius</i> Bartsch,1978	
* Shallow water species. Not included in Green & Macquitty, 1987	
North Sea	
<i>A. similis</i> Bartsch,1976	
* Not included in Green & Macquitty, 1987	
?	
<i>A. septentrionalis</i> Bartsch,1991	
* In sublittoral sediments.	
Bohuslän	
<i>Arhodeoporus</i> Newell,1947	
<i>A. gracilipes</i> (Trouessart,1889)	
Bohuslän, North Sea, Tromsö, England	
<i>A. minor</i> Bartsch,1977	
Bretagne, Yorkshire, Ireland	
<i>Coloboceras</i> Trouessart,1889	
<i>C. drachi</i> Monniot,1962	
Ireland	
<i>C. longiusculus</i> Trouessart,1889	
* Not included in Green & Macquitty, 1987	
Bohuslän, French atlantic coast	
<i>C. koehleri</i> Trouessart,1896	
* Not included in Green & Macquitty, 1987	
Biscay in 1410m among corals	
<i>Copidognathus</i> Trouessart,1888	
<i>C. brevirostris</i> Viets,1927	
Baltic, Bohuslän, Bergen, North Sea	
<i>C. brifaciuss</i> Bartsch,1989	
Bohuslän	
<i>C. dentatus</i> Viets,1940	
Grimsby, Kent	
<i>C. fabricii</i> (Lohmann,1889)	
= <i>Halocarus fabricii</i> Lohmann,1889	
Baltic, Halland, Bohuslän, North Sea, Tromsö, Iceland, Faeroes, Ireland, W France, Mediterranean, Black Sea, Azores, Cape Verde	
<i>C. gibbus</i> (Trouessart,1889)	
Helgoland, Engl. Channel	
<i>C. granulatus</i> (Hodge,1863)	
England, French atlantic coast	
<i>C. glyptoderma</i> Trouessart, ?	
Plymouth??	

- C. hartwigi* Bartsch,1979
Irish Sea, Bermuda
- C. lamellosus* (Lohmann,1893)
= *C. tabellio* (Trouessart,1894)
Engl. Channel, Yorkshire, Ireland
- C. latisetus* Viets,1940
Bohuslän, North Sea, Bretagne, Yorkshire, Kent, Ireland
- C. loricifer* André,1946
Baltic, Engl. Channel
- C. magnipalpus* (Police,1909)
* Not included in Green & Macquitty, 1987
Bohuslän, Mediterranean
- C. oculatus* (Hodge,1863)
Baltic, Kattegatt, Bohuslän, North Sea, Engl. Channel
- C. poucheti* (Trouessart,1894)
= *Halacarus pouchetti* Trouessart,1894
Iceland, Labrador
- C. remipes* (Trouessart,1894)
Engl. Channel, Ireland
- C. reticulatus* (Trouessart,1893)
= *Halacarus reticulatus* Trouessart,1893
Bohuslän (Svartskär N of Kälggrund - Bartsch, unpubl.), Norway, Spitzbergen, Iceland, Ireland
- C. rhodostigma* (Gosse,1855)
Baltic, Halland, Bohuslän, North Sea, England, Scotland, Ireland
- C. septentrionalis* (Hallert,1915)
= *Halacarus (Copidognathus) lamellosus* Lohmann,1893 var. *septentrionalis* Hallert,1915
Engl. Channel, Clew Bay
- C. stevici* Bartsch,1976
@ Ectoparasite on eggs of *Maia squinado*
Engl. Channel
- C. tectirostris* Bartsch,1979
Bohuslän, Bretagne, Yorkshire, Irish Sea, Mediterranean
- C. gracilipes* (Trouessart,1889)
* Not included in Green & Macquitty, 1987
N. Kattegatt, Bohuslän, Helgoland, Bergen
- C. caudani* (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Biscay
- C. humerosus* (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Engl. Channel
- C. latus* Viets,19??
* Not included in Green & Macquitty, 1987
Öresund, Bergen
- Halacarus* Gosse,1855
- H. bisulcus* Viets,1927
Bohuslän, Helgoland, Bergen, Iceland, Faeroes, Yorkshire, Devon, Ireland
- H. anomalus* Trouessart,1894
* Not included in Green & Macquitty, 1987
Baltic, Helgoland
- H. ctenopus* Gosse,1855 (TM)
North Sea, Shetland, Engl. Channel, Ireland
- H. actenos* Trouessart,1889
Engl. Channel, Wales, Ireland
- H. borealis* Trouessart,1893
NW Iceland
- Thalassarchna* Packard,1871
= *Halocarellus* Schulz,1933
- T. baltica* (Lohmann,1889)
Baltic, Bohuslän, North Sea
- T. basteri* (Johnston,1836)
= *Acarus basteri* Johnston,1836
= *T. verrilli* Packard,1871
= *Halacarus spinifer* Lohmann,1889
Baltic, Bohuslän, North Sea, Iceland, Murmansk, Novaja Zemlya, Greenland, Faeroes, Ireland, E & S England, France, U.S.A.
- T. capuzina* (Lohmann,1893)
= *Halacarus (Halacarus) areolatus* Hallert,1915
Baltic, Öresund, Bohuslän, North Sea, England, Scotland, Ireland
- T. petiti* (Angelier,1950)
Kieler Bucht, Bretagne, Humber estuary
- T. procura* (Viets,1927)
Baltic, Halland, Helgoland, Bretagne, Ireland
- T. southerni* (Hallert,1915)
= *Halacarus (Halacarus) southerni* Hallert,1915
Bohuslän, Yorkshire, Clew Bay, Irish Sea
- T. striatus* (Lohmann,1889)
= *Halacarus similis* Viets,1927
Kieler Bucht, Bohuslän, Bergen, Bretagne, England, Ireland
- T. subterranea* (Schulz,1933)
Baltic, North Sea, Norway, Bretagne, Essex
- T. bergensis* (Viets,1927)
* Not included in Green & Macquitty, 1987
Kieler Bucht, Bergen
- T. floridearum* (Lohmann,1889)
* Not included in Green & Macquitty, 1987
Baltic
- T. subcrispus* (Bartsch,1978)
* Not included in Green & Macquitty, 1987
?
- T. inermis* (Trouessart,1888)
* Not included in Green & Macquitty, 1987
Baltic (e.g. Greifswald), Halland, North Sea
- T. longipes* (Trouessart,1888)
* Not included in Green & Macquitty, 1987
Baltic, Bohuslän, North Sea
- T. micropectinatus* (Bartsch,1972)
* Not included in Green & Macquitty, 1987
Schleswig-Holstein
- T. hexacanthus* (Viets,1927)
* Not included in Green & Macquitty, 1987
Kieler Bucht, Öresund, Kattegatt, Bohuslän, Bergen
- T. abyssorum* (Trouessart,1896)
* Not included in Green & Macquitty, 1987
Biscay in 400-1410m among corals
- Camactognathus* Newell,1984
* Correct subfamily?
- C. borealis* Bartsch,1991
Bohuslän
- Acarochelopodinae* Bartsch,1977
- Acarochelopodia* Angelier,1954
- A. delamarei* Angelier,1954
* Not included in Green & Macquitty, 1987
Europe
- Simognathinae* Viets,1927
- Simognathus* Trouessart,1889
- S. leiomerus* Trouessart,1894
= *S. andrei* Monniot,1961
Engl. Channel, Ireland
- S. minutus* (Hodge,1863)
= *Pachygratulus sculpus* Brady,1875
Kullen (NW Skåne), Helgoland, Bergen, England, Scotland
- Acaromantis* Trouessart & Neumann,1893
- A. squilla* Trouessart & Neumann,1893
* Not included in Green & Macquitty, 1987
French atlantic coast
- A. minutus* Bartsch,1976
* Not included in Green & Macquitty, 1987
?
- Atelopsalis* Trouessart,1896
- A. tricuspidis* Trouessart,1896
* Not included in Green & Macquitty, 1987
Biscay in 1410m among corals
- Lohmannellinae* Viets,1927
- Lohmannella* Trouessart,1901
- L. falcatata* (Hodge,1863)
= *Leptogratulus falcatus* Hodge,1863
Baltic, Bohuslän, Norway, North Sea, Shetlands, Hebrides, Ireland, Britain, Iceland, W France, Mediterranean, Black Sea, E North America, (possibly cosmopolitan)
- L. kervillei* (Trouessart,1894)
Engl. Channel, Wales
- L. multiseta* Bartsch,1986
* Not included in Green & Macquitty, 1987
Bohuslän, Engl. Channel, Mediterranean
- L. norvegica* Viets,1927
* Not included in Green & Macquitty, 1987
Bergen
- Porolohmannella* Viets,1933
- P. violacea* (Kramer,1879)
* NB! Freshwater species
Widespread in Europe from Belgium to U.S.S.R.
- Scaptognathus* Trouessart,1889
- S. tridens* Trouessart,1889
Engl. Channel, French atlantic coast
- S. trouessarti* Hallert,1915
Dingle Bay (Ireland)
- S. haltezi* Trouessart,1894
* Not included in Green & Macquitty, 1987
Calais
- Limnohalacarinae* Viets,1927
- Limnohalacarus* Walter,1917
- L. wackeri* (Walter,1914)
* NB! Freshwater species
Widespread in Europe
- Parasoldanellonyx* Viets,1929
- P. parviscutatus* (Walter,1917)
* NB! Freshwater species
Belgium, England, Scotland, Switzerland, Italy, Germany, U.S.S.R.
- Soldanellonyx* Walter,1917
- S. chappuisi* Walter,1917
* NB! Freshwater species

England, Scotland, France to Roumania and U.S.S.R.

S. monardi Walter,1919

* NB! Freshwater species

England, Wales, Scotland, France to Roumania and U.S.S.R.

Porohalacarinae Viets,1933

Caspihalacarus Viets,1928

C. hyrcanus Viets,1928

* NB! Brackish water species

Wales, Caspian and Black Seas, Dnieper

Lobohalacarus Viets,1939

L. dolgarae Green,1954

* NB! Freshwater species

England

L. gallicus (Migot,1926)

* NB! Freshwater species

French Pyrenees

L. weberi (Romijn & Viets,1924)

* NB! Freshwater species

England, Scotland, Widespread in European inland waters

Porohalacarus Thor,1923

P. alpinus (Thor,1910)

* NB! Freshwater species

England, Wales, Widespread in European and N. African waters

ANYSTINA

Anystoidea

Anystidae Oudemans,1902

Chaussieria Oudemans,1937

C. sp.

Britain

Teneriftiidae

Heteroteneriffia Hirst,1926

PARASITENGONINA Oudemans,1909

Erythraeidea

Erythraeidae Oudemans,1902

Balaustium von Heyden,1826

B. sp.

Britain

Erythraeus Latreille,1806

E. sp.

Britain

Trombidioidea

Podothrombiidae

Podothrombium Berlese,1910

P. filipes (Koch,18??)

Norway, Germany, Ireland, Britain

Trombiculidae

Trombiculidae Leach,1814

Rhyncholophus Dugès,1834 (

* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?

R. rubripes Berlese & Trouessart,1889

= *R. minicatus* var. *rubripes* Berlese & Trouessart,1889

= *Ritteria hirsuta* George,1910

@ Commensal on *Semibalanus balanoides*.

W Ireland, Britain, France, ?Germany

R. norvegicus (Sig. Thor,1900)

= *Ritteria norvegica* Sig. Thor.

Norway, Ireland

R. tardus Halbert,1915

W Ireland

R. passerinii (Berlese,1904)

= *Erythraeus passerini* Berlese,1904

W Ireland, Italy

R. araneoides (Berlese,1910)

= *Achorolophus araneoides* Berlese,1910

W Ireland, Sicily

Microtrombidium Haller,1882

M. pusillum (Hermann,18??) var. *major* Halbert,1920

W Ireland

Hydryphantoidae

Hydryphantidae

@ larvae on Odonata, Dipera, Hemiptera, Plecoptera & Trichoptera

Hydryphantes Koch,1842

Hydrodromidae

@ larvae on Chaoboridae, Culicidae & Chironomidae

Hydrodroma Koch,1837

Hygrobatoidea

Pontarachnidae Sig Thor,1929

* In the littoral zone of marine waters, but probably not N of the Mediterranean.

Pontarachna Philippi,1840

Litarachna Walter,1925

Pionidae

Piona Koch,1842

ELEUTHERENGINA Oudemans,1909

RAPHIGNATHAE

Raphignathoidea

Raphignathus Dugès,1834

* This genus is not found in Hayward & Ryland (Eds.) 1990.

R. scutatus Halbert,1920

W Ireland

Stigmaeidae Oudemans,1931

Stigmæus Koch,1835

S. rhodomelas var. *fissuricola* Halbert,1920

W Ireland

Cheylostigmæus Willmann,19??

C. sp.

Britain

Tetranynchoidea

Tetranynchidae

Monoceronychus McGregor,1945

Schizotetranychus Trägårdh,1915

Temnopalnidæ

=? Phytontpalnidæ

=? Pseudolepidae

Dolichotetranychus Sayed,1938

Pseudoleptus Bruyant,1911

CRYPTOSTIGMATIDA

= ORIBATIDA

= ORIBATEI Dugès,1834

Zachvatkina ??,19??

* Correct suborder?; Family?

Z. sp. cf. stercorarii (Dubinin,1949)

@ *Stercorarius parasiticus*

British Columbia

Cymbaeremaeidae

* Which superfamily?

Cymbaeremæus Berlese,1896

C. sp.

@ *Uria aalge*

British Columbia

CIRCUMDEHISCENTIAEINA

?Superfamily?

Oribata Latreille,1802

* The species in this genus should probably go to other oribateid genera.

O. setosa C.L. Koch,18??

W. Ireland

O. quadricornuta Michael,18??

W Ireland

O. lucasii Nicolet,18??

W Ireland

Ameronothroidea

Ameronothridae Willmann,1931

Ameronothrus Berlese,1896 (TM *Eremaeus lineatus* Thorell,18??)

A. marinus Banks,18??

* On rocks in intertidal zone.

France

A. bilineatus Michael,18??

* Algidorous species restricted to littoral or estuarine situations.

Br. Isles

A. schneideri (Oudemans,1905)

* Coastal

Borkum

A. sp. aff. oblongatus Sitnikova,1975

@ *Uria aalge*

British Columbia

Fortuninia van der Hammen,1960

Hygroribates Jacot,1934

Selenoribates Strenzke,1961

Carabodoidea

Carabodidae Willmann,1931

<i>Carabodes</i> C.L. Koch, in Panzer, 1835 (TM <i>C. coriaceus</i> Koch, in Panzer, 1835)	<i>P.</i> sp. Britain
<i>C.</i> sp. Britain	<u>Oribatuloidea</u>
<i>Passalozetes</i> Grandjean, 1932	<u>Oribatulidae</u> Jacot, 1929
<u>Ceratozetoidea</u>	<i>Halaribatula</i> Schuster, 1957
<u>Ceratozetidae</u> Jacot, 1925	<i>Liebstadia</i> Oudemans, 1906 (TM <i>Notaspis similis</i> Michael, 1888)
<i>Ceratozetes</i> Berlese, 1908 (TM <i>Oribata gracilis</i> Michael, 1884)	<i>L.</i> sp. Britain
<i>C.? quadrivertex</i> (Halbert, 1920) = <i>Oribata quadrivertex</i> Halbert, 1920 W Ireland	<i>Oribatula</i> Berlese, 1895 (TM <i>Notaspis tibialis</i> Nicolet, 1855)
<i>Trichoribates</i> Berlese, 1910 (TM <i>Murcia trimaculata</i> C.L. Koch, 1836)	<i>O. saxicola</i> Halbert, 1920 W Ireland
<i>T.</i> sp. Britain	<i>O. similis</i> Michael, 18?? * Is thus identical with the genotype of <i>Liebstadia</i> Oudemans? W Ireland
?Families?	<u>Oribatulidae</u> Grandjean, 1933
<i>Chamobates</i> Hull, 1916 (TM <i>Oribata cuspidata</i> Michael, 1884)	<u>Scheloribatidae</u> Grandjean, 1933
<i>C.</i> sp. Britain	<i>Scheloribates</i> Berlese, 1908 (TM <i>Zetes latipes</i> C.L. Koch, 1844)
<i>Euzetes</i> Berlese, 1908 (TM <i>Oribata globula</i> Nicolet, 1855)	<i>S.</i> sp. Britain
<i>E.</i> sp. Britain	<u>Passalozetoidea</u>
<i>Fuscozetes</i> Sellnick, 1928 (TM <i>Oribates fuscipes</i> C.L. Koch, 1844)	<i>Scutovortex</i> Michael, 1879 (TM <i>S. sculptus</i> Michael, 18??)
<i>F.</i> sp. Britain	<i>S. spoofi</i> Oudemans, 1900 = <i>S. bilineatus</i> : Oudemans, 1901, non Michael, 18?? = <i>Ameronothrus spoofi</i> Auct. * Alivorous species restricted to littoral or estuarine situations. Finland, Br. Isles
<i>Minunthozetes</i> Hull, 1916 (TM <i>Oribata fusigera</i> Michael, 1884 = <i>Zetes semirufus</i> C.L. Koch, 1840)	<i>S. bidactylus</i> Coggi, 1900 * in maritime sand dunes Norderney, Ireland, Netherlands, Italy
<i>M.</i> sp. Britain	<i>S. corrugatus</i> Michael, 18?? = <i>Eremaeus lineatus</i> Thorell, 18?? W Ireland
<i>Mycobates</i> Hull, 1916	<i>S. perforatus</i> Berlese, 1910 Ireland
<i>M. parmeliae</i> (Michael, 1884) (TM) = <i>Oribata parmeliae</i> Michael, 1884 * Under lichens growing on granite rocks Cornwall, W Ireland	<i>S. maculatus</i> Michael, 18?? @ in <i>Lichina pygmaea</i> W Ireland
<i>Punctoribates</i> Berlese, 1908 (TM <i>Oribates punctum</i> C.L. Koch-Berl., 1840)	<u>Pelopoidea</u>
<i>P.? avenifera</i> Michael, 18?? = <i>Oribata avenifera</i> Michael, 18?? W Ireland	<u>Pelopsidae</u> Ewing, 1917
<i>Sphaerozetes</i> Berlese, 1885 (TM <i>Oribates orbicularis</i> C.L. Koch, 1836)	<i>Pelops</i> C.L. Koch, 1835, non Gistl, 1834 (Coleoptera) (TM <i>Notaspis hirsutus</i> C.L. Koch, 18??)
<i>S.</i> sp. Britain	<i>P.</i> sp. Britain
<u>Galumnoidea</u>	<i>Eupelops</i> Ewing, 1917
<i>Galumnidae</i> Grandjean, 1936	<i>E.</i> sp. Britain
<i>Galumna</i> von Heyden, 1826 (TM <i>Notaspis akatus</i> Hermann, 1804)	<i>Peloptulus</i> Berlese, 1908
<i>G.</i> sp. Britain	<u>EUPTYCTIMINA</u>
<u>Liacaroidea</u>	<i>Phitracaroidea</i>
<i>Adoristes</i> Hull, 1916 (TM <i>Oribates ovatus</i> C.L. Koch, 1840)	<i>Steganacarus</i> Ewing, 1917
<i>A.</i> sp. Britain	<i>S.</i> sp. Britain
<i>Xenillus</i> Robineau-Desvoidy, 1839 (TM <i>X. clypeator</i> R.-D., 1839 = <i>Notaspis tegeocranus</i> Hermann, 1804)	<u>NOTHRONATINA</u>
<i>X.</i> sp. Britain	<i>Nanhermannidae</i> * Which superfamily?
<u>Oppioidea</u>	<i>Nanhermannia</i> Berlese, 1913
<i>Oribella</i> Berlese, 1908 (TM <i>Notaspis pectinata</i> Michael, 1885)	<u>Nothriidea</u>
<i>O.</i> sp. Britain	<i>Camisiidae</i> Sellnick, 1928
<u>Eremaeidae</u>	<i>Camisia</i> von Heyden, 1826 (TM <i>Notaspis segnis</i> Hermann, 1804)
<i>Halozetes</i> Berlese, 1917	<i>C.</i> sp. Britain
<i>Oppia</i> C.L. Koch, 1835 (TM <i>O. nitens</i> C.L. Koch, 1835)	<i>Platynothrus</i> Berlese, 1913 (TM <i>Nothrus peltifer</i> C.L. Koch, 1840)
<i>O. maritima</i> (Willmann, 1929) @ in seaweed Cuxhaven	<i>P.</i> sp. Britain
<i>O.</i> sp. Britain	<u>Nothridae</u>
<i>Thalassozetes</i> Schuster, 1963	<i>Nothrus</i> Koch, in Panzer, 1835 * This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?
<u>Oribatelloidea</u>	<i>N. invenustus</i> Michael, 18?? @ in <i>Lichina pygmaea</i> W Ireland
<i>Oribatellidae</i> Jacot, 1925	<u>Hermannioidea</u>
<i>Oribatella</i> Banks, 1895 (TM <i>O. quadridentata</i> Banks, 1895?)	<i>Hermanniaidae</i> Sellnick, 1928
<i>O. maritima</i> v. <i>litoralis</i> Strenzke, 1952 ?	
<i>O.</i> sp. Britain	
?Family?	
<i>Parachipteria</i> van der Hammen, 1952	

- | | |
|--|--|
| <i>Hermannia</i> Nicolet,1855 (tm <i>Hermannia crassipes</i> Nicolet,1855 = <i>Nothrus gibbus</i> Koch,1840) | Britain |
| <i>H. scabra</i> (L. Koch,18??)
= <i>H. nodosa</i> Michael,18??
Finland, Sweden, Germany, Br. Isles, W Ireland, Novaja Zemlja, Siberia | <i>Pergamasus</i> Berlese,1904 |
| <i>H. reticulata</i> Thorold,18??
Netherlands, Britain, W Ireland, E Greenland, Spitsbergen, Novaja Zemlja | <i>P. sp.</i>
Britain |
| <u>ANACTINOTRICHIDA</u>
= <u>PARASITIFORMES</u> Reuter,1909
= <u>GAMASIFORMES</u>
= <u>ANACTINOTRICHIDA</u> | <i>Trachygamasmus</i> Berlese,1906 |
| <u>MESOSTIGMATIDA</u> Canestrini,1891
= <u>GAMASIDA</u> | <i>T. sp.</i>
Britain |
| <u>UROPODINA</u> Kramer,1881 | <i>Vulgarogamasus</i> Tichomirov,19?? |
| <u>Uropodidae</u> Berlese,1917 "Sköldkvalster" | <i>V. sp.</i>
Britain |
| <i>Dendrouropoda</i> Willmann,1959 | <u>DERMANYSSINA</u> |
| <i>Dentibaiulus</i> ??,19?? | <u>Dermanyssoidae</u> |
| <i>Phaulodinychus</i> Berlese,1905
= <i>Hahropoda</i> Halbert,1915 | <u>Halacaridae</u> |
| <i>P. minor</i> (Halbert,1915)
= <i>Hahropoda minor</i> Halbert,1915
= <i>Trachiropoda minor</i> Halbert,1920
* In decaying wrack.
W Ireland | <i>Halarachne</i> Allman,1847 |
| <i>P. orchesitiidarum</i> (Barrois,1887)
= <i>Uropoda orchesitiidarum</i> Barrois,1887
* In decaying wrack.
W Ireland | <i>H. halichoeri</i> Allman,1847
@ nostrils of <i>Halichoerus gryphus</i>
Baltic, Ireland |
| <i>P. repletus</i> Berlese,1905
= <i>Hahropoda interrupta</i> Halbert,1915 (tm of <i>Hahropoda</i> Halbert,1915 - Ireland) | <i>H. miroungae</i> Ferris,1925
@ <i>Phoca vitulina</i>
US Pacific |
| <i>P. littoralis</i> (Trouessart,1902)
= <i>P. berlesei</i> Halbert,1915
Ireland, Britain | <i>Orthohalarachne</i> Newell,1947
@ walruses & eared seals |
| <i>Dinychus</i> Kramer,1886 | <i>O. diminuata</i> (Doetschman,1944)
@ <i>Callorhinus ursinus</i> & <i>Zalophus californianus</i>
Bering Sound-C California |
| <i>D. sp.</i> : Halbert,1920
W Ireland | <i>O. attenuata</i> (Banks,1910)
@ <i>Callorhinus ursinus</i> & <i>Eumetopias jubata</i>
Bering Sound-Oregon |
| <i>Uropoda</i> Latreille,1806 | <u>Laelanidae</u> Berlese,1892 |
| <i>U. sp.</i>
Britain | <i>Arctoseius</i> Thor,1930
* Described from Spitsbergen |
| <i>Urosternella</i> Berlese,1918 | <i>Aspidilaelaps</i> Trägårdh,1949 |
| <u>Thinozerconidae</u> Halbert,1915 | <i>Gammaridacarus</i> Canaris,1962
@ <i>Orchestoidea</i> |
| <i>Thinozercon</i> Halbert,1915 | <i>Laelaps</i> Koch,1836
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family? |
| <i>T. michaeli</i> Halbert,1915
Ireland | <i>L. dentatus</i> Halbert,1920
W Ireland |
| <u>Polyspididae</u> Berlese,1917 | <i>Hypoaspis</i> Canestrini,1884 |
| <i>Trachytes</i> Michael,1894 | <i>H. sp.</i>
Britain |
| <i>T. sp.</i>
Britain | <i>Ololaelaps</i> Berlese,1905 |
| <u>PARASITINA</u> | <i>O. sp.</i>
Britain |
| <u>Parasitidae</u> Oudemans,1901 "Parasitkvalster" | <i>Pseudoparasitus</i> Oudemans,1902 |
| <i>Eugamasus</i> Berlese,1892
* This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? It was used as a subgenus together with <i>Gamasus</i> by Halbert,1915. | <i>P. sp.</i>
Britain |
| <i>E. immanis</i> (Berlese,1903)
* Collected from tidal drift feeding on small Oligochaetes.
Norway, Millport, Northumberland, Ireland | <u>Rhinonyssidae</u> |
| <i>E. trouessarti</i> (Berlese,1892)
= <i>G. thalassinus</i> Berlese & Trouessart,1889
= <i>G. fucorum</i> var. β Berlese & Trouessart,1889
Norway, Northumberland, Ireland, France | <i>Rhinonyssus</i> Trouessart,1884 |
| <i>Parasitus</i> Latreille,1795 | <i>R. sp.</i>
@ <i>Uria aalge</i>
British Columbia |
| <i>P. kembensi</i> Oudemans,19??
* In decaying wrack.
Br. Isles | <u>Rhodacaroidea</u> |
| <i>P. flevensis</i> Oudemans,1922
* In wrack.
Zuidersee coast | <i>Rhodacaridae</i> Oudemans,1902 |
| <i>Cornigamasus</i> Evans & Till,19?? | <i>Rhodacarinae</i> Oudemans,1902 (sensu Lee,1970) |
| <i>C. sp.</i>
Britain | <i>Rhodacarus</i> Oudemans,1902 |
| <i>Gamasodes</i> Oudemans,1939 | <i>R. strenzkei</i> Willmann,1957
* intertidal
Kiel |
| <i>G. sp.</i>
Britain | <i>R. roseus</i> Oudemans,1902
@ "orange lichen zone"
Ireland |
| <i>Holoparasitus</i> Oudemans,1936 | <i>R. pallidus</i> Hull,1918
* In intertidal crevices.
Kiel, Northumberland, Scotland, W Ireland, Spain |
| <i>H. sp.</i>
Britain | <i>R. corniculatus</i> Willmann,1935 |
| <i>Paragamasus</i> Hull,19?? | <i>R. silesiacus</i> Willmann,1936
@ intertidal sand
?Europe? |
| <i>P. sp.</i> | <i>R. rhodacaropsis</i> Willmann,1935 |
| | <i>R. inexpectata</i> Willmann,1935
@ coastal ground water
Kiel |
| | <u>Cyrtolaelaniidae</u> |
| | <i>Hydrogamasus</i> Berlese 1892 |

<i>H. salinus</i> (Laboulbène,1851?) =? <i>H. littoralis</i> (G. & R. Canestrini,1885) * On <i>Ascophyllum</i> and in rock-crevices of high water zone; feeds on Collembola. W Ireland	Britain
<i>H. giardi</i> (Berlese & Trouessart,1889) = <i>Sinus giardi</i> Berlese & Trouessart,1889 W. Ireland, ?French Atlantic coast	
<u>Eviphidoidea</u>	
<u>Eviphididae</u>	
<i>Thinoseius</i> Halbert,1920 * In decaying seaweed.	
<i>T. ramsayi</i> Evans,19?? Europe	
<i>T. fucicola</i> (Halbert,1920) =? <i>T. berlesii</i> Halbert,1920 W Ireland, SW England	
<i>T. spinosus</i> Willmann,19?? Europe	
Pachylaelapidae Vitzthum,1931	
<i>Pachylaelaps</i> Berlese,1888	
<i>P. littoralis</i> Halbert,1915 W Ireland	
<i>Onchodellus</i> Berlese,1904	
<i>O.</i> sp. Britain	
Machrochelidae Vitzthum,1930	
<i>Machrochelus</i> Latreille, in Cuvier,1829 * <i>Macrochelus</i> is the only spelling in Neave	
<i>M. marginatus</i> var. <i>littoralis</i> (Halbert,1915) = <i>Holostaspis marginatus</i> var. <i>littoralis</i> Halbert,1915 Ireland	
<u>Ascoidea</u>	
Halolaelapidae	
<i>Gamasellus</i> Berlese,1892 (TM <i>G. falciger</i> (G. & R. Canestrini,1885??)) * This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?	
<i>G. inermis</i> Halbert,1920 W Ireland	
<i>Gamasus</i> Latreille,1802 * This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?	
<i>G. (Gamasus) kempersi</i> Oudemans,1902 Norway, Netherlands, Northumberland, Ireland	
<i>G. (Gamasus) lunaris</i> Berlese,1906 = <i>G. rubescens</i> G. & R. Canestrini,1882 * Possibly not a real shore-mite Northumberland, Ireland, Italy	
<i>G. coleoptratorum</i> (Linnaeus,1758) = <i>Acarus coleoptratorum</i> Linnaeus,1758 Ireland	
<i>G. crassipes</i> (Linnaeus,1758) var. <i>longicornis</i> Berlese,1??? * Possibly not a real shore-mite W Ireland	
<i>Gamasoides</i> Berlese,1903 (TM <i>Poecilochirus carabi</i> G. & R. Canestrini,1885)	
<i>G. spinipes</i> (C.L. Koch,1844) = <i>G. brachiosus</i> (G. Canestrini,1883-89) W Ireland	
<i>Halolaelaps</i> Berlese & Trouessart,1889	
<i>H. marinus</i> (Brady,1875) = <i>Gamasus marinus</i> Brady,1875 =? <i>H. glabrioculus</i> Berlese & Trouessart,1889 = <i>Zercon marinus</i> : Monez,1890 = <i>Parastinus marinus</i> : Oudemans,1902 W Ireland, Britain, France, Spain	
<i>H. celticus</i> Halbert,1915 W Ireland	
<i>H. incisus</i> Hyatt,1980? Br. Isles	
<i>Sapromasellus</i> Berlese & Trouessart,1???	
<i>S.</i> sp. Britain	
<u>Antennoseiidae</u>	
<i>Antennoseius</i> Berlese,1916	
<i>A.</i> sp. Britain	
<u>Ascidae</u>	
<i>Episeius</i> Hull,1918 * This genus is not found in Hayward & Ryland (Eds.) 1990. Synonymized? - Correct family?	
<i>E. grandis</i> (Berlese,1916) = <i>Lasioseius grandis</i> Berlese,1916 W Ireland	
<i>Arctoseioides</i> Willmann,1949	
<i>A.</i> sp.	
	Britain
	<i>Arctoseius</i> Thor,1930
	<i>A.</i> sp. Britain
	<i>Iphidozercon</i> Berlese,1904
	<i>I.</i> sp. Britain
	<i>Leioseius</i> Berlese,1917
	<i>L.</i> sp. Britain
	<i>Neajordensis</i> Evans,1957
	<i>N.</i> sp. Britain
	<i>Platyseius</i> Berlese,1917
	<i>P.</i> sp. Britain
	<u>Phytoseiidae</u>
	<i>Amblyseius</i> Berlese,1915
	<i>A.</i> sp. Britain
	<i>Lasioseius</i> Berlese,1917 (TM <i>Seius muricatus</i> (C.L. Koch,18??))
	<i>L. salinus</i> Halbert,1920 W Ireland
	<i>Seius</i> Agassiz,1845 = <i>Sejus</i> Koch,1836
	<i>Tychlodromus</i> Scheuten,1???
	<i>T.</i> sp. Britain
	<u>Veigaioidae</u>
	<i>Veigaiidae</i> Oudemans,1939
	<i>Cyrthydrolaelaps</i> Berlese,1905
	<i>C. hirtus</i> Berlese,1905 * In intertidal crevices Wales, W Ireland, NW France, Spain
	<i>Gamasolaelaps</i> Berlese,1904
	<i>G. excisus</i> (L. Koch,1879) = <i>Seius excisus</i> L. Koch,1879 = <i>Cyrtolaelaps (?) aurantiacus</i> Berlese,1903 = <i>Metaparacitus suboles</i> Oudemans & Voigts,1904 Bremen, W Ireland, Siberia
	<u>IXODIDA</u>
	<u>Ixodoidea</u>
	<i>Ixodidae</i> = <i>Isoodidae</i>
	<i>Ixodes</i> Latreille,1795
	<i>I. uriae</i> (White,1???) @ <i>Uria aalge</i> British Columbia
	<u>NOTOSTIGMATIDA</u>
	* Not in Scandinavian maritime or marine zone?
	<u>HOLOTHYRIDAE</u>
	* Not in Scandinavian maritime or marine zone?
	<u>PANTOPODA</u> Gerstaecker,1863
	= <u>PYCNOGONIDA</u> Latreille,1810
	= <u>PODOSOMATA</u> Leach,1815
	<i>PEGMATA</i> Fry,1978
	<i>Nymphonidae</i> Wilson,1878
	<i>Nymphon</i> J.C. Fabricius,1794
	<i>N. brevirostre</i> Hodge,1863 = <i>N. rubrum</i> Hodge,1865 = <i>N. gracile</i> : G.O. Sars Öresund, Isefjord, Bohuslän, Denmark, all Norway, British Isles, Bretagne
	<i>N. brevitarse</i> Kroyer,1844 Greenland, Strait of Matotschkin Sharr
	<i>N. hirtum</i> J.C. Fabricius,1794 = <i>Chaetonyphon hirtum</i> : Auctt. Skagerak, Troms, Belgium, Northumberland, Scotland, Shetlands, Iceland, E & W Greenland, Spitzbergen
	<i>N. glaciale</i> Lilljeborg,1870 Varangerfjord
	<i>N. gracile</i> Leach,1814 non = <i>N. gracile</i> : G.O. Sars Bergen - Nordland, Morocco
	<i>N. grossipes</i> (O. Fabricius,1780) Lofoten - Varangerfjord
	<i>N. mixtum</i> Kroyer,1844-45 = <i>grossipes</i> Auctt. non? (O. Fabricius,1780) * <i>N. grossipes</i> , <i>N. mixtum</i> & <i>N. glacialis</i> Lilljeborg,1851 seem to be a species complex, of which only <i>N. mixtum</i> can be found S. of the Arctic. Öresund - Bohuslän - Lofoten, Denmark, Shetland

- N. leptochelus* G.O. Sars,1888
Bohuslän-Skagerrak, all Norway, British Isles
- N. gracilipes* Heller,1875
= *N. giganteum* Goodsir,1844
* Is very close to, or possibly identical with *N. stroemi*.
Skagerrak (off Arendal) in 150-250 m - NE Finnmark, Northumberland
- N. stroemi* Kroyer,1844-45
Kattegatt - Bohuslän - W Finnmark
- N. macrum* Wilson,1880
W. Skagerrak, Bergen - NE Finnmark, W Greenland
- N. longitarse* Kroyer,1844-45
Bergen, Varangerfjord, Northumberland, Arctic
- N. longimanum* (G.O. Sars,1888)
W. Skagerrak??, Jan Mayen, Kara Sea
- N. spinosum* Goodsir,1842
= *Chatonymphon s.*
Bohuslän-Skagerrak, all Norway
- N. spinosissimum* Norman,1894
= *Chatonymphon s.*
Bergen - Varangerfjord, Spitsbergen, Iceland, S & W Greenland
- N. hirtipes* Bell,1853
= *Chatonymphon h.*
Lofoten - NE Finnmark, NE of Shetland
- N. macronyx* G.O. Sars,1877
= *Chatonymphon m.*
NE Finnmark, Faroe-Shetland Channel, Spitzbergen, E & W Greenland
- N. tenellum* (G.O. Sars,1888)
= *Chatonymphon tenellum* G.O. Sars,1888
W. of Finnmark at 1134m to Faroe Channel
- N. serratum* G.O. Sars,1879
W. Norway, Hudson Bay
- N. megalops* G.O. Sars,1887
Bergen - S Troms, between Norway and Shetland
- N. elegans* H.J. Hansen,1887
N. Norway, Shetland-Faroe Channel
- N. sluteri* Hoek,1881
Varangerfjord - NE Finnmark, Spitzbergen, Arctic
- N. microrhynchum* G.O. Sars,1888
Novaja Zemlya, Arctic
- Boreonymphon* G.O. Sars,1888
- B. abyssorum* (Norman,1873)
= *Nymphon abyssorum* Norman,1873
More, Norwegian Sea
- B. ossiansarsi* Knaben, in Just,1972
Off W. Norway
- B. compactum* Just,1972
E & NW Greenland
- B. robustum* (Bell,1855)
= *Nymphon robustum* Bell,1855 (TM)
= *Nymphon hians* Heller,1875
between Faeroes & Iceland, off N Iceland, E & W Greenland, Spitsbergen, Kara Sea
- Acheliidae Fry,1978
- Achelia* Hodge,1864
- A. echinata* Hodge,1864
@ *Fistula folacea*
Bohuslän, Hirshals, Helgoland, Bergen - E Finnmark, British Isles
- A. longipes* (Hodge,1864)
= *Ammothea longipes* Hodge,1864
NW Finnmark - Porsangerfjord, British Isles, Atlantic coast of France, W Ireland
- A. hispida* Hodge,1864
* May be synonymous with *A. longipes*.
W Finnmark - Porsangerfjord, SW Britain, W Ireland
- A. laevis* Hodge,1864
Trondelag S of Trondheimsfjord, Helgoland, British Isles
- A. simplex* (Giltay,1934)
Isle of Man, Galway, Millport
- A. vulgaris* (Costa,1861)
Britain
- Ammotheidae Dohrn,1881
= *Phoxichilididae* G.O. Sars,1891 (p.p.)
- Ammothea* Leach,1814
- A. hilgendorfi* (Böhm,1879)
- Eurycyde Schiödte, in Rink,1857
- E. hispida* (Kroyer,1844-45)
Norway (Nordmøre and northwards), Denmark?
- Phoxichilidium* Milne-Edwards,1840
- P. femoratum* (Rathke,1799)
@ reported from cysts in *Syncoyne eximia*
Bohuslän, all Norway, Britain, Greenland, France
- P. tubulariae* Lebour,1947
@ gastric cavity of *Tubularia larynx*
Bohuslän, Holland, British Isles
- P. viresceus* Hodge,1864
- @ reported from cysts in *Syncoyne eximia*
British Isles, France
- Paranymphon* Caulery,1896
* Correct family?
- P. spinosum* Caulery,1896
off Wales in >90m
- Tanystylum* Miers,1879
* Correct family?
- T. conirostre* (Dohrn,1881)
? Britain, Mediterranean
- Anoplodactylidae
- Anoplodactylus* Wilson,1878
- A. angulatus* (Dohrn,1881)
SW Britain, France, Mediterranean
- A. petiolatus* (Kroyer,1844)
@ manubrium of *Obelia* sp. medusae
Öresund - Bohuslän - S Troms, North Sea, British Isles
- A. pygmaeus* (Hodge,1864)
@ gastric cavity of *Obelia geniculata* polyps & on *Dynamena pumila*
Denmark, British Isles, Helgoland, Holland, France, Azores
- A. exiguis* (Dohrn,1881)
Bohuslän? (refound recently (Oct. 1994) in 30-38 m. at Lilleskärsslättan), Atlantic coast of France
- A. typhlops* G.O. Sars, 1891?
Norway (More - Nordland), Off W. Ireland
- Endeidae Norman,1908
- Endeis* Philippi,1843
- E. spinosa* (Montagu,1808)
= *Phoxichilus s.*
= *E. laevis* (Grube,1871)
Bohuslän, Stavanger - More, British Isles
- E. charybdaea* (Dohrn,1881)
= *E. spinosa* sensu King, 1974, non (Montagu)
Stavanger area (Dusevigen), Roscoff, British Isles, Mediterranean
- Collosendeidae Hoek,1881
- Collossendeis* Jarzynsky,1870
- C. proboscidea* (Sabine,1824)
Varangerfjord, Scotland-Faroe Channel
- C. angusta* G.O. Sars,1877
Between Norway and Shetland at 670-820m
- Callipallenidae Hilton,1942
- Callipallene* Flynn,1929
- C. brevirostris* (Johnston,1837)
Kieler Bucht - Bohuslän - S Nordland, Hirshals, British Isles
- C. phantoma* (Dohrn,1881)
= *Pallene producta* G.O. Sars,1888
Bergen - NW Sor-Trondelag, British Isles
- C. spectrum* (Dohrn,1881)
Britain-Mediterranean
- C. tiberi* (Dohrn,1881)
Britain-Mediterranean
- C. emaciata* (Dohrn,1881)
Atlantic coast of France, England, Wales
- Cordylochete* G.O. Sars,1888
- C. longicollis* G.O. Sars,1888
Skagerrak at 343m (off Arendal) - Lofoten
- C. malleolata* (G.O. Sars,1879)
Off N. Norway at 168-200m
- C. brevicollis* G.O. Sars,1888
Vadsø, Faeroes
- Pseudopallene* Wilson,1878
- P. circularis* (Goodsir,1842)
= *Pallene intermedia* Kroyer,1844-45
= *Pallene discoidea* Kroyer,1844-45
Kullen - Skagerrak - all Norway, Scotland
- P. spinipes* (O. Fabricius,1780)
Bergen - NE Finnmark
- PYNOGONOMORPHIDA Fry,1978
- Pycnogonidae* Wilson,1878
- Pycnogonum* Brönnich,1764
- P. littorale* (Ström,1762)
@ *Actinia equina*, *Metridium senile* & *Urticina felina*; may also be associated with *Cucumaria frondosa*
Öresund - Bohuslän - all Norway, British Isles
- P. crassirostre* G.O. Sars,1888
@ Sea anemones
North Sea, Hardangerfjord - Trondheimsfjord - S Troms
- ASCORHYNCHOMORPHIDA Fry,1978
- * Not represented in the area.
- RHOPALORHYNCHOMORPHIDA Fry,1978
- * Not represented in the area.
- RHYNCHOTHORACIMORPIDA Fry,1978

* Not represented in the area.

UNIRAMIA Snodgrass,1935

* Most taxa of this group treated in this list, are not truly marine. Some Diptera larvae are obligate or facultative estuarine or marine inhabitants. The rest of the species represented in the check-list may be referred to as thalassobionts (bound to the marine littoral fringe) or thalassophilic (facultative inhabitants, of the sea shore, preferring this environment to others). Many other organisms may be characterized as either indifferent sea shore taxa (may as well be found in other environments) or as temporal taxa in the marine littoral. Organisms of indifferent and temporal occurrence in the marine supralitoral zone have, in most cases, been completely left out from this list. Some insect orders, e.g. Hymenoptera, to which some sea shore dwellers belong (e.g. the sand dune species *Pomphius* Fabricius,1798 *phloebeus* (Fabricius,1787)), have been omitted in this list, on reasons of difficulties in bordering between maritime and non-maritime species. Thus, this tableau must be regarded as very incomplete as a check-list. Certainly, several taxa are missing, and some of the encountered species may possibly show to be temporal visitors to the marine littoral, and should thus perhaps be weeded out.

MYRIPODA "Tusenfotinger"

CHILOPODA "Enkelfotinger"

* Only thalassobionts and a few thalassophilic taxa included

ANAMORPHA

LITHOBIIIDA "Stenkrypare"

LITHOBIIDEA

Lithobioidae

Lithobius Leach,1814

L. melanops G. Newport,1845

* Thalassophilic.

Scandinavia, Germany, Iceland

L. forficatus (Linnaeus,1758)

* Thalassophilic.

Scandinavia (e.g. Bohuslän-Uppland), Iceland, France

L. calcaratus ?Koch,1???

* In wrack.

Småland archipelago

L. erythrocephalus ?Koch,1???

* In wrack.

Blekinge-Uppland, Åland

L. curtipes ?Koch,1???

* In wrack.

Uppland, Åland, Finland

Henicopidae

Lamycetes Meinert,1868

* Does this genus belong in that family?

L. fulvicornis Meinert,1868?

* Thalassophilic.

Iceland

EPIMORHA

GEOPHILIDA "Jordkrypare"

Schendylidae

Schendyla Bergsöe & Meinert,1866

S. nemorensis (C.L. Koch,1836)

* Thalassophilic. In Wrack.

Bohuslän

Hydroschendyla Brölemann & Ribaut,1911

H. submarina (Grube,1870)

Sweden, Bretagne, S. England, Mediterranean

Brachyschendyla Brölemann & Ribaut,1911

B. monodi Brölemann,1924

Finistère, Arcachon

Geophilidae

Strigamia Gray,1843

= *Scoloplanes* Bergsöe & Meinert,1866

S. maritima (Leach,1817)

Gotland, Vinga, Bohuslän, S. Norway, Denmark, Helgoland, Br. Isles, France

Geophilus Leach,1814

G. fucorum ?1??? *seurati* Brölemann,1930?

Plymouth, Scotland, Ireland

G. proximus Koch,1???

* In wrack.

Uppland, Åland, Finland

G. algarum ?1??? *decipiens* Brölemann,19??

N France, Arcachon

Necrophloeophagus Newport,1843

N. longicornis (Leach,1815)

= *Geophilus longicornis* Leach,1815

* Thalassophilic.

Scandinavia, Britain

Chilopodidae

Pachymeriinae

Pachymerium C.L. Koch,1847

P. ferrugineum (C.L. Koch,1835)

* Thalassophilic.

Blekinge, Uppland, Denmark, Åland, Åbo, Germany, Arcachon

SYMPHYLA "Dvärgfotinger"

Scolopendrellidae

Scolopendrella Gervais,1839

S. notacantha Gervais,1839

* Thalassophilic.

Wales

PAUROPODA "Fäfotinger"

TETRAMEROERCERATA

ECTOMORPHA

Pauropodidae

Thalassopauropus Bagnall,1935

T. remyi Bagnall,1935

E. Scotland, French Mediterranean

Altopauropus Berlese,1902, ex Silvestri MS

A. thalassophilus Remy,19??

E. Scotland

A. danicus (Hansen,1???)

E. Scotland

A. littoralis Bagnall,1935?

E. Scotland

A. stephani Bagnall,1935?

E. Scotland

DIPLOPODA "Dubbelfotinger"

* Only thalassobionts and a few thalassophilic taxa included

PENICILLATA

= PSEPHALOGNATHA Latzel,1884 "Penselfotinger"

POLYXENIDA

= SCHIZOCEPHALA Verhoeff,1926

Polyxenoidea

Polyxenidae Gray & Jones,1842

Polyxenus Latreille,1802

P. lagurus (Linnaeus,1758)

* Thalassophilic.

Bohuslän, Skåne, Åland, Germany, Denmark, Norway, Br. Isles, etc.

PENTAZONIA

* No maritime species

HELMINTHOMORPHA

DIPLOCHETA

JULIDA

Nemasomatidae

Thalassoisobates Verhoeff,1908

= *Isobates* Menge,1801 (p.p.)

T. littoralis (Silvestri,1903)

* In wrack on rocky shores.

Bohuslän, Br. Isles, Mediterranean

Blaniulidae Attems,1909

Choneiulus Brolemann,1921

C. palmatus (Nemec,1895)

* Occasionally in wrack.

Bohuslän, S Norway, Denmark, Finland, Germany, Netherlands, Britain

Proteroiulus Silvestri,1897

P. fuscus (Am Stein,1857)

* Occasionally in wrack, normally a forest species.

S Sweden, Norway, Åland, S Finland, Germany, Netherlands, Belgium, France, Britain

Julidae Leach,1815

Cylindroiuliinae Verhoeff,1930

Cylindroiulus Verhoeff,1894

C. (Aneuloboiulus) latestratus (Curtis,1844)

= *C. (Aneuloboiulus) frisius* (Verhoeff,1891)

* Thalassophilic. In wrack.

Gotland, Bohuslän, Norway, Denmark, Germany, Faroes, Netherlands, Br. Isles, Ireland, Arcachon

C. (Aneuloboiulus) punctatus (Leach,1815)

= *C. (Aneuloboiulus) silvarum* (Meinert,1868)

* Occasionally in wrack.

Småland archipelago, Uppland, W Sweden, Norway, Denmark, Finland, Britain, Netherlands, Belgium, France

Julinae Verhoeff,1930

Julus Linnaeus,1758

J. terrestris Porat,1889

* Thalassophilic

Finland, Åland, Gotland, E. & S. Sweden, Bornholm, Sealand, Bohuslän, Germany

Schizophyllinae Verhoeff,1910

Omnatoiulus Latzel,1884

= *Schizophyllum* Verhoeff,1895 (p.p.)

= *Archithus* Berlese,1886, non Scudder,1873 (p.p.)

- O. sabulosus** (Linnaeus,1758)
 = *Schizophyllum* Verhoeff,1895 (*Bothriulus* Verhoeff,1894) *sabulosum* (Linnaeus,1758)
 * Thalassophilic
 S. Finland, Denmark, Germany, Sweden, S Norway, Br. Isles etc.
- Tachypodoiulus** Verhoeff,1894
- T. niger** (Leach,1815)
 = *Julus albipes* C.L. Koch,1838
 * Thalassophilic.
 Normandie, Netherlands, Belgium, Germany, Br. Isles, Ireland
- Brachytulina** Verhoeff,1910
- Brachiyulus** Berlese,1884
- B. pusillus** (Leach,1814)
 = *B. littoralis* Verhoeff,1898
 Öland, Gotland, Skåne-Blekinge-Småland, Estonia-Germany, Britain, Ireland, France, Spain
- HEXAPODA** Latreille,1825
 * Probably a polyphyletic taxon. Some entognaths may be closer to Myriapoda.
- ENTOGNATHA**
- DIPLURATA** "Larvborstsvansar"
 * Not represented in the area with any sea shore species.
- MYRIENTOMATA**
 = **PROTURA** "Trevfotinger"
 * Not represented in the area with any sea shore species.
- OLIGOENTOMATA**
- COLLEMBOLA** Lubbock,1870 "Hoppstjärtar"
 * Chiefly thalassobionts and a few thalassophilic species have been listed here.
- PODUROMORPHA** Börner,1913
- Hypogastruridae** Börner,1906
- Hypogastrura** Bourlet,1839
- H. viatica** (Tullberg,1872)
 * Thalassophilic. Considered to be cosmopolitan in wrack and salt marshes.
 Bohuslän-all Baltic coasts, Denmark, Norway, German North Sea coast, Br. Isles, France
- H. vernalis** (Carl,1901)
 * Littoral & in wrack.
 Sörmaland-Uppland, Åland
- H. gisini** (Strenzke,1954)
 along sandy sites in Baltic Shores & Sylt
- Ceratophysella** Börner, in Brohmer,1942
- C. armata** (Nicole,1842)
 = *Hypogastrura armata* : Auctt.
 * Ubiquistic & in wrack.
 Bohuslän-Uppland, Finland
- Xenylla** Tullberg,1871
- X. humicola** (O. Fabricius,1780)
 * Thalassophilic. Considered to be presumably cosmopolitan in wrack and salt marshes.
 Bohuslän-all Baltic coasts, Norway, S. North Sea coasts, Br. Isles, France, Mediterranean
- X. maritima** Tullberg,1871
 * Thalassophilic. Considered to be presumably cosmopolitan in wrack and salt marshes.
 Finland, W. Sweden, Uppland, Norway, German North Sea coast, Br. Isles, France, Mediterranean
- Neanuridae** Börner,1901
- Neanurinae** Börner,1901
- Neanura** Macgillivray,1893
 = *Anoura* Gervais,1843, *non* Gray,1838, (Mammalia) *nec* Gray,1840 (Mollusca)
- Pseudachorutinae** Börner,1906
- Anurida** Laboulbène,1865
- A. maritima** (Guérin-Méneville,1836) Laboulbène,1865
 Kattegatt-Bohuslän, Norway, Germany, Netherlands, Belgium, Br. Isles, Arcachon
- A. tulbergi** Schott,1891
 * thalassophilic
 W. Sweden, Sörmaland, Finland, German Baltic & North Sea coasts, Norway, Britain, France, Mediterranean
- A. granaria** (Nicole,1847) Tullberg,1869
 * In wrack & humus
 Skåne-Småland, Finland
- A. thalassophila** (Bagnall,1939) Stach,1949
 = *Anuridella thalassophila* : Auctt.
 Br. Isles
- Pseudachorutella** Stach,1949
- P. remyi** (Denis,1933) Massoud,1967
 = *Anurida remyi* : Auctt.
 Finland, Skåne, German Baltic & North Sea coasts, Jan Mayen, Greenland
- Gastranuridae** Bagnall,1949
- G. denisi** (Bagnall,1939) Massoud,1967
 = *Anurida denisi* : Auctt.
 Br. Isles, France
- Anuridella** Willem,1906
- A. marina** Willem,1906
 Br. Isles, France
- A. calcarata** Denis,1925
 Br. Isles, Mediterranean
- A. germanica** (Gisin,1951)
 German Baltic & North Sea coast
- A. submarina** Bagnall,1934
 * A synonym of which other species?
 Br. Isles
- A. immsiana** Bagnall,1939
 * A synonym of which other species?
 Br. Isles
- Frieseinae** Massoud,1967
- Friesea** Dalla Torre,1895
- F. acuminata** (Denis,1925) Goto & Lawrence, in Kloet & Hincks,1964
 Br. Isles, Mediterranean
- F. mirabilis** (Tullberg,1871) Willem,1901
 * In deep wrack layers & humus
 Vinga-Uppland, Åland, Finland
- Brachystomellidae** Stach,1949
- Brachystomella** Ågren,1903
- Onychiuridae** Börner,1901
- Onychiurus** Gervais,1841
- Protaphorura** Absolon,1901
- P. debilis** (Moniez,1889) Salmon,1964
 = *Onychiurus debilis* : Auctt.
 German Baltic & North Sea coast, Br. Isles, Arcachon, Mediterranean
- P. thalassophila** (Bagnall,1937) Hopkin,1997
 = *Onychiurus thalassophila* : Auctt.
 Scotland, England
- P. halophila** (Bagnall,1937) Salmon,1959
 = *Onychiurus halophila* : Auctt.
 Northumberland, Yorkshire
- P. armata** (Tullberg,1869) Börner,1909
 = *Onychiurus armata* : Auctt.
 * In deep wrack layers.
 Bohuslän-Uppland, Finland
- P. tullbergi** (Bagnall,1935) Kloet & Hincks,1945
 = *Onychiurus tullbergi* : Auctt.
 Scotland
- Tullbergiidae** Bagnall,1935
- Tullbergia** Lubbock,1876
- Odontellidae** Massoud,1967
- ENTOMOBRYOMORPHA** Börner,1913, sensu Mateos,1988
- Ispotomoidea** Schäffer,1896 (Börner,1913 (Szeptycki,1979))
- Isotomidae** Schäffer,1896 (Börner,1913)
- Isotominae** Schäffer,1896
- Bagnallella** Salmon,1951
- Isotoma** Bourlet,1839, *non* Dejean,1834 (Coleoptera)
- Desoria** Nicolet, in Desor,1841
- D. tigrina** Nicolet,1842
 * In deep wrack layers.
 Bohuslän-Uppland, Åland, All Europe, Arctic, Antarctic, Australia
- D. olivacea** (Tullberg,1871)
- * In deep wrack layers & eurytopic
 Bohuslän-Uppland, Åland, All Europe, Arctic
- Halisotoma** Bagnall,1949
- H. maritima** (Tullberg,1871) Bagnall,1949
 = *Isotoma maritima* Tullberg,1871
 * In wrack and under stones.
 W. Sweden, Skåne, all Baltic coasts, Norway, S. North Sea coasts, Br. Isles, France, Mediterranean
- H. boneti** (Delamare-Debouteville,1954)
 Portugal-Senegal, S Spain, Mediterranean
- H. poseidonis** (Bagnall,1939)
 = *Isotoma poseidonis* Bagnall,1939
 Br. Isles, Ireland, Biscay
- Isotomiella** Bagnall,1939
- Isotomurus** Börner,1903
- Axelsonia** Börner,1906
- A. littoralis** (Moniez,1890) Handschin,1929
 * In wrack and littoral subsoil.
 North Sea shores, Br. Isles, French North Sea coast, Arcachon, Mediterranean, S America, Australia
- Proisotominae**
- Folsomia** Willem,1902
- F. sexoculata** (Tullberg,1871) Handschin,1929
 Skåne-Uppland, Åland, Norway, German Baltic & North Sea coast, Finland, Denmark, Br. Isles
- F. thalassophila** Bagnall,1940
 Scotland, Ireland
- F. garetti** Bagnall,19??
 * Possibly not a sea shore species? Not recognized in Janssens' check-list.
 Br. Isles
- Proisotoma** Börner,1901
- P. buddenbrocki** Strenzke,1954
 Tvärminne, Kieler Bucht

- P. admiritalia* Murphy,1953
Scotland
- Ballistura* Börner,1906
- B. schoetti* (von Dalla Torre,1895) Stach,1947
* Thalassophilic. In wrack and under stones.
W. Sweden, Sörmland-Uppland, Finland, German North Sea coast, Br. Isles
- B. crassicauda* Tullberg,1871
* Thallasophilic. In wrack and under stones. Not recognized in Janssens' check-list.
Finland, German Baltic coast, W. Sweden, Br. Isles
- Anurophorinae
- Archisotoma* Axelson,1912
* in littoral subsoil and salt marshes
- A. besseli* (Packard,1877) Willem,1925
Sweden, ?Norway, German Baltic & North Sea coast, Finland, ?Br. Isles, France, Mediterranean
- A. pulchella* (Moniez,1890) da Gama,1968
Denmark, German North Sea coast, Arcachon, ?Br. Isles
- A. megalops* Bagnall,1939
Br. Isles
- A. nigricans* Bagnall,1939
Br. Isles
- A. subbrucei* Delamare-Debouteville,1954
French Atlantic coast
- Actaletidae Börner,1902
- Actaletes* Giard,1889
- A. neptuni* Giard,1889
On rocks in the intertidal zone of the French North Sea coast, ?Br. Isles
- Coenaeletidae Bellinger,1985
- Isotogastruridae* Thibaud & Najt,1992
- Entomobryoidae Schött,1891 (Womersley,1934, sensu Szepetki,1979)
- Entomobryidae Schott,1891
- Entomobryinae* Schott,1891 (Schäffer,1896)
- Entomobrya* Rondani,1861
= *Degerria* Nicolet,1841, non Meigen,1838 (Diptera)
= *Mydonius* Gistl,1848
- Seira* Lubbock,1870
- Willowsia* Shoebotham,1917
- Lepidocyrinidae
- Lepidocyrtus* Bourlet,1839
- L. cyaneus* (Tullberg,1871) Tullberg,1871
* In wrack surface layers.
Bohuslän-Uppland, Finland
- L. tanuginosus* (Gmelin,1788) Tullberg,1871
* In wrack surface layers & ubiquitous.
Blekinge-Uppland, Åland, Finland
- Pseudosinella* Schäffer,1897
- P. halophila* Bagnall,1939
Sylt, England
- P. maritima* Bagnall,1941
Br. Isles, Ireland
- Orchesellinae
- Orchesella* Templeton,1835
- O. litoralis* Brown,1925
Yorkshire
- Microfalciliidae Massoud & Betsch,1966
- Paronellidae Börner,1913
- Cyphoderidae Börner,1913
- Oncopoduridae Carl & Lebedinsky,1905
- Tomoceroidea Schäffer,1896 (Szepetki,1979)
- Tomoceridae Schäffer,1896
- METAXYPLEONA Salmon,1964
- Poduridae Latreille,1802-03
- SYMPHYPLEONA Börner,1901
- Sminthuridae Lubbock,1862
- Deuterostominturus* Börner,1901
= *Deuterostominturus* Börner,1901
- Sminthurides* Börner,1900
- Sminthurinus* Börner,1901
- Smynthurus* Latreille,1802
- Bourletellidae Börner,1913
- Bourletiella* Banks,1899
- Mackenziellidae Yosii,1961
- Sminthuriidae Börner,1906
- Kataniidae Börner,1913, sensu Stach,1956
- Sturmidae Bretfeld,1994
- Arthropalpitidae Stach,1956
- Spinothecidae Delamare Deboutteville,1961
- Dicrtonidae Börner,1906
- NEELIPLEONA Massoud,1971
- Neelidae Folsom,1896
- INSECTA
= ECTOGNATHA
- ARCHEOGNATHA
- APTERYGOTA
= ZYGENTOMA
- THYSANURA Latreille,1804 "Fjällborstsvensar"
- ZYGOENTOMATA
* Not represented in the marine littoral in the area.
- ARCHEOGNATHA
= MICROCORYPHIA
- Machiloidea
- Machilidae Verhoeff,1912 "Nattsmýgar", "Silverfiskar"
- Machilinae
- Petrobius* Leach,1817 "Kustnattsmýgar"
- P. brevistylis* Carpenter,1913
= *P. maritimus* Auct., non (Leach,1815)
Baltic-Bohuslän, Helgoland
- P. maritimus* (Leach,1809)
= *P. lohmanderi* Agrell,1944 (synonymized by Davies 1988)
E. Sweden (on rocky shores), Denmark, Atlantic coasts (on rocky shores), e.g. Br. Isles
- PTERYGOTA
= PTILOTA "Yrfän"
- PAL(A)EOPTER(YGOT)A
* The first order of this infraclassis, the Ephemeroptera, has no really marine or maritime species in the area. However larvae of some species within Odonata may be found in estuaries. A list of such species found in the Baltic by Lindberg,1948 thus has been used for making this list.
- EPHEMEROPTERA "Dagsländer"
- Ephemeroidae
- Ephemeridae
- Baetidae
= Heptagenioidea
- Baetidae
- Callibaetis* Eaton,1881
- ODONATA "Trollslandor"
- ZYGOPTERA
- Lestinoidea
- Lestidae
- Lestes* Leach,1815
- L. sponsa* (Hansemann,1823)
N. Baltic
- Coenagrionoidea
- Coenagrionidae
- Ischnura* de Charpentier,1840
- I. elegans* (P.L. v. d. Linden,1820)
N. Baltic
- Enallagma* de Charpentier,1840
- E. cyathigerum* (de Charpentier,1840)
N. Baltic
- Erythromma* de Charpentier,1840
- E. najas* (Hansemann,1823)
N. Baltic
- Arionoidea
= Calopterygoidea
- Ariionidae
= Calopterygidae
- Agrion* Fabricius,1775
= *Calopteryx* de Charpentier, in Burmeister,1839
- A. armatum* de Charpentier,1840
N. Baltic
- A. pulchellum* P.L. v. d. Linden,1825
N. Baltic
- A. hastulatum* de Charpentier,1825
N. Baltic
- ANISOPTERA

AeshnoideaAeshnidae

Aeshna Fabricius,1775
= *Aeschna* Illiger,1802

A. coerulea (H. Strom,1783)
N. Baltic

A. juncea (Linnaeus,1758)
N. Baltic

A. grandis (Linnaeus,1758)
N. Baltic

A. cyanea (O.F. Müller,1764)
N. Baltic

Brachytron Evans,1845

B. hafniense (O.F. Müller,1764)
N. Baltic

LibelluloideaLibellulidae

Libellula Linnaeus,1758

L. quadrimaculata Linnaeus,1758
N. Baltic

Sympetrum Newman,1833

S. striolatum (de Charpentier,1840)
N. Baltic

S. vulgatum (Linnaeus,1758)
N. Baltic

S. flaveolum (Linnaeus,1758)
N. Baltic

S. danae (J.H. Sulzer,1776)
N. Baltic

Leucorrhinia Brittinger,1850

L. rubicunda (Linnaeus,1758)
N. Baltic

Orthetrum Newman,1833

O. cancellatum (Linnaeus,1758)
N. Baltic, SW Sjælland in ≈10 ppt S

NEOPTERAORTHOPTEROIDEA

* As far as known, by the compiler there are no truly maritime species.

ORTHOPTERA

= SALTATORIA "Hopprätvingar"

CAELIFERA

Acrididae "Markgräshoppor"

Onocestus Bolívar,1878

O. haemorrhoidalis (?17???)
Germany

ENSIFERA

= GRYLLOPTERA

Tettigoniidae "Vårtbitare"

Decticus Audinet-Serville,1831

D. verrucivorus (Linnaeus,1758) "Stor vårtbitare"
= *Grylhus verrucivorus* Linnaeus,1758
Sweden, Germany

DERMAPTERAFORFICULINALabiduroideaCarcinophoridaeLabiduridaeHEMIPTEROIDEAPHTIRAPTERA "Löss"

ANOPLURA "Åkta löss"

Echinophthiriidae

Echinophthirius Giebel,1871

E. horridus (von Olfers,1816) "Sallus"

= *E. sericaria* Meinerz,18???

= *E. setosus* ??17???

= *E. phocaiae* Lucas,17???

@ earless seals, e.g. *Phoca vitulina*
Europe; e.g. around Sweden, Denmark, Netherlands, Arctic Seas

Antarctophthirus Enderlein,1906

A. trichechi (Bohemann,17???)
@ *Odobenus rosmarus*
Arctic Seas

A. microchir (Trouessart & Newman,1888)
@ *Otariidae* & *Phoca vitulina*
California

A. callorhini (Osborn,1899)

@ *Callorhinus ursinus*

Alaska

Proechinophthirus Ewing,1923

P. fluctus (Ferris,1916)

@ *Callorhinus ursinus* & *Eumetopias jubatus*

Arctic Seas, Alaska

MALLOPHAGA "Fjäderätere", "Fagellöss" (& "Pälsätere")

@ birds

* No check-list, but names of a few genera that may be found on sea birds, have been listed here.

This is a group, which is considered very difficult, even by specialists.

AMBLYYCERAMenoponidae

Eidmanniella Keler,1938

@ e.g. *Sula*, *Phalacrocorax*

Sweden, etc.

Actornitophilus Ferris,1916

@ e.g. *Larus*, *Rissa*

Sweden, etc.

Colpocephalum Nitzsch,1818

@ e.g. *Larus*, *Rissa*

Sweden, etc.

Comatomenopon Uchida,1920

@ *Laridae*

Sweden, etc.

Ancistrona Westwood,1874

@ Procellariiformes

Sweden, etc.

Procellariiphaga Eichler,19???

@ Procellariiformes

Sweden, etc.

Piagetiella Neumann,1906

@ e.g. *Phalacrocorax*

?

Austromenopon Bedford,1939

@ e.g. *Larus*, *Rissa*, *Alca*, *Fulmarus*

Sweden, etc.

A. crocatum Nitsch,18???, Giebel

@ *Haematopus*

Br. Isles

Austromenopea ??,19???

* Is this *Austromenopon* wrongly spelled?

A. uriae Timmermann,1???

@ *Uria aalge*

N Atlantic, British Columbia

ISCHNOCERAPhilopteridae

Anaticola Clay,1935

Anatoecus Cummings,1916

Cunningsiella Ewing,1930

= *Dollabella* Cummings,1916, *menopon* de Lamarck,1801 (Mollusca)

Quadraceps Clay & Meinertzhagen,1939

@ e.g. *Larus*, *Rissa*, *Alca*

Sweden, etc.

Q. obliquus (Mjöberg,1???)

@ *Uria aalge*

North Atlantic, British Columbia

Saemundssonia Timmermann,1936

@ e.g. *Larus*, *Rissa*, *Alca*, *Fulmarus*

Sweden, etc.

S. calva (Kellogg,1???)

@ *Uria aalge*

North Atlantic, British Columbia

Perineus Thompson,1936

@ Procellariiformes, Laridae

Sweden, etc.

Halipeurus Thompson,1936

@ Procellariiformes.

Sweden, etc.

Pectinopygus Mjöberg,1910

@ e.g. *Sula*, *Phalacrocorax*

Sweden, etc.

P. bassani (O.F. Müller,1776)

@ *Sula bassana*

Br. Isles

Philopterus Nitzsch,1818

P. gonothorax Giebel,18???

@ *Larus marinus*

Br. Isles

HEMIPTERA= HETEROPTERA "Halvvingar"

* The "most marine" species of this order is to be found in the suborder Gerromorpha, where four families, Gerridae, Veliidae, Mesoveliidae and Hermatobatidae together have a dozen genera, e.g. *Halobates*, with both estuarine and oceanic species. However none of these more than 80 species is to be found in Europe. They occur in warmer seas.

GERROMORPHAGerroidea

- Gerridae "Vattenlöpare"
- Gerris** J.C. Fabricius,1794
- G. odontogaster** (Zetterstedt,1828)
* A typical brackish water species. Limnohaline.
Baltic, Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- G. thoracicus** Schummel,1832, ?non Stephens,1829
* A typical brackish water species.Pseudohalobiont. Halophilic.
Baltic, Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- G. rufoscutellatus** Latreille,1807
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- G. argentatus** Schummel,1832
Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, France, ?Br. Isles
- G. lacustris** (Linnaeus,1758)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- Veliidae "Bäcklöpare"
- Microvelia** Westwood,1834
- M. reticulata** (Burmeister,18??)
* A typical brackish water species.
Baltic
- Mesovelidae** Douglas & Scott,1867
- Mesovelinae** China & Miller,1959
- Mesovelia** Mulsant & Rey,1852
- M. furcata** Mulsant,1???
* A typical brackish water species. Limnohaline.
Baltic, Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, France, Br. Isles
- NEPOMORPHA
- Nepoidea
- Nenidae "Vattenskorpcioner"
- Nepa** Linnaeus,1758
- N. rubra** Linnaeus,1758
* A typical brackish water species.
Baltic
- N. cinerea** Linnaeus,1758
Baltic
- Notonectoidea
- Notonectidae "Ryggsmärra"
- Notonecta** Linnaeus,1758
- N. viridis** Delcourt,18??
* Halophilic.
Germany, Netherlands, England
- N. glauca** Linnaeus,1758
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Corixoidea
- Corixidae "Buksimärra"
- Cymatinae
- Cymaria** Flor,1860
- C. bonsdorffii** (Sahlberg,1819)
Baltic
- C. coleoptrata** (J.C. Fabricius,1777)
Baltic, Sweden, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- C. rogenhoferi** Fieber,18??
Europe
- Micronectinae
- Micronecta** Kirkaldy,1897
- M. meridionalis** (A. Costa,1847)
Europe
- M. minutissima** (Linnaeus,1758)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Corixinae
- Arctocoris** Wallengren,1894
- A. carinata** (Sahlberg,1819)
Baltic
- Callicorixa** B. White,1873
- C. praecusta** (Fieber,1848)
Baltic, Sweden, Norway, Denmark, Finland, Balticum, Germany, North Sea Shores, France, Br. Isles
- C. producta** (Reuter,1???)
Baltic, Sweden, Finland
- Corixa** Geoffroy,1762
- C. affinis** Leach,1817
German Baltic coast, England
- C. panzeri** (Fieber,1848)
Sweden, Denmark, Germany, North Sea Shores, France, Br. Isles
- Hesperocorixa** Kirkaldy,1908
* Should spelling be *Hesperocrixa* ?
- H. sahlbergi (Fieber,1848)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- Sigara** J.C. Fabricius,1775
- S. assimilis** (Fieber,1848)
Europe
- S. concinna** (Fieber,1848)
Europe
- S. dorsalis** (Leach,1817)
England
- S. fatteni** (Fieber,1848)
England, German Baltic coast
- S. lateralis lateralis** (Leach,1817)
Europe, e.g. Baltic
- S. selecta** (Fieber,1848)
Sweden, Norway, Denmark, Germany, North Sea Shores, France, Br. Isles
- S. stagnalis** (Leach,1817)
= *S. lugubris* Fieber,18??
* A typical brackish water species. Halophilic-Halobiont.
Baltic, Netherlands, Br. Isles
- S. distincta** (Fieber,1848)
* Halophilic.
Sweden, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- S. striata** (Linnaeus,1758)
* A typical brackish water species. Limnohalin.
Baltic, Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- S. venusta** Douglas & Scott,1???
France
- LEPTOPODOMORPHA
- Leptopodoidea
- Salidae "Strandstinkflyn"
- Saldinae
- Salda** Fabricius,1803
- S. littoralis** (Linnaeus,1758)
* Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Chartoscirta** Stål,1868
- C. cincta** (Herrich-Schaeffer,1841)
Europe, e.g. Sweden
- C. elegantula** (Fallén,1807)
Sweden
- Halosalda** Reuter,1912
- H. lateralis** Reuter,1912
* Halobiont.
Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, France, Br. Isles
- Saldula** Van Duzee,1914
- S. pilosella** (Thomson,18???)
* Halobiont.
Sweden, Norway, Finland, Balticum, Br. Isles
- S. scotica** (Curtis,18???)
* Halophilic.
Sweden, Norway, Finland, Balticum, Br. Isles
- S. saltatoria** (Linnaeus,1758)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- S. pallipes** (J.C. Fabricius,1781)
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Micracanthia** Reuter,1912
* Right subfamily?
- M. fennica** Reuter,1912
* Halobiont.
Sweden, Finland, Balticum, ?Norway
- Chiloxyanthinae
- Chiloxyanthus** Reuter,1891
- C. pilosus** (Fallén,1807)
* Halobiont.
Sweden, Denmark, Germany, Netherlands, France, England
- Pentacora** Reuter,1912
- Aepophilinae "Havstinkflyn"
- Aepophilus** Signoret,1879
- A. bonnairei** Signoret,1879, in Puton
* Halobiont.
Br. Isles, N. France, Ireland, Iceland
- CIMICOMORPHA
- Miroidea
- Miridae**
= *Capsidae* Burmeister "Ängsstinkflyn"
- Mirinae** Dohrn,1858
- Mirini** Dohrn,1858
- Poeciloscytus** Fieber,1858
- P. vulneratus** (Panzer,1798)

- = *Polymerus* Hahn,1831 (*Poeciloscytus*) *vulneratus* (Panzer,1798)
 @ e.g. *Salsola*, *Galium*, *Corispermum*, etc. * Halophilic.
 Sweden, Denmark, Germany, Netherlands, France, England
- Stenodemini* China,1943
- Teratocoris* Fieber,1858
- T. saundersi* Douglas & Scott,1867
 @ *Phragmites* * Halophilic.
 Sweden, Norway, Finland, Denmark, Balticum, Germany, Netherlands, Br. Isles
- Trigonotylus* Fieber,1858
- T. psammæcolor* Reuter,1885
 @ *Ammophila* & *Elymus* * Halophilic.
 Sweden, Denmark, Germany, Netherlands, England
- Phylinae Handlirsch,1925
- Phylini Douglas & Scott,1845
- Conostethus* Fieber,1858
- C. salinus* (J. Sahlgren,1870)
 @ Halophytes
 Sweden, Denmark, Netherlands, England, German Baltic coast
- C. frisicus* E. Wagner,1952
 @ *Limonium*
 German North Sea coast
- C. brevis* Reuter,1877
 @ *Limonium*
 Scotland
- PENTATOMOMORPHA
- Piesmatoidea
- Piesmatidae
- Piesma* Peletier & Serville,1825
- P. quadrata* (Fieber,1844?)
 * Halophilic.
 Sweden, Norway, Finland, Denmark, Germany, North Sea Shores, England
- P. salsolae* Becker,18??
 * Halophilic.
 Finland, France
- Lygaeoidea
- Lygaeidae
- Henestaris* Spinola,1837
- H. halophilus* (Burmeister,1835)
 * Halobiont.
 Germany, France, England
- H. laticeps* (Curtis,1836)
 * Halobiont.
 France, England
- Trapezonotus* Fieber,1860
- T. distinguendus* Flor,18??
 * Halophilic.
 Sweden, Norway, Finland, Denmark, Balticum, Germany, Netherlands, England
- Beosus Amyet & Serville,1843
- B. maritimus* (Scopoli,1763)
 * Halophilic.
 Germany, Netherlands, France, England
- HOLOMETABOLA
- NEUROPTERA "Nätvingar"
- MEGALOPTERA
- Sialidae "Sävsländor"
- Sialis* Latreille,1802
- S. sp.*
 Baltic
- TRICHOPTERA "Nattsländor"
 * Some species represented by larvae only in very "diluted" estuarine water, have not been listed here.
- Hydropsychoidea
- Hydropsychidae
- Hydropsyche* Pictet,1834
- H. sp.*
 Baltic
- Polycentropodidae
- Polycentropus* Curtis,1835
- P. flavomaculatus* Pictet,18??
 Baltic
- Limnephiloidea
- Limnephilidae
- Limnephilus* Leach,1815
- L. affinis* (Curtis,1827?)
 * The only true European estuarine-marine species. Tolerates a salinity of up to 24 ppt.
 Baltic, Br. Isles
- L. marmoratus* Curtis,1830
- N. Baltic
- L. decipiens* (Kolenati,1848)
 N. Baltic
- L. lunatus* Curtis,18??
 N. Baltic
- L. ignavus* McLachlan,18??
 N. Baltic
- Phryganeidae
- Phryganea* Linnaeus,1758
- P. grandis* Linnaeus,1758
 N. Baltic
- P. striata* Linnaeus,1758
 N. Baltic
- Agrypnia* Curtis,1835
- A. pagetana* Curtis,1835
 N. Baltic
- Agrypnetae* McLachlan,1876
- A. sp.*
 Baltic
- Molannidae
- Molanna* Curtis,1834
- M. angusta* (Curtis,1828?)
 N. Baltic
- Sericostomatidae
- Lepidostoma* Rambur,1842
- L. hirtum* (Fabricius,17??)
- Leptoceridae
- Athripsodes* Billberg,1820
- A. aterrimus* (Stephens,18???)
 Baltic
- Oecetis* McLachlan,1877
- O. ochracea* Curtis,1925
 SW Sjælland, in ≈ 10 ppt S.
- O. furva* (Rambur,1842?)
 N. Baltic
- ?Superfamilies? & ?Families?
- Grammotaulius* Kolenati,1848
- G. atomarius* (J.C. Fabricius,1793), ?non *Phryganea atomaria* J.F. Gmelin,1790
 N. Baltic
- Colpotauelius* Kolenati,1848
- C. incisus* (Curtis,1834)
 N. Baltic
- Ptoceridae
 * Taxa from a "Baltic Invertebrate Code-list". Not listed in Parker (Ed.),1982
- Lycentropus* ?,1???
- L. sp.*
 Baltic
- Raylea* ?,1???
- R. sp.*
 Baltic
- Agraylea Curtis,1834
- A. sp.*
 Baltic
- LEPIDOPTERA "Fjärilar"
- * This part of the list is certainly much incomplete. Only some, more or less obligate, associates of maritime herbs are listed.
- DITRYSIA
- Tortricoidea "Vecklare"
- Tortricidae Stephens,1829
- Olethreutinae* Walsingham,1897
- Olethreutini* Obraztsov,1946
- Bactra* Stephens,1834 (TM *Olethreutes lanceolana* Hübner,1796-99)
- B. robustana* Christoph,1872
 @ *Scirpus maritimus*
 Bohuslän, Halland, Skåne, Denmark, Germany
- Lobesia* Guenée,1845 (TM *Asthenia reliquana* Hübner,1825)
- L. (Lobesia) littoralis* Westwood & Humphreys,1845
 @ *Armeria maritima*
 Bohuslän, Halland, Skåne, Denmark, Norway, Germany, Netherlands
- Eucosmini* Obraztsov,1946
- Eucosma* Hübner,1823 (TM *E. circulana* Hübner,1823)
- E. (Phaneta* Stephens,1852) *tripoliana* Barrett,1880 (TM of *Phaneta* is *Cochylis pauperana*
 @ *Aster tripolium* Duponchel,1842)
 W Jylland, N. Germany

- E. (Phaneta* Stephens,1852) *krygeri* Rebel,1937
@ *Artemisia maritima*
Sjælland (Amager)
- E. (Phaneta* Stephens,1852) *maritima* Westwood & Humphreys,1845
@ *Artemisia* spp.
N Germany
- Tortricinae
- Aethes* Billberg,1820
- A. ditucidana* ?,1???
@ *Angelica archangelica litoralis*
Bohuslän, Halland, Netherlands, Britain
- Phalonidia* Le Marchand,1933
* Placed in correct superfamily & family?
- P. affinitana* Dgl.,1???
@ *Aster tripolium*
Bohuslän, Halland, Skåne, Denmark
- P. vectisana* Humphreys & Westwood,1845
@ *Triglochin maritimum*
Halland, Skåne, Denmark, Finland
- Pterophoroidea
- Pterophoridae "Fjädermott"
- Agdistis* Hübner,1825
- A. benneti* (Curtis,18???)
@ *Limonium*
S. Sjælland
- Pyraloidea
- Pyralidae "Mott"
- Acentropus* Curtis,1834
- A. niveus* (Olivier,1791)
* in sheltered, vegetation-rich brackish water (up to ≈20 ppt S) lagoons (& in fresh water)
Baltic, Danish estuaries
- Crambinae
- Pediasia* Hübner,1825
- P. aridella* (Thunberg,1788) "en gräsmotts-art"
* Halophilic.
Halland
- Tineoidea "Malar"
- Bucculatrigidae
- Bucculatrix* Zeller,1839
- B. maritima* Stainton,1851
@ *Aster tripolium*
Bohuslän, Halland-S Baltic shores, Denmark, Finland
- Gelechioidae
- Oecophoridae
- Depressiniae
- Agonopterix* Hübner,1825
- A. zephyrella* ?,1???
@ *Angelica archangelica litoralis*
Halland, Germany, Netherlands, Britain
- Gelechiidae
- Gelechiinae
- Scrobipalpa* Janse,1951
- S. salicorniae* (E. Hering,1???)
@ *Salicornia*, *Suaeda*, etc.
Halland, central Baltic shores, Denmark, Finland
- S. stangei* (E. Hering,1889)
@ *Triglochin maritimum*
Bohuslän, Halland-central Baltic Shores, Denmark, Finland
- S. plantaginella* (Stainton,18???)
@ *Plantago maritima*
Bohuslän, Halland-S Baltic shores, Denmark, Finland
- S. nitentella* (Fuchs,1902)
@ *Atriplex*, *Suaeda* & *Chenopodium*
Göteborg archipelago, Skåne-central Baltic shores, Denmark, Finland
- Aristotelia Hübner,1825
- A. brizella* Treitschke,1833
@ *Armeria maritima*
Halland-central Baltic shores, Norway, Denmark, Finland
- Coleophoridae "Säckdrägmalar"
- Coleophora* Hübner,1822
- C. atriplicis* Meyrick,1928
@ *Atriplex*, *Suaeda* & *Chenopodium*
Halland-central Baltic shores, Finland
- C. adjunctella* Hodgkinson,1882
@ *Juncus gerardi*
Halland-S & N Baltic shores, Denmark, Finland
- C. salicorniae* Wocke,1876
@ *Salicornia*
Halland, S Baltic shores, Denmark, Finland
- C. asteris* Mühlig,18???
@ *Aster tripolium*
Göteborg archipelago, Skåne, Denmark
- C. suaedivora* Durrant,1???
@ *Suaeda maritima*
É Småland, England
- Elachistidae
- Elachista* Treitschke,1833
- E. scirpi* Stainton,1???
@ *Scirpus maritimus* & *Juncus gerardi*
Bohuslän-S Baltic shores, Denmark
- DIPTERA "Tvåvingar"
* See also Yerbury, J. W., 1919. Seashore Diptera. *J. mar. biol Ass. U.K.* **12** : 141-145.
- NEMATOCERA "Myggor"
- Tipuloidea
- Tipulidae "Harkrankar"
- Tipulinae "Egentliga harkrankar"
- Tipula* Linnaeus,1758
- T. solstitialis* ?,1???
* In brackish water ditches.
Britain
- Limoniiinae "Småharkrankar"
- Limonini
- Limonia* Meigen,1803
- L. (Geranomyia* Haliday,1833) *unicolor* Haliday,1833
* Larvae in tufts of *Lichen pygmaea* and *Catenella repens*
S. England
- L. (Geranomyia* Haliday,1833) *bazzii* A. & L.,19??
* Larvae assumed to feed on *Enteromorpha*
Suffolk-Dorset, N. Africa
- L. (Dicranomyia* Stephens,1829) *goritiensis* (Mik,1???)
* In tiny streams, which make the cliff wet.
S. English coast
- L. sera* ?,1???
* Salt-marshes.
Britain
- L. complicata* ?,1???
* Salt-marshes.
Britain
- L. ventralis* ?,1???
* In brackish water ditches.
Britain
- Eriopterini
- Erioptera* Meigen,1803
- E. strictita* ?,1???
* In salt-marshes.
Britain
- E. bivittata* ?,1???
* In brackish water ditches.
Britain
- Gonomyia Meigen,1818
- G. conoviensis* ?,1???
* In tiny streams, which make the cliff wet.
Britain
- Psychodoidea
- Ptychopteridae
= *Liriopidae* "Glansmyggor"
- Ptychoptera* Meigen,1803
- P. contaminata* van der Wulp,18???
* In brackish water ditches.
Sweden, Britain
- Culicoidea
- Chironomidae
= *Tendipedidae* "Fjädermyggor"
- Telmatogotoninae Brundin,1966
- Thalassomyia* Schiner,1856
- T. frauendorfii* Schiner,1856
Br. Isles, France, Mediterranean
- Psamathomyia* Deby,1889
- P. pectinata* Deby,1889
Br. Isles, French Mediterranean coast
- Telmatogeton* Schiner,1866
=? *Parachano* Kieffer,1911

- T. remanei* Remmert,1963
Kiefer Bucht
- Orthocladiinae* Edwards,1929
- Ctinio* Haliday,1855
- C. marinus* Haliday,1855
Baltic-Bohuslän, Belgium, France, Br. Isles
- Halocladius* Hirvenoja,1973
* Includes some former *Cricotopus* species with larvae in saline waters. Probably some of those beneath.
- H. fucicola* (Edwards,1926)
= *Cricotopus fucicola* (Edwards,1926)
* Intertidal, on rocky shores.
Br. Isles
- H. varians* (Staeger,1839)
= *Cricotopus halophilus* Kieffer,1909
= *Cricotopus marinus* Alverdes,1911
* Brackish water species. Muns,1967 found *Chironomus halophilus* Kieffer (Strenzke) in most Danish estuaries, between 1-16 ppt S. Is he referring to this species?
Kiefer Bucht, Öresund, Bergen, Helgoland, Br. Isles, Belgium, Normandie, ?Faroes, ?Iceland
- Cricotopus* van der Wulp,1874
- C. obnixus* (Walker,1856)
SW. England
- C. annulator* Goetghebuer,1927
Baltic, Belgium
- C. bicinctus* (Meigen,1818)
= *Trichocladus dizonias* (Meigen,1830)
Baltic, Belgium
- C. caducus* Hirvenoja ,19??
Baltic
- C. festivellus* (Kieffer,1906)
Baltic, Belgium
- C. flavocinctus* (Kieffer,1924)
Baltic, Belgium
- C. laetus* Hirvenoja,19??
Baltic
- C. obtusus* Hirvenoja,19??
Baltic
- C. ornatus* (Meigen,1818)
Baltic, Belgium
- C. pilatensis* (Zetterstedt,1850)
Baltic, Belgium
- C. pilosellus* Brundin,19??
Baltic
- C. sylvestris* (Fabricius,1794)
Baltic, Belgium
- C. triannulatus* (Macquart,1826)
Baltic, Belgium
- Psectrocladius* Kieffer,1906
- P. barbimanus* Edwards,1929
* Brackish water species
England
- P. sordidellus* (Zetterstedt,1838)
Baltic, Belgium
- P. ventricosus* Kieffer,1???
Baltic
- Smittia* Holmgreen,1869
- S. rupicola* (Kieffer, in Mercier,1923)
Germany, Belgium, France
- S. bacilligera* (Kieffer,1924)
German Baltic
- Pseudosmittia* Goetghebuer,1932
- P. brevifurcata* (Edwards,1926)
England
- Thalassosmittia* Strenzke & Remmert,1957
- T. thalassophila* (Bequaert & Goetghebuer,1913)
Belgium, England
- Corynoneura* Winnertz,1846
- C. marina* Kieffer,1924
Germany
- C. scutellata* Winnertz,1846
Baltic
- Metroclemens* van der Wulp,1874
- M. picipes* (Meigen,1818)
Scandinavia, Germany, Netherlands, Belgium, France, SW. England
- Paraphaenocladius* Thienemann,1924
- P. impensus* (Walker,1856)
SW. England
- Abiskomyia* Edwards,1937
- A. virgo* Edwards,19??
Baltic
- Acricotopus* Kieffer,1921
- A. thienemanni* Goetghebuer,19??
Baltic
- Camptocladius* van der Wulp,1874
- C. sp.*
Baltic
- Eukiefferiella* Thienemann,1926
- E. sp.*
Baltic
- Heterotanytarsus* Spärck,1923
- H. apicalis* Kieffer,19??
Baltic
- Heterotriassocladus* Spärck,1923
- H. grimshawi* (Edwards,1929)
Baltic, Belgium
- H. marcidus* (Walker,1856)
Baltic, Belgium
- H. maeae* Brundin,19??
Baltic
- H. subpilosus* (Kieffer,19???)
Baltic
- Limnophyes* Eaton,1875
- L. sp.*
Baltic
- Microcricotopus* Thienemann,1932
- M. balticus* Palmen,19??
Baltic
- M. bicolor* (Zetterstedt,18???)
Baltic
- Orthocladius* van der Wulp,1874
- O. excavatus* Brundin,19??
Baltic
- Parakiefferiella* Thienemann,1936
- P. bathopila* Kieffer,19??
Baltic
- Rheorthocladius* Thienemann,1935
- R. sp.*
Baltic
- Syncretotopus* Brundin,1956
- S. rufiventris* (Meigen,18???)
Baltic
- Trissocladius* Kiefer,1908
- T. sp.*
Baltic
- Tanypodinae Thienemann & Zavrel,1916
- Ablabesmyia* Johannsen,1905
- A. monilis* (Linnaeus,1758)
Baltic
- A. barbitarsis* (Zetterstedt,18???)
Baltic
- A. cingulata* (Walker,1???)
Baltic
- A. melanops* (Meigen,18???)
Baltic
- Anatopynia* Johannsen,1905
- A. plumipes* (Fries,1823)
Baltic, Belgium
- Clinotanytarsus* Kieffer,1913
- C. nervosus* (Meigen,1818)
Baltic, Belgium
- Macropelopia* Thienemann, in Thienemann & Kieffer,1916
- M. nebulosa* (Meigen,1804)
Baltic, Belgium
- Arctopelopia* Fittkau,1962
- A. griseipennis* (Van der Wulp,1858)
Baltic
- Procladius* Skuse,1889
- P. breviatus* Remmert,1???
- P. barbatus* Brundin,19??
Baltic
- P. crassinervis* (Zetterstedt,18???)
Baltic
- P. flavifrons* Edwards,19??
Baltic

- P. lundstroemi* Goetghebuer,18??
Baltic
- P. nigriventris* Kieffer,1???
Baltic
- P. nudipennis* Brundin,19??
Baltic
- Psectrotanypus* Kieffer,1909
- P. varius* (Fabricius,1787)
Baltic, Belgium
- Holotanypus* Roback,1982
- H. choreus* (Meigen,1804)
Baltic, Belgium
- H. signatus* (Zetterstedt,1850)
Baltic, Belgium
- Psilotanypus* Kieffer,1906
- P. sp.*
Baltic
- Tanypus* Meigen,1803
= *Tanypus* Rondani,1856
- T. punctipennis* Meigen,1818
Baltic
- Pentaneurini
- Thienemannimyia* Fittkau,1957
- T. fusciceps* Edwards,1???
Baltic
- T. tentiginosa* Fries,1???
Baltic
- T. vitellina* (Kieffer,18???)
Baltic
- Diamesinae* Kieffer,1923
- Dianesa* Meigen,1830
- D. sp.*
Baltic
- M. ekmani* Brundin,19??
Baltic
- Protanypus* Kieffer,1906
- P. morio* (Zetterstedt,18???)
Baltic
- Pothastia* Kieffer,1922
- P. sp.*
Baltic
- Prodiamesinae* Sæther,1976
- Prodiamesa* Kieffer,1906
- P. olivacea* (Meigen,1818)
Baltic, SW England
- Monodiamesa* Kieffer,1921
- M. bathyphila* Kieffer,1???
Baltic
- Chironominae Macquart,1838
- Camptochironomus* Kieffer,1918
- C. pallidivittatus* Malloch,1???
Baltic
- C. tentans* (Fabricius,1805)
Baltic, Belgium
- Chironomus* Meigen,1803
= *Tendipes* Meigen,1800
- C. satinarius* Kieffer,19??
* The larvae thrive in laboratory cultures with up to 50 ppt S.
Baltic, dominating species in Danish estuaries.
- C. halophilus* Meigen,18??
Baltic, Westfalen
- C. bicornutus* ?,1???
Westfalen
- C. aprilinus* Hoffmannsegg, in Meigen,1830
/Br. Isles
- C. annularius* Meigen,1818
= *C. annularis* (De Geer,1776)
Baltic, Belgium
- C. plumosus* (Linnaeus,1758)
Baltic
- C. anthracinus* Zetterstedt,18??
= *C. bathophilus* Kieffer,19??
Baltic
- C. thummi* Kieffer,1911
=? *C. gregarius* Meigen,18??
Baltic
- C. dorsalis* Meigen,18??
Baltic
- C. tenuistylus* Brundin,19??
Baltic
- Cryptochironomus* Kieffer,1918
- C. supplicans* (Meigen,1830)
Baltic, Belgium
- C. albifasciatus* (Staeger,1839)
Baltic, Belgium
- C. psittacinus* (Meigen,1830)
Baltic, Belgium
- Cryptocladopelma* Lenz,1941
- C. viridula* (Fabricius,17???)
Baltic
- C. edwardsi* Krus.,19??
Baltic
- Cryptotendipes* Lenz,1941
- C. usmaensis* (Pagast,1931?)
* Is this not a synonym of *Tanytarsus usmaensis* Pagast,1931?
Baltic
- C. paripes* Edwards,19??
Baltic
- Demicryptochironomus* Lenz,1941
- D. vulneratus* (Zetterstedt,1838)
Baltic, Belgium
- Einfeldia* Kieffer,1924
- E. insolita* Kieffer,19??
=? *E. dissidens* (Walker,1856)
Baltic
- Endochironomus* Kieffer,1918
- E. dispar* (Meigen,1830)
Baltic, Belgium
- E. impar* (Walker,18???)
Baltic
- E. intextus* (Walker,18???)
Baltic
- E. tendens* (Fabricius,1775)
Baltic, Belgium
- E. atipennis* (Meigen,1830)
=? *E. maritimus* Kieffer,1921
Belgium
- Fleuria* Kieffer,1924
- F. sp.*
Baltic
- Harnischia* Kieffer,1921
- H. curtilamellata* (Malloch,1915)
= *H. pseudosimplex* Goetghebuer,1923
Baltic, Belgium
- H. viridula* (Linnaeus,1767)
Baltic
- Kribioxenus* Kieffer,1921
- K. brayi* Goetghebuer,19??
Baltic
- Lauterborniella* Thienemann & Bause,1913
- L. agrayloides* (Kieffer,1911)
Baltic
- Lenzia* Kieffer,1922
- L. flavipes* (De Geer,1776)
Baltic
- L. punctipes* Wiedemann,1???
Baltic
- Dicrotendipes* Kieffer,1913
= *Limnochironomus* Kieffer,1920
- D. nervosus* (Staeger,1839)
Baltic, Belgium
- D. pulsus* (Walker,1856)
Baltic
- Microchironomus* Kieffer,1918
=? *Leptochironomus* Pagast,1931?
- M. tener* (Kieffer,1918)
= *Leptochironomus balticus* Pagast,1931?
Baltic, Belgium
- Microtendipes* Kieffer,1915
- M. brevitaris* Brundin,19??
Baltic
- M. chloris* (Meigen,1818)
Baltic, Belgium
- M. confinis* (Meigen,1830)
Baltic, Belgium
- M. pedellus* (De Geer,1776)
Baltic

<i>Pagastiella</i> Brundin,1949	<i>T. brundini</i> Lindeberg,19???
<i>P. orophila</i> (Edwards,1929) Baltic, Belgium	Baltic
<i>Parachironomus</i> Lenz,1921	
<i>P. arcuatus</i> (Goetghebuer,1919) Baltic, Belgium	
<i>P. mauricii</i> (Kruseman,1933) Baltic, Belgium	
<i>P. swammerdami</i> (Kruseman,19???) Baltic	
<i>P. varus</i> (Goetghebuer,1921) Baltic, Belgium	
<i>Paracladopelma</i> Harnisch,1923	
<i>P. campistolabis</i> (Kieffer,1913) Baltic	
<i>P. nigritula</i> Goetghebuer,19???	
<i>P. obscura</i> Brundin,19???	
<i>Paratendipes</i> Kieffer,1911	
<i>P. albimanus</i> (Meigen,1818) Baltic	
<i>P. nudisquama</i> Edwards,19???	
Baltic	
<i>Paralauterborniella</i> Lenz,1941	
<i>P. nigrohalteralis</i> Malloch,19???	
Baltic	
<i>Pentapeditum</i> Kieffer,1913	
<i>P. sordens</i> (van der Wulp,1874) Baltic, Belgium	
<i>P. tritum</i> (Walker,1856) Baltic, Belgium	
<i>Polypeditum</i> Kieffer,1912	
<i>P. birenatum</i> Brundin,19???	
Baltic	
<i>P. convictum</i> (Walker,1856) Baltic, Belgium	
<i>P. cultellatum</i> Goetghebuer,1931 Baltic, Belgium	
<i>P. lactum</i> (Meigen,1818) Baltic, Belgium	
<i>P. nubeculosum</i> (Meigen,1804) Baltic, Belgium	
<i>P. pullum</i> (Zetterstedt,1838) Baltic, Belgium	
<i>P. scataenum</i> (Schrank,1803) Baltic, Belgium	
<i>Pseudochironomus</i> Malloch,1915	
<i>P. prasinatus</i> (Staeger,1839) Baltic, Belgium	
<i>Sergentia</i> Kieffer,1922	
<i>S. coracina</i> (Zetterstedt,18???) Baltic	
<i>S. longiventris</i> Kieffer,19???	
Baltic	
<i>Stenochironomus</i> Kieffer,1919	
<i>S. sp.</i> Baltic	
<i>Stictochironomus</i> Kieffer,1919	
<i>S. crassiforceps</i> Kieffer,19???	
Baltic	
<i>S. histrio</i> (Fabricius,1794) Baltic, Belgium	
<i>S. pictulus</i> (Meigen,1830) Baltic, Belgium	
<i>S. rosenschoeldi</i> (Zetterstedt,18???) Baltic	
<i>Xenochironomus</i> Kieffer,1921	
<i>X. xenolabis</i> (Kieffer,1916) Baltic, Belgium	
<i>Glyptotendipes</i> Kieffer,1913	
<i>G. barbipes</i> (Staeger,1839) Baltic, Norway	
<i>Tanytarsus</i> van der Wulp,1874	
<i>T. brunnipes</i> (Zetterstedt,1850) Scandinavia, SW. England	
	<i>T. brundini</i> Lindeberg,19???
	Baltic
	<i>T. chinensis</i> Goetghebuer,1934
	Baltic, Belgium
	<i>T. curticornis</i> Kieffer,19???
	Baltic
	<i>T. decipiens</i> Lindeberg,19???
	Baltic
	<i>T. eminutus</i> (Walker,1856)
	Baltic, Belgium
	<i>T. gregarius</i> Kieffer,1909
	Baltic, Belgium
	<i>T. heusdensis</i> Goetghebuer,1923
	Baltic, Belgium
	<i>T. usmaensis</i> Pagast,1931
	Baltic, Belgium
	<i>T. holochlorus</i> Edwards,1929
	Baltic, Belgium
	<i>T. testagei</i> Goetghebuer,1922
	Baltic, Belgium
	<i>T. lugens</i> Kieffer,19???
	Baltic
	<i>T. norvegicus</i> Kieffer,19???
	Baltic
	<i>T. signatus</i> (van der Wulp,1859)
	Baltic, Belgium
	<i>T. socialis</i> Lindeberg,19???
	Baltic
	<i>T. telmaticus</i> Lindeberg,19???
	Baltic
	<i>Cladotanytarsus</i> Kieffer,1922
	<i>C. atridorsum</i> Edwards,19???
	Baltic
	<i>C. mancus</i> (Walker,1856)
	Baltic, Belgium
	<i>C. wexionensis</i> Brundin,19???
	Baltic
	<i>Constempellina</i> Brundin,1947
	<i>C. brevicosta</i> Edwards,19???
	Baltic
	<i>Corynocera</i> Zetterstedt,1837
	<i>C. ambigua</i> Zetterstedt,18???
	Baltic
	<i>Lauterbornia</i> Kieffer,1911
	<i>L. coracina</i> (Zetterstedt,18???)
	Baltic
	<i>Micropsectra</i> Kieffer,1909
	<i>M. groenlandica</i> Andersen,1937?
	Baltic
	<i>M. monticola</i> Edwards,19???
	Baltic
	<i>M. praecox</i> (Meigen,18???)
	= <i>M. merripes</i> Kieffer,19???
	Baltic
	<i>Paratanytarsus</i> Tienemann & Bause,1913
	<i>P. confusus</i> Palmen,19???
	Baltic
	<i>P. inoperatus</i> (Walker,1856)
	Baltic, Belgium
	<i>P. intricatus</i> (Goetghebuer,1921)
	Baltic, Belgium
	<i>P. iaccophilus</i> Edwards,19???
	Baltic
	<i>P. natvigi</i> Goetghebuer,19???
	Baltic
	<i>P. penicillatus</i> Goetghebuer,19???
	Baltic
	<i>P. tenuis</i> (Meigen,1830)
	Baltic, Belgium
	<i>Rheotanytarsus</i> Thienemann & Bause,1913
	<i>R. sp.</i>
	Baltic
	<i>Stempellina</i> Thienemann & Bause,1913
	<i>S. bausei</i> (Kieffer,19???)
	Baltic
	<i>S. subgabriennnis</i> Brundin,19???
	Baltic
	<i>Stempellinella</i> Brundin,1947
	<i>S. brevis</i> Edwards,19???

Baltic	
<i>S. minor</i> (Edwards,1929) Baltic, Belgium	Sweden, Br. Isles, European mainland
<i>Zavrelia</i> Kieffer, in Thienemann & Bause, 1913	<i>C. (Culiseta) subochrea</i> (Edwards,1921) * Larvae may tolerate slightly brackish water? Scandinavia, Br. Isles, European mainland
<i>Z. pentatoma</i> Kieffer, 1913 Baltic, Belgium	<u>Anophelinae</u>
<u>Heleidae</u> = Ceratopogonidae "Svidknott"	<i>Anopheles</i> Meigen, 1818
<i>Culicoides</i> Latreille, 1809	<i>A. (Anopheles) atroparvus</i> van Thiel, 1927 * The only North European malaria vector. Earlier considered to be a subspecies of <i>A. maculipennis</i> Meigen, 1818. Coastal distribution. Tolerates high salinities. Sweden, S. Baltic, German North Sea coast, Netherlands
<i>C. furens</i> (Poey, 18???) Br. Isles	<i>A. (Anopheles) algeriensis</i> Theobald, 1903 * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, N. Germany, Mediterranean countries
<i>C. sp.</i> Baltic	<i>A. (Anopheles) claviger</i> (Meigen, 1804) * Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, Ireland, European mainland
<i>Dasyhelea</i> Kieffer, 1911	<u>Chaoboridae</u> = Corethridae "Tofsmyggor"
<i>D. flavoscutellata</i> (Zetterstedt, 1850) Scandinavia, Britain	<u>Chaoborinae</u>
<i>Monoculicoides</i> Khalaf, 1954	<i>Chaoborus</i> Lichtenstein, 1800
<i>M. nubeculosus</i> (Meigen, 18???) Britain	<i>C. (C.) crystallinus</i> (De Geer, 1776) = <i>C. plumicornis</i> (Fabricius, 1794) = <i>C. fuscus</i> (Staeger, 1839) Baltic, Belgium
<i>Bezzia</i> Kieffer, 1899	<i>C. (C.) flavicans</i> (Meigen, 1830) Baltic, Belgium
<i>B. sp.</i> Baltic	BRACHYCERA "Flugor" * Almost only halobionts have been listed
<i>Palpomyia</i> Meigen, 1818, ex Megerle MS	ORTHORRHAPHA
<i>P. sp.</i> Baltic	<u>Tabanoidea</u>
<i>Sphaeromais</i> ?,1???	<u>Tabanidae</u> "Bromsar"
<i>S. sp.</i> Baltic	<u>Tabaninae</u>
<i>Leptoconops</i> Skuse, 1889	<i>Tabanus</i> Linnaeus, 1758
<u>Dixidae</u>	<i>T. bovinus</i> Linnaeus 1758 "Fäbroms", "Nötbroms" Sweden, German Baltic coast
<i>Dixa</i> Meigen, 1818	<i>Hybomitra</i> Enderlein, 1922
Subgen.: <i>Dixella</i> Dyar & Shannon, 1924	<u>Chrysopsinae</u> "Blinningar"
<i>D. (Dixella) attica</i> Pandazis, 19???	<i>Chrysops</i> Meigen, 1800
Britain	<i>C. relicta</i> Hoffmannsegg, in Meigen, 1820 "Blindbroms", "Guldögonbroms" * In salt-marshes Sweden, Germany
<u>Culicidae</u> "Stickmyggor"	<i>Haematopota</i> Meigen, 1803 "Egentliga blinningar"
<u>Culicinae</u>	<i>H. grandis</i> ?Macquart, 1834? * In salt-marshes Britain
<i>Aedes</i> Meigen, 1818, ex Hoffmannsegg MS	<i>H. bigota</i> ?,1???
<i>A. detritus</i> (Haliday, 1833) = <i>A. salinus</i> Ficalbi, 1896 = <i>Ochlerotatus</i> Lynch Arribalzaga, 1891 <i>detritus</i> (Haliday, 1833) * Chiefly coastal. Tolerates high salinities. S. Baltic, Denmark, Br. Isles, Ireland, France	* In salt-marshes Britain
<i>A. caspius</i> (Pallas, 1771) = <i>Ochlerotatus</i> Lynch Arribalzaga, 1891 <i>caspicus</i> (Pallas, 1771) * A salt tolerant species which also commonly is found in non-saline inland waters. along the European coast from Finland-England-Atlantic-Mediterranean	<i>Silvius</i> Meigen, 1820
<i>A. dorsalis</i> (Meigen, 1830) = <i>Ochlerotatus</i> Lynch Arribalzaga, 1891 <i>dorsalis</i> (Meigen, 1830)	<u>Stratiomyidae</u> "Vapenflugor"
* Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, N. & central Europe	<u>Stratiomyinae</u>
<i>A. flavescentes</i> (O.F. Müller, 1764)	<i>Stratiomys</i> Geoffroy, 1762
* Not coastal, but larvae may develop in low-medium salinity brackish water. Br. Isles, N. & central Europe	<i>S. longicornis</i> ?,1???
<i>A. leucomelas</i> (Meigen, 1804)	* In brackish water ditches. Britain
* Not coastal, but larvae may develop in low-medium salinity brackish water. Br. Isles, N. Europe	<i>S. furcata</i> Fabricius, 1775 * In brackish water ditches. Sweden, Britain
<i>Culex</i> Linnaeus, 1758	<u>Citellarinae</u>
<i>C. (Barraudius</i> Edwards, 1921) <i>modestus</i> Ficalbi, 1889	<i>Nemotelus</i> Geoffroy, 1762
* Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, Mediterranean countries	<i>N. uliginosus</i> ?,1???
<i>C. (Culex) pipiens</i> Linnaeus, 1758	* Larvae develop in brackish water. Sweden, Britain
* Not coastal, but larvae may develop in low? salinity brackish water. Sweden, Br. Isles, temperate areas of the world	<i>N. notatus</i> ?,1???
<i>C. (Culex) torrentium</i> Martini, 1925	* Larvae develop in brackish water. Britain
* Not coastal, but larvae may develop in low salinity brackish water. Br. Isles, N. & central Europe	<u>Asiloidae</u>
<i>C. (Neoculex</i> Dyar, 1925) <i>territans</i> Walker, 1856	<i>Asilidae</i> Leach, 1819 "Rovflugor"
* Not coastal, but larvae may develop in low salinity brackish water. Baltic area, Br. Isles, European mainland	<u>Asilinae</u> Macquart, 1838, emend. Schiner, 1862
<i>Cutiseta</i> Felt, 1904 = <i>Theobaldia</i> Neveu-Lemaire, 1902	<i>Philonicus</i> Loew, 1849
<i>C. (Allotheobaldia</i> Brolemann, 1919) <i>longiareolata</i> (Macquart, 1838)	<i>P. albiceps</i> (Meigen, 1820) * Coastal sand dunes. Prey is peculiar to seashore. ?Scandinavia, Britain, Germany
* Larvae may develop in both fresh and brackish water. Br. Isles, Mediterranean countries	<u>Empidoidea</u>
<i>C. (Culisetta</i> Felt, 1904) <i>litorea</i> (Shute, 1928)	<i>Empididae</i> "Dansflugor"
* Larvae tolerates slightly brackish water. Br. Isles, France, Spain	<i>Hilara</i> Meigen, 1822
<i>C. (Culisetta) morsitans</i> (Theobald, 1901)	<i>H. lundbecki</i> Frey, 19???
* Larvae tolerates slightly brackish water. Scandinavia, Br. Isles, European mainland	
<i>C. (Culiseta) annulata</i> (Schrank, 1776)	
* Larvae tolerates brackish water. Our largest Culicidae species.	

Sweden, Denmark, Germany, Netherlands, Br. Isles

Chersodromia Walker,1849, Haliday MS

* In wrack-beds (in moist wrack).

C. arenaria (Haliday,1833)

Sweden, Denmark, Norway, Finland, Germany, Netherlands, S. England

C. difficilis Lundbeck,1910

Sweden, Denmark, Finland, France, S. England

C. incana Walker,18??

Sweden, Denmark, Netherlands, France, S. England

C. speculifera Walker,18??

Denmark, Germany, France, S. England

C. curvifrons (Zetterstedt,1842)

Bohuslän, Denmark, Finland, Netherlands, France, S. England

Coloboneura Melander,1902

C. hirta (Walker,18???)

= *Chersodromia hirta* Walker,18???

Sweden, Denmark, Germany, Netherlands, France, S. England

Dolichopodidae "Stylflugor"

Dolichopodinae

Dolichopus Latreille,1796

D. sabinus ?,1???

* In salt-marshes

Britain

D. plumipes (Scopoli,1763)

= *Musca plumipes* Scopoli,1763

* In salt-marshes

Sweden, Britain

Hygrocoleuthus Loew,1857

H. latipennis (Fallén,1823)

Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium

H. diadema (Haliday,1832)

Sweden, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

Hydromorphinae

Hydromorphorus Fallén,1823

H. bisetus Loew,18???

Denmark, Germany, Netherlands, France, Br. Isles

H. praecox (Lehmann,18???)

Sweden, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, S. England

H. oceanus (Macquart,1???)

* Bare mud in salt-marshes.

S. England

Liancalus Loew,1857

L. virens (Scopoli,1763)

= *Musca virens* Scopoli,1763

* In tiny streams, which make the cliff wet.

Sweden, Britain

Thinophilus Wahlberg,1844

T. flavipalpis Zetterstedt,1844, ?non Schiodte,1844

Sweden, Finland, Balticum, Germany, Netherlands, France, Br. Isles

T. ruficornis Haliday,18???

Sweden, Finland, Balticum, Denmark, Germany, Netherlands, France, Br. Isles

Schoenophilus Mik,1878

S. versutus Walker,18???

Sweden (Skåne), Denmark, Germany, Netherlands, Belgium, France, Br. Isles

Subfamily?

Muscidideicus Becker,1917

M. praetextatus Haliday,18???

German North Sea coast, Belgium, France, Br. Isles

Orthoceratium Schrank,1803

O. lacustre (Scopoli,1763)

German North Sea coast, Belgium, Br. Isles

Aphrosylus Haliday, Walker,1851

* Larvae may be predators on barnacles

A. ferox Haliday, Walker,1851

Netherlands, France, S. England

A. mitis Verrall,18???

France, S. England

A. celtiber Haliday,1851?

* Is the name *A. celicer*?

France, S. England

A. raptor Haliday, Walker,1851

France, S. England

A. venator Loew,18???

France

Machaerium Haliday,1832

M. maritimae Haliday,1832

* In salt-marshes

Denmark, Germany, France, S. England

CYCLORRAPHA

ASCHIZA

Phoroidea

Phoridae "Puckelflugor"

Syrphoidea

Syrphidae "Blomflugor"

Lathyrophthalmus Mik,1897

L. aeneus (Scopoli,1763)

= *Musca aena* Scopoli,1763

= *Eristalinus* Rondani,1845 *aeneus* (Scopoli,1763)

* In salt-marshes. Larvae probably in small pools in the splash zone.
Britain, Germany

Eristalis Latreille,1804 "Slamflugor"

E. abusivus ?,1???

* In salt-marshes.

Britain

E. sepulchralis (Linnaeus,1758)

= *Musca sepulchralis* Linnaeus,1758

@ on thrift blossoms (*Armeria maritima*)

Sweden, S. England

Helophilus Leach,1817 (Emend. pro *Elophilus* Meigen,1903)

H. parallelus ?,1???

* In brackish water ditches.

Britain

Lejops Rondani,1857

= *Liops* Verrall,1901, non Fieber,1870 (Hemiptera)

L. vittata (Meigen,18???)

* In brackish water ditches.

Sweden, Britain

Tropidia Meigen,1822

T. scita ?,1???

* In brackish water ditches.

Britain

SCHIZOPHORA

Tephritoidea

Otitidae

= Otafidae "Fläckflugor"

Melieria Robineau-Desvoidy,1830

= *Meliera* Becker,1903

M. omissa Meigen,18???

Finland, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

M. picta (Meigen,1826)

Germany, Netherlands, France, Br. Isles

M. cana Loew,18???

Germany, Netherlands, France, Br. Isles

Ceroxys Macqurt,1835

C. urticae ?,1???

* In brackish water ditches.

Britain

Tephritidae

= Trypetidae = Trypaneidae "Fruktflugor", "Borrflugor"

Paroxyna Hendel, Lindner,1927

P. plantaginis ?,1???

@ *Aster tripolium*

Britain

Sciomyzoidea

Sciomyzidae "Kärrflugor"

Salicella Robineau-Desvoidy,1830

S. fasciata (Meigen,1830)

N. France, Ireland, S. Europe

Helcomyzidae

Helcomyza Curtis,1825

H. ustulata Curtis,1825, Meigen MS

* One of the main species in wrack-beds (in fairly dry wrack)

Sweden, Finland, Denmark, Germany, Netherlands, Belgium, France, Scotland

Heterochaeta Rondani,1857

H. buccata (Fallén,1820)

* One of the main species in wrack-beds (in moist wrack).

Skåne, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

Chamaemyiidae "Markflugor"

Chamaemyia Meigen,1803

C. maritima (Zetterstedt,1846)

Sweden, Denmark, Germany, Netherlands, Belgium, France, Br. Isles

C. flavigalpus Haliday,18???

Britain

Coelopidae "Tångflugor"

* ≈ 6 Swedish species.

Coelopa Meigen,1830

= *Fucomyia* Haliday, Curtis,1837

- C. frigida* (J. C. Fabricius,1805)
 = *C. parvula* Haliday,18??
 = *C. eximia* Stenhammar,18??
 * One of the main species in wrack-beds (in moist wrack).
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- C. pilipes* Haliday,1838
 * One of the main species in wrack-beds (in moist wrack).
 Sweden, Finland, Germany, Belgium, France, S. England, Scotland
- Malacomysia* Westwood,1840, ex Haliday MS
Malacomysia Curtis,1837, ex Haliday MS, non Wesmael,1836 (Neuroptera)
- M. sciomyzina* (Haliday,1833)
 = *Coelopa furcata* Zetterstedt,1847
 * One of the main species in wrack-beds (in moist wrack).
 Bohuslän, Balticum, Germany, Netherlands, France, S. England
- Oedoparea* Loew,1862
 = *Heteromyza* Fallén,1820 (p.p.)
- O. buccata* (Fallén,1820)
 = *Heteromyza buccata* Fallén,1820
 * One of the main species in wrack-beds. Is this species possibly identical with *Heterochaeta b.* ?
 Sweden, S. England, Scotland
- Sensidae* "Svängflugor"
- Ornatinae*
- Orygma* Meigen,1830
- O. luctuosa* Meigen,1830
 * One of the main species in wrack-beds (in moist wrack).
 W. Sweden, Norway, Finland, Denmark, Germany, Netherlands, Belgium, France, S. England, Scotland
- Drosophiloidae
- Drosophilidae
- Drosophila* Fallén,1823
- D. funebris* (J.C. Fabricius,1787) "Stora attikflugan"
 * In banks of decaying seaweed.
 Sweden, Scotland
- D. phalerata* Meigen,18??
 S. England
- Enhydriidae "Vattenflugor"
- Hecamede* Haliday, in Curtis,1838
- H. albicans* (Meigen,1830)
 Sweden, Denmark, Balticum, Germany, Netherlands, Belgium, France, S. England
- Clanoneurum* Becker,1903
- C. cimiciformae* (Haliday,18???)
 Germany, Belgium, France, England, Mediterranean
- Atissa* Haliday, in Curtis,1837
- A. limosina* Becker,1???
 Norway, Finland, France, Br. Isles
- A. pygmaea* (Haliday,1833)
 ?Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Glenanthe* Haliday,1839
- G. fuscinervis* Becker,1???
 Norway (Molde), Finland, Germany
- G. ripicola* Haliday,1839
 Finland, Germany, Belgium, France, Br. Isles
- Philygria* Stenhammar,1844
- P. ochracea* Oldenberg,1???
 Borkum
- Ephydria* Fallén,1810 "Saltflugor"
- E. riparia* Fallén,1813
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- E. salinae* Zetterstedt,1846
 Sweden (so far only in inland salt water)
- Scatella* Robineau-Desvoidy,1830
- S. stagnalis* Fallén,18??
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- S. subguttata* (Meigen,1830)
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Halmopota* Haliday,1856
- H. salinarum* Behé,1???
 Germany, Br. Isles
- Lamproscatella* Hendel,1917
- L. quadrisetosa* (Becker,1896)
 European littoral
- Parydroptera* Collin,1913
- P. discomyzina* ?,1???
 * In salt-marshes.
 Britain
- Chloropoidea
- Tethinidae
- Pelomyia* Williston,1893
- P. cinerella* (Haliday,18???)
- Finland, ?Denmark, Germany, Netherlands, France, Br. Isles
- P. angustifacies* de Meij,1???
 Germany, Netherlands, Br. Isles
- Tethnia* Haliday, in Curtis,1836
 = *Tethina* Haliday,1838
- T. cinerea* Loew,18???
 ?Denmark, Germany, Netherlands, France, Br. Isles
- T. grisea* (Fallén,1810?)
 Sweden, Finland, Germany, Netherlands, Br. Isles
- T. illota* (Haliday,1836?)
 Finland, ?Denmark, Amrum, Netherlands, France, Br. Isles
- T. griseola* v. d. Wulp,18???
 Netherlands, Belgium, France, Br. Isles
- T. pictipes* Becker,1???
 Br. Isles
- T. nigripes* Cz.,1???
 Germany
- T. longirostris* Loew,18???
 France
- Canaceidae "Surfflugor"
- Canace* Haliday, in Curtis,1838
- C. nasica* Haliday,18???
 * Breeding in *Enteromorpha*
 Belgium, France, Br. Isles
- Xanthocanace* Hendel,1914
- X. ranula* Loew,18???
 ?Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Chloropidae "Fritflugor"
- Calamoncosis* Enderlein,1911
- C. tomentosa* (Macquart,1???)
 S. England
- Muscoidea
- Scathophagidae
 = *Cordylundiae* "Kolvflugor"
- Cordylurinae
- Scopeuma* Meigen,1800
 = *Scathophaga* Meigen,1803
- S. litoreum* Fallén,1823?
 * In moist wrack. ?Identical with *Scatomyza l.* ?
 Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- S. decipiens* (Haliday, in J. Curtis,1832)
 Germany, Netherlands, Br. Isles, Ireland, N. France
- S. islandicum* Becker,18???
 Norway, Iceland, N. France
- S. villipes* (Zetterstedt,1846)
 * Under dead seaweed.
 Sweden, Norway, S. England, Scotland
- S. rufiventre* Villeneuve,19???
 France
- S. calida* ?,1???
 * In moist wrack.
 Britain
- Ceratinostoma* Meade,1885
- C. ostiorum* (Haliday, in Curtis,1832)
 * In moist wrack.
 Sweden, Norway, ?Denmark, Germany, Netherlands, Belgium, France, S. England
- Calliphoridae "Spyflugor"
- Sarcophaga* Meigen,1826
- S. (Belliaria* Robineau-Desvoidy,1863) *melanura* Meigen,1826
 S. England
- Muscidae "Egentliga flugor"
- Mydaeinae
- Limnophorini
- Limnophora* Robineau-Desvoidy,1830
- L. (Neolimnophora* Schnabl,1902) *maritima* (v. Röder,1887)
 ?Denmark, German North Sea coast, France, Br. Isles
- L. (N.) virgo* Villeneuve,1906
 German North Sea coast, France, Br. Isles
- L. (Spilogona* Schnabl,1911) *baltica* Ringdahl,1918
 Sweden, Finland, ?Denmark, Germany, Netherlands, Br. Isles
- L. (S.) glauca* Stein,1916
 Sweden, Finland, Germany, Belgium
- L. (S.) aerea* (Fallén,1825)
 Sweden, Norway, Finland, Denmark, Germany, Netherlands, France, S. England
- L. (S.) biseriata* Stein,1916
 Sweden, ?Denmark, Borkum, Netherlands, France, Br. Isles
- L. (S.) marina* Collin,18???
 ?Denmark, Germany, Netherlands, France, S. England

- L. (Villeneuvia) Schnabl,1911) aestuum* Villeneuve,1902
?Denmark, German North Sea coast, Belgium, France, Br. Isles
- L. (Limnophora) setinervia* Schnabl,1911
= *L. excisra* Stein,1916
France, S. England
- L. insularis* Collin,18??
Norway, Br. Isles
- Lispe* Latreille,1796
= *Lispa* Fallén,1810
- L. caesia* Meigen,1826
Sweden, Balticum, ?Denmark, German North Sea coast, Netherlands, Belgium, France, Br. Isles
- L. hydromyzina* Fallén,1825
Sweden, Finland, Balticum, ?Denmark, German coasts, Netherlands, Br. Isles
- L. litorea* Fallén,1825
Sweden, Finland, Balticum, Denmark, German North Sea coast, Netherlands, Belgium, France, Br. Isles
- L. toewi* Ringdahl,1922
Sweden, ?German North Sea coast, France, Br. Isles
- L. tentaculata* (C. De Geer,1776)
Britain
- Mydaeini
- Helina* Robineau-Desvoidy,1830
- H. protuberans* (Zetterstedt,1845)
Sweden, Norway, Finland, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- Coenosimae*
- Chelisiini
- Limnospila* Schnabl,1902
- L. albifrons* (Zetterstedt,1849)
Sweden, Denmark, Germany, France, Br. Isles
- Dexiopsis* Pokorný,1893
- D. lacustris* Karl,1???
Amrum, Br. Isles
- Coenosimi
- Coenosia* Meigen,1826
- C. salinarum* Stein,1897
Germany, Netherlands, France, Br. Isles
- Anthomyiidae*
- Fucellini
- Fucellia* Robineau-Desvoidy,1842
- F. maritima* (Haliday,1838)
* one of the main species in wrack-beds
Sweden, Norway, Finland, Balticum, Denmark, Germany, Netherlands, Belgium, France, Br. Isles
- F. griseola* (Fallén,1819)
Sweden, Finland, Balticum, Denmark, German Baltic coast,
- F. fucorum* (Fallén,1819)
Sweden, Norway, Finland, Denmark, German Baltic coast, S. England
- F. marina* ?1???
* Maybe just a lapsus for *F. maritima*?
Sweden, Denmark
- Families yet lacking undoubted systematic placement
- Heleomyzidae* "Myllflugor"
= *Helomyzidae*
- Neoleria* Malloch,1919
- N. maritima* Villeneuve,19??
N. France
- Sphaeroceridae "Hoppflugor"
= *Borbonidae*
- Thoracochaeta* Duda,1918
- T. zosterae* (Haliday,1833)
= *Leptocera* Olivier,1813 *zosterae* (Haliday,1833)
* One of the main species in wrack-beds (in moist wrack).
Bohuslän, Norway, ?Denmark, Germany, Netherlands, France, S. England, Scotland
- T. brachystoma* (Stenhammar,18???)
* One of the main species in wrack-beds (in moist wrack).
S. England
- Scatomyza* Fallén,1810
- S. litorea* Fallén,1810
Sweden, S. England
- Collinellula* Strand,1928
- C. limosa* (Fallén,1820?)
= *Limosina limosa* Fallén,1820?
Balticum, ?Denmark, Germany, France, S. England
- Coprophila* Duda,1918
- C. vagans* (Haliday,1833)
S. England
- COLEOPTERA** "Skalbaggar"
- * See also Keys, James H., 1918. A List of the Maritime, Sub-Maritime and Coast-frequenting Coleoptera of South Devon and South Cornwall, with especial reference to the Plymouth District. *J. mar. biol. Ass. U.K.*, 11 : 497-513
- ADEPHAGA**
- Carabidae "Jordlöpare"
* Around 50 British coastal species
- Cicindelinae* Leconte & Horn,1883 "Sandjägare"
- Cicindela* Linnaeus,1758
- C. maritima* Latreille,1822-24 "Strandsandjägaren"
= *C. hybrida* Linnaeus,1758 var. *maritima* Latreille,1822-24
* Halophilic
Skåne, Halland, Värmland, Dalarna, Medelpad-Norrboten, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- C. hybrida* Linnaeus,1758 "Brun sandjägare"
Skåne-Bohuslän, Denmark
- Carabinae* Horn,1881 & Harpalinae "Egentliga jordlöpare"
- Broscus* Panzer,1813
- B. cephalotes* (Linnaeus, 1758) "Läderbagge"
* Coastal. In maritime sand dunes.
Skåne-Gästrikland, Denmark, Germany, S. England
- Dyschirius* Panzer,1813, ex Bonnelli
- D. obscurus* (Gyllenhal,1827)
* Halophilic
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, ?Br. Isles
- D. thoracicus* (P. Rossi,1790)
= *D. arenosus* Stephens,1827
* On sand
Sweden, Denmark, Finland, German North Sea coast
- D. impunctipennis* Daws.,1854
* Halobiont.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- D. extensus* Putzeys,1846
* Halobiont.
England, Germany
- D. chalcurus* Erichson,1837
* Halobiont.
Sweden, Denmark, Germany, Netherlands, Belgium
- D. salinus* Schaum.,1843
* Halobiont.
Sweden, Norway, Denmark, Balticum, Germany, Netherlands, Belgium, Br. Isles, Arcachon
- D. globosus* Herbst,1???
* Eurytopic with preference for sandy shores.
Bohuslän-Uppland, Åland, Finland
- Bembidion* Latreille,1810
= *Bembidium* Gyllenhal,1810
- B. (Actedium* Motschoulsky,1864) *pallidipenne* (Illiger,1802)
* Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- B. (Periphyphus* Stephens,1828) *saxatile* Gyllenhal,1827
* Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, Netherlands, Br. Isles
- B. (Periphyphus* Stephens,1828) *femoratum* (Sturm,1825)
= *B. maritimum* Stephens,18???
* Coastal.
S. England, N Germany
- B. (Notaphus* Stephens,1828) *varium* (Olivier,1795)
* Sub-maritime
Sweden, Denmark, Germany
- B. (Emphanes* Motschoulsky,1864) *normannum* (Dejean,1821)
* Halobiont.
Denmark, Germany, Netherlands, Belgium, S. England
- B. (Ocypus* Stephens,1828) *harpaloides* (Serville,1821)
S. England, Germany
- B. (Notaphemphanes* Netolitzky,1920) *ephippium* (Marsham,1802)
* Halobiont.
Denmark, Germany, Netherlands, Belgium, England
- B. (Talanea* Motschoulsky,1864) *aspericollis* (Germar,1829)
* Halobiont.
Germany, Belgium, France
- B. (Philochthus* Stephens,1828) *aeneum* (Germar,1824)
* Halobiont.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Br. Isles
- B. (Diplocampa* Bedel,1896) *fumigatum* (Creutzer, in Duftschmid,1812)
* Halobiont.
Sweden, Denmark, Germany, Netherlands, Belgium, England
- B. concinnum* (Stephens,18???)
* Halobiont.
Germany, Netherlands, Belgium, Br. Isles
- B. iricolor* Bedel,18???
* Halobiont.
Netherlands, Belgium, France, England
- B. assimile* Gyllenhal,18???
* Hygrophilous. Especially abundant on sea shores.
Bohuslän-Skåne, Finland
- Clivina* Latreille,1802
- C. fossor* (Linnaeus,1758)
Bohuslän, Finland
- Trechus* Schellenberg,1806
- T. quadristratus* Schrank,1???
* Eurytopic on heath & shore.
Bohuslän, Finland

- Cillenus* Samouelle,1819, ex Leach MS
- C. lateralis* Samouelle,1819
* An amphipod hunter; e.g. *Corophium & Talitrus*. Halobiont. Fully intertidal. Denmark, German North Sea coast, Netherlands, Br. Isles, Arcachon
- Aepus* Samouelle,1819, ex Leach MS
- A. marinus* (Ström,1783)
* Maritime. Halobiont. Lives permanently between the tidemarks in crevices & beneath stones. S. Norway, Denmark, S. England, Scotland
- Dicheirotrichus* J. Du Val,1855
- D. gustavii* Crotch,18??
= *Carabus pubescens* G. von Paykull,1798, non O.F. Müller,1776
* Sub-maritime. Halobiont. Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Br. Isles
- D. obsoletus* (Dejean,1829)
* Halobiont. Germany, Netherlands, Belgium, S. England
- Aepopsis* Jeannel,1922
- A. robinii* (Laboulbène,1850)
* Maritime. Halobiont. Lives permanently between the tidemarks in crevices & beneath stones. Br. Isles, France, Spain, Morocco
- Lymanaeum* Stephens,1828
- L. nigropiceum* (Marsham,1802)
* At marine shores beneath stones. Germany, Netherlands, S. England
- Tachys* Melsheimer,1806
? 3 British species at marine shores beneath stones.
- T. (Tachys) scutellaris* Germar,1829
= *T. taurica* V. v. Motschulsky,1850
* Halobiont. German North Sea coast, Netherlands, Belgium, France, England
- T. (Tachysura) V. v. Motschulsky,1862) parvulus* (Dejean,1831)
* Sub-maritime. S. England, Germany
- Trogophloeus* Mannerheim,1831
- T. schneideri* Ganglbauer,18??
* Halobiont. Denmark, Borkum, Netherlands
- T. halophilus* von Kiesenwetter,1844
* Halobiont. Sweden, Denmark, Germany, Netherlands, France, England
- Perileptus* Schaum,1860
- P. areolatus* (Creutz,1799)
* Coastal. Sweden, S. England, Germany
- Pogonus* Dejean,1821, ex Ziegler MS
- P. luridipennis* (Germar,1822-25)
* Under pieces of algae cast ashore on salt marsh shores. Halobiont. Bohuslän, Denmark, Germany, Netherlands, France, ?Scotland
- P. chalceus* (Marsham,1802)
* Sub-maritime. Halobiont. Denmark, Germany, Netherlands, Belgium, S. England, Scotland, Arcachon
- P. luridipennis* E.A. Nicolai,1822
* Halobiont. Germany
- P. litoralis* (Megerle, in Duftschmid,1812)
* Halobiont. Netherlands, Br. Isles
- Panagaeus* Latreille,1802
- P. bipustulatus* (J.C. Fabricius,1775) "Mindre korslöpare"
* Coastal. Skåne-Östergöt., Öland, Gotland, Nericke, S. England
- Chlaenius* Dejean,1826, ex Bonelli
- C. vestitus* (G. von Paykull,1790)
* Coastal. S. England
- Harpalus* Latreille,1802
- H. tenebrosus* (Dejean,1821)
* Coastal. S. England
- H. attenuatus* Stephens,1828
* Coastal. S. England
- H. tardus* (Panzer,1796)
Sweden, S. England
- Amara* Stephens,1827, ex Bonelli,1810
- A. ovata* (J.C. Fabricius,1792)
* Coastal. On gravel. Sweden, S. England
- A. incida* (Andersch, in Duftschmid,1812)
* Coastal. S. England
- A. convexiuscula* (Marsham,1802)
* Halophilic. Sweden, Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles
- Agonum* Stephens,1827, ex Bonelli,1810
- A. ruficornue* (Goeze,1777)
= *Platynus* Stephens,1827, ex Bonelli,1810 (*Anchromerus* Stephens,1827, ex Bonelli,1810) *ruficornis* Bohuslän, S. England (Goeze,1777)
- Microlestes* Schmidt-Goebel,1846
- M. maurus* (Sturm,1827)
* Coastal. S. England, Germany
- Lionychus* Wissmann, in Schmidt-Goebel,1846
- L. quadrillum* (Cretzter, in K. Duftschmidt,1812)
* Sub-maritime. Sweden, S. England
- Anisodactylus* Dejean,1829
- A. pocilloides* (Stephens,1828)
* Halobiont. Sweden, Denmark, Germany, Netherlands, Belgium, England
- Acupalpus* Latreille, in Cuvier,1829
- A. (Manicellus) V. v. Motschulsky,1864) elegans* (Dejean,1829)
* Halobiont. Denmark, Balticum, Germany, Netherlands, Belgium, England
- Nebria* Latreille,1802
- N. complanata* (Linnaeus,1767)
* Sandy shores Britain, Spain
- N. (Nebria) livida* (Linnaeus,1758) var. *lateralis* ?,???, German Baltic & North Sea coast
- Haliphilidae* Aube,1836 "Vattentrampare"
- Haliphus* Latreille,1802 (TM *Dytiscus impressus* Fabricius,1787 = *Haliphus (Haliphus) ruficollis* (De Geer,1774))
Subgen.: *Haliphus* (s. str.)
- H. lineolatus* Mannerheim,1844
* A typical brackish water species. Limnohalin. N. Baltic
- H. immaculatus* Gerhardt,1877
* A typical brackish water species. Pseudohalobiont. Baltic, Norway
- H. apicalis* Thomson,1868
* A typical brackish water species. Pseudohalobiont. Baltic, Britain
- H. fluvialis* Aubé,1836, in Dejean & Boisduval German Baltic coast
Subgen.: *Haliplidius* Guignot,1928
- H. obliquus* (J.C. Fabricius,1787)
* A typical brackish water species. Pseudohalobiont. Baltic
- H. confinis* Stephens,1829 *pallens* Fowler,1887
* A typical brackish water species. Pseudohalobiont. Baltic
Subgen.: *Liaphthus* Guignot,1928
- H. fulvulus* (J.C. Fabricius,1801) *lapponum* Thomson,1854
- H. flavicollis* Sturm,1834
* A typical brackish water species. Pseudohalobiont. Baltic, Norway
- Dytiscidae* De Geer,1774 "Dykarbagger"
- Hygrotes* Stephens,1828
- H. (Coelambus) Thomson,1860) inaequalis* (J.C. Fabricius,1777)
* A typical brackish water species. Limnohaline. * Coastal. Baltic, S. England
- Agabus* Leach,1817
- A. conspersus* (Marsham,1802)
* Coastal. A typical brackish water species. Pseudohalobiont. Baltic, S. England
- A. nebulosus* Forster,1???
* A typical brackish water species. Pseudohalobiont. Baltic, Norway
- A. undulatus* Schrank,1???
German Baltic coast
- Ilybius* Erichson,1832
- I. fuliginosus* (J.C. Fabricius,1792)
* A typical brackish water species. Limnohaline. Baltic
- I. subaeneus* Erichson,1837
* A typical brackish water species. altic
- Noterus* Schellenberg,1806
- N. clavicornis* (C. De Geer,1774)
* A typical brackish water species. Baltic, Skåne-Bohuslän, SE. Norway

- Laccophilus* Leach,1817
- L. variegatus* Germar,1???
- * A typical brackish water species
Baltic, S.E. Norway
- Bidessus* Sharp,1881
- B. geminus* (J.C. Fabricius,1798)
- * A typical brackish water species. Limnohaline.
Baltic, Norway
- Coelambus* Thomson,1860
- C. impressopunctatus* Schaller,18??
- * A typical brackish water species. Limnohaline
Baltic
- C. parallelogrammus* (Ahrens,1812)
- * A typical brackish water species. Pseudohalobiont Halophilic.
Bohuslän, Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles
- Deronectes* Sharp,1881
- D. depressus* (J.C. Fabricius,1794) *latescens* Falkenström,1???
- * A typical brackish water species. Pseudohalobiont.
Baltic, North Sea coasts
- Platambus* Thomson,1860
- P. maculatus* (Linnaeus,1758)
- Baltic, Skåne-Torne Lappmark
- POLYPHAGA** Emery,1886
- Staphylinidae** Latreille,1802
- Hydraenidae** Mulsant,1844
- Ochthebiinae** Thomson,1859
- Ochthebius* Leach,1815
- O. dilatatus* Stephens,1829, ex Leach MS
- * Coastal.
S. England
- O. metalleascens* Rosenhauer,1847
- * Coastal.
S. England
- O. marinus* (G. von Paykull,1798?)
- * A typical brackish water species. Sub-maritime. Halophilic-Halobiont.
Bohuslän, S. Norway, Finland, Denmark, Balticum, North Sea Shores, France, Br. Isles
- O. viridis* Peyron,1???
- * Coastal. Halophilic.
S. England, France, Germany
- O. lejolisi* Mulsant & Rey,18??
- * Sub-maritime.
S. England
- O. auriculatus* Rey,18??
- * Halobiont. Should the species name be *auricularius*?
Denmark, Germany, Netherlands, France, England
- O. impressicollis* Dejean,1821
- * Halophilic.
Germany, Netherlands, France, Br. Isles
- O. punctatus* Stephens,1829
- * Halophilic.
Netherlands, France, England
- Leiodidae**
- Leiodes* Latreille,1796
- L. ciliaris* (W.L.E. Schmidt, in Germar,1841)
- * Halophilic.
Sweden, Denmark, Finland, Germany, Netherlands
- L. dubia* (Kugelann,1794?)
- * Sandy coastal habitats.
Britain
- Hydnobius* Schmidt, in Germar,1841
- H. perrisi* (Fairmaire,18???)
- * Sandy coastal habitats.
Britain
- Silphidae** "Skinnarbaggar", "Asbaggar"
- Ablattaria** Reitter,1885
- A. laevigata* (J.C. Fabricius,1775)
- * Coastal.
S. England
- Ptiliidae** Erichson 1845/Motschulsky 1845 "Fjädervingar"
- Ptenidiom** Erichson,1845
- P. punctatum* (Gyllenhaal,1827)
- * Sub-maritime. Halobiont. In deep wrack layers.
Bohuslän-Uppland, Åland, Finland, Norway, Denmark, Balticum, Germany, Netherlands, Belgium, S. England
- Acrotrichis* Motschulsky,1848
- A. fratercula* Matthews,18???
- * In deep wrack layers.
Bohuslän-Uppland, Åland, Finland
- Actidiom* Matthews,1868
- A. coarctatum* (Haliday,1839?)
- * Halobiont. Sandy shores.
Skåne, Denmark, Germany, Netherlands, France, England
- Actinopteryx** Mathews,1872
- A. fucicola* (Allibert,1844)
- * Halobiont. Beneath seaweed.
Skåne, ?Norway, Denmark, Netherlands, England
- Staphylinidae** Latreille,1802 "Kortvingar"
- Aleocharinae**
- Actocharis* Fauvel,1870
- A. marina* Fauvel,1870
- = *A. readingi* Shp.,1???
- * Maritime.
S. England
- Phytosus* Curtis,1838, ex Rudd MS
- P. balticus* Kraatz,1???
- * Sub-maritime. Halobiont.
Sweden, Denmark, Germany, Netherlands, Belgium, S. England, Scotland
- P. spinifer* G.T. Rudd, in J. Curtis,1838
- * Sub-maritime. Halobiont.
Denmark, German North Sea coast, Netherlands, France, S. England, Scotland
- P. nigritrinitatis* (Chevrolat,1843)
- * Sub-maritime. Halobiont.
Germany, Netherlands, Belgium, France, S. England
- Arena* Fauvel,1862
- A. octavia* Fauvel,1862?
- * Halobiont.
Netherlands, S. England
- Athetida* Thomson,1858
- A. thinobioides* (Kraatz,1???)
- * Coastal.
S. England
- A. (Aloconota* Thomson,1858) *sulcifrons* (Stephens,1829)
- S. England
- A. (Pseudopasilia* Ganglbauer,1895) *tabida* (Kiesenwetter,18???)
- * Sub-maritime. Halobiont.
Germany, S. England, Scotland
- A. (Halobrecta* Thomson,1858) *flavipes* Thomson,18???
- * Sub-maritime. Halobiont.
Sweden, Germany, Netherlands, France, S. England, Scotland
- A. (Halobrecta* Thomson,1858) *puncticeps* Thomson,18???
- * Coastal. Halobiont.
Sweden, Norway, Germany, Netherlands, S. England, Scotland
- A. (Microdota* Mulsant & Rey,1873) *indubia* (Sharp,18???)
- * Coastal.
S. England
- A. triangulum* (Kraatz,1???)
- * Coastal. Halophilic.
Sweden, German Baltic, Netherlands, France, Br. Isles
- A. setigera* Sharp,18???
- * Halophilic.
Sweden, Finland, Balticum, Germany, England
- A. marina* Rey,18???
- * Halobiont.
Sweden, Finland, Germany, Netherlands, France, England
- A. meridionalis* Rey,18???
- * Halobiont.
Sweden, Denmark, Germany, Netherlands, England
- A. varendorffiana* Bernhauer,18???
- * Halobiont.
German North Sea coast
- A. (Thinobaena* Thomson,1859) *vestita* (Gravenhorst,1???)
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Finland, Germany, Netherlands, S. England, Scotland
- A. (Acrotona* Thomson,1859) *fungi* (Knob, in Gravenhorst,1806) *orbata* (Erichson,1837)
- * Coastal.
S. England
- Heterota** Mulsant & Rey,1873
- H. plumbea* (Waterhouse,1???)
- * Sub-maritime, under seaweed. Halobiont.
Hallands Väderö, Germany, France, S. England
- Myrmecopora** Sauley,1865
- M. lohmanderi* Bernhauer,19??
- * Under wrack
Bohuslän
- M. brevipes* Butler,18???
- S. England
- M. sulcata* (Kiesenwetter,1???)
- * Halobiont.
S. England, France
- M. uvida* Erichson,18???
- * Halobiont.
Netherlands, Belgium, England
- Diglotta* Champion,1899
- = *Diglossa* Haliday,1837, non Wagler,1832
- * With 2 maritime Scandinavian species.
- D. meresa* (Haliday,1837)
- * Maritime. Halobiont.
Denmark, Germany, Netherlands, France, S. England, Scotland
- D. marina* (Haliday,18???)

- * on rocky & sandy shores
Britain
- D. submarina* Fairmaire,1???
- Arcachon
- Ocalea* Erichson,1837
- O. rivularis* Miller,1???
- S. England
- O. picata* Stephens,18??
- Uppland
- Drusilla* Samouelle,1819, Leach MS
- D. canaliculata* Dillwyn,18??
- Br. Isles
- Oxypoda* Mannerheim,1831
- O. tarda* Sharp,1???
- England
- O. exoleta* Erichson,1839
- * Coastal.
S. England
- Aleochara* Gravenhorst,1802
- A. algarum* Fauvel,18??
- * Sub-maritime. Halobiont.
W. Sweden, Norway, Denmark, Germany, Netherlands, Belgium, France, S. England
- A. grisea* Kraatz,1???
- * Sub-maritime. Halobiont.
Sweden, Norway, Finland, Germany, Netherlands, S. England, Scotland
- A. obscurula* Gravenhorst,1806
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Finland, Germany, Netherlands, Belgium, France, S. England, Scotland
- Omaliniae
- Omalium* Gravenhorst,1802
- O. rivulare* (G. von Paykull,1789)
- Sweden, Denmark
- O. laeviusculum* Gyllenhal,1827
- * Sub-maritime. Halophilic.
Bohuslän, Norway, Finland, Germany, Netherlands, S. England, Scotland
- O. riparium* Thomson,18???
- * Sub-maritime. Halobiont.
Skåne, Norway, Denmark, Finland, Germany, Netherlands, Belgium, France, S. England, Scotland
- O. rugulipenne* Rye,18???
- * Halobiont.
Netherlands, France, Br. Isles
- Micralymma* Westwood,1837
- M. marinum* (Strom,1783)
= *M. brevipenne* (Gyllenhal,1810)
- * Maritime. Halobiont.
Bohuslän, W. Norway, Denmark, German North Sea coast, Netherlands, Belgium, France, Br. Isles
- Oxvteliniae
- Cryptobium* Mannerheim,1831
- C. fracticornis* (G. von Paykull,1790)
- Sweden, Denmark
- Bledius* Samouelle,1819, Leach MS
- B. spectabilis* Kraatz,1???
- * Halobiont.
Norway, Denmark, Germany, Netherlands, Arcachon, S. England, Scotland
- B. arenarius* (G. von Paykull,1800)
- * Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Arcachon, Br. Isles
- B. tricornis* (Herbst, Fuessly,1784)
- * Halophilic.
Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, England
- B. taurus* (?Dejean,1821)
= *B. furcatus* (Olivier,1???)
- * Halobiont.
Halland, Bohuslän, Denmark, Germany, France, England
- B. diota* ?,18???
- Sweden, Denmark
- B. bicornis* (?L.H. v. Bloch, Becker,1799?)
- * Halobiont.
Denmark, Germany, Netherlands, England
- B. unicornis* (Dejean, Germar,1825?)
- * Halobiont.
Germany, Belgium, Arcachon, England
- Oxytelus* Gravenhorst,1802
- O. perrisi* Fauvel,18???
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Germany, Netherlands, Belgium, France, S. England, Scotland
- O. complanatus* Erichson,1839
- * Coastal.
S. England
- Proteininae
- Proteinus* Latreille,1796
- P. brachypterus* Fabricius,1???
- Sweden, Denmark
- Xantholiniae
- Hypongryus* Tottenham,1940
- * Should it be spelled *Hyponygrus* ?
- H. punctulatus* (G. von Paykull,1789)
- Denmark
- Onedinae
- Heterothops* Stephens,1829, Kirby MS
- H. binotatus* (Gravenhorst,1806?)
- * Sub-maritime. Halobiont.
Sweden, Norway, Denmark, Germany, Netherlands, France, S. England, Scotland
- Steninae
- Stenus* Latreille,1796
- S. atratus* Erichson,1839
- * Coastal.
Sweden, S. England
- S. nanus* Stephens,1829
- S. England
- Paederinae
- Medon* Stephens,1833
- M. brunneus* Erichson,18???
- Denmark
- M. ripicola* (Kraatz,1???)
- * Sub-maritime.
S. England
- M. pociferus* Peyron,18???
- * Sub-maritime. Halobiont.
S. England
- Astenus* Dejean,1833
- A. filiformis* (Latreille,18???)
- * Coastal.
S. England
- Lathrobium* Gravenhorst,1802
- L. angustatum* Boiduval & Lacordaire,18???
- S. England
- Tachyporinae
- Tachyporus* Gravenhorst,1802
- T. chrysometinus* (Linnaeus,1758)
- all of Sweden, S. England
- Staphylininae Latreille,1802
- Cafius* Stephens,1829, Leach MS
- C. xantholoma* (Hellwig, J.L.C. Gravenhorst,1806)
- * Sub-maritime. Halobiont.
Uppland-Skåne-Bohuslän, Norway, Denmark, Finland, Germany, S. North Sea coast, France, Br. Isles
- C. fucicola* (Curtis,1???)
- * Sub-maritime. Halobiont.
S. England, Scotland
- Philonthus* Stephens,1829
- P. salinus* von Kiesenwetter,1844
- * Halobiont.
Denmark, Germany, Netherlands
- Orthidus* Mulsant & Rey,1876
- O. cribratus* Erichson,1840
- Europe
- Xantholina* Serville,1825
- X. longiventris* Heer,1839
- S. England
- Remus* F. Holme,1838
- R. sericeus* F. Holme,1838
- * Under wrack. Halobiont.
Swedish westcoast, Norway, Denmark, Germany, Netherlands, S. England
- Quedius* Stephens,1829
- Q. tristis* (Gravenhorst,18???)
- S. England
- Q. umbrinus* Erichson,1839
- * Coastal.
Sweden, S. England
- Q. semiaeneus* (Stephens,1829)
- * Coastal.
S. England
- Staphylinus* Linnaeus,1758
- S. ater* Gravenhorst,1802
- * Coastal.
S. England
- S. globulifer* Geoffroy, A.F. de Fourcroy,1785
- S. England
- Ocypus* Samouelle,1819
- O. olens* Müller,1???
- Amrum
- ?Subfamilies?

- Olophrum** Erichson,1839
- O. piceum** (Gyllenhal,1810)
Sweden, S. England
- O. consimile** (Gyllenhal,1810)
* Halophilic.
Sweden, ?Norway, Germany
- Trogophloeus** Mannerheim,1831
- T. halophilus** H. v. Kiesenwetter,1844
* Maritime.
S. England
- T. unicolor** Sharp,1???
= *T. anglicanus* Sharp,1???
* Sub-maritime.
S. England
- Pselaphidae** "Klubbhornsbaggar"
- Brachygluta** Thomson,1859
- B. simplex** (Waterhouse,1???)
* Sub-maritime.
S. England
- B. helferi** Schm.-Göb,18??
* Halobiont.
Sweden, Norway, Denmark, Germany, Netherlands, England
- Reichenbachia** Leach,1826
- R. waterhousi** Rye,18??
* Halobiont.
England
- Hydrophiloidae** De Geer,1774
- Hydrophilidae** De Geer,1774 "Vattenbaggar", "Palpbaggar"
- Sphaeridiinae** Latreille,1802
- Cercyon** Leach,1817
- C. littoralis** (Gyllenhal,1808)
* Sub-maritime. Halobiont. In deep wrack layers.
Bohuslän-Uppland, Åland, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Britain
- C. depressus** Stephens,1829
* Sub-maritime. Halobiont. In deep wrack layers.
Bohuslän-Uppland, Finland, Norway, Denmark, Germany, Netherlands, S. England, Scotland
- C. marinus** Thomson,1???
* Littoral.
Bohuslän
- Hydrophilinae** De Geer,1774
- Hydrophilini** De Geer,1774
- Enochrus** Thomson,1859
- E. bicolor** (J.C. Fabricius,1792)
* A typical brackish water species. Halophilic-Halobiont.
Sweden, ?Norway, Finland, ?Denmark, Balticum, North Sea Shores, France, Br. Isles
- E. melanoccephalus** (Olivier,1792)
* A typical brackish water species. Pseudohalobiont.
Baltic- Bohuslän, S. Norway
- E. testaceus** (J.C. Fabricius,17??)
* A typical brackish water species. Limnohaline.
Baltic
- E. quadrripunctatus** (Herbst,1797) *fuscipennis* Thomson,18??
* A typical brackish water species. ?Pseudohalobiont.
Baltic, Norway
- Berosini Mulsant,1844
- Berosus** Leach,1817
- B. spinosus** Stephens,18??
* A typical brackish water species. Halobiont.
Sweden (Skåne-Gotland), S. Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, Br. Isles
- Oocyclini Hansen,1991
- Laccobius** Erichson,1837
- L. decoratus** (Gyllenhal,1810)
* A typical brackish water species. Halobiont.
Baltic
- L. minutus** (Linnaeus,1758)
* A typical brackish water species. Limnohaline.
Baltic
- Elaphorus** J.C. Fabricius,1775
= *Helophorus* Illiger,1801
- E. minutus** J.C. Fabricius,1775
* A typical brackish water species.
Baltic
- Anacaenini Hansen,1991
- Paracymus** Thomson,1867
- P. aeneus** Germar,18??
* Brackish water species. Halobiont.
Skåne, Germany, Netherlands, France, England
- Histeridae** "Stumpbaggar", "Pansarbaggar"
- Halacritus** Schmidt,1893
- H. punctum** (Aubé,1843)
- = *Acritus* Leconte,1853 *punctum* (Aubé,1843)
* Sub-maritime. Halobiont.
German North Sea coast, Netherlands, S. England
- Baeckmanniulus** Reichardt,1926
- B. maritimus** (Stephens,1829, ■ Kirby MS)
= *Pachylopus* Erichson,1834 *maritimus* (Stephens,1829, ■ Kirby MS)
* Sub-maritime. Halobiont.
Denmark, Germany, Netherlands, France, S. England, Scotland
- Scarabaeoidea**
- Scarabaeidae** "Bladhorningar"
* 7 coastal species occurs in Britain
- Aphodius** Illiger,1798
- A. nitidulus** (J.C. Fabricius,1792)
* Coastal.
S. England
- Psammobius** Heer,1841
- P. porcicollis** (Illiger,1803)
* Coastal.
S. England
- Aegialia** Latreille,1807
- A. arenaria** (J.C. Fabricius,1787)
* Sub-maritime. Halobiont.
Sweden, Norway, ?Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles
- Byrrhoidea** Latreille,1804
= **Dryopoidea**
- Heteroceridae** MacLeay,1825 "Strandgrävbaggar"
- Heterocerinae** MacLeay,1825
- Heterocerus** Fabricius,1792
- H. flexuosus** Stephens,1828
* Halobiont.
Sweden, Norway, Denmark, Balticum, Germany, Netherlands, Arcachon, Br. Isles
- H. (Littorinus) Gozis,1885) maritimus** Guérin-Menéville,1844
* Halobiont.
Netherlands, Arcachon
- H. obsoletus** Curtis,1828
* Halophilic.
Sweden, Denmark, Germany, Belgium, Br. Isles
- Elateroidea** Leach,1815
- Elateridae** Leach,1815
= *Dicromychidae*
= *Lissomidae*
= *Plastoceridae* "Knäppare"
- Agriotes** Eschscholtz, ■ Thon,1829
- A. sordidus** (Illiger,1807)
* Sandy coasts.
Britain
- Throscidae**
= *Trixagidae*
- Trixagus** Kugelann,1794
- T. elateroides** (Heer,1841)
* Salt-marshes.
Britain
- Cantharoidea**
- Cantharidae** "Flugbaggar"
- Cantharis** Linnaeus,1758
- C. rufa** Linnaeus,1758
* Beneath strandline algae
Sweden, Britain
- C. (Telephorus) Schaeffer,1766) darwiniana** Crotch,18??
* Halobiont.
Germany, Belgium, Scotland
- Paratinus** Abeille de Perrin,1891
- P. femoralis** (Erichson,18???)
* Halophilic.
Sweden, Denmark, Finland, Germany
- Bostrichoidea**
- Ptinidae** "Tjuvbaggar"
- Ptinus** Linnaeus,1767
- P. palliatus** (Perris,18???)
Britain
- Cleroidea** Latreille,1802
- Melyridae** Leach,1815
- Malachius** Fabricius,1775
- M. sardous** Erichson,1840
* Halobiont.
Germany, Netherlands
- M. vulneratus** (Abeille,18???)
* Salt-marshes
Britain
- Dolichosoma** Stephens,1830

<i>D. lineare</i> (Rossi,17??) * Salt-marshes Britain	<i>A. humilis</i> Germar,1825 * Halobiont. Sweden, Norway, Denmark, Germany, Netherlands, England
<u>Cucujoidea</u> = <u>Clavicornia</u>	<i>A. crotchi</i> Pic.,19?? * Halobiont. England, France
<u>Corylophidae</u> = <u>Orthoperidae</u>	<i>A. binaculatus</i> (Illiger,1801) * Halophilic. Sweden, Finland, Denmark, Balticum, Germany, Netherlands, France
<i>Corylophus</i> Leach, in Stephens,1833	<i>A. flavigipes</i> (Panzer,17??) * Halophilic. Sweden, Finland, Denmark, Balticum, Germany, Netherlands, France
<i>C. cassidoides</i> (Marsham,1802)? * "Only British species occurring on the coast" according to Hayward & Ryland (Eds.),1990. Synon. of next?	<i>A. antherinus</i> (Linnaeus,1761) * In wrack. Sweden
Britain	<u>Oedemeridae</u> "Blombaggar"
<i>C. sublaevipennis</i> Du Val,18?? * Coastal. S. England	<i>Nacerdes</i> Dejean,1834, ex Steven MS = <i>Nacerda</i> Faldermann,1857
<u>Cryptophagidae</u> "Fuktbaggar"	<i>N. melanura</i> (Linnaeus,1758)? * Coastal. Sweden, S. England
<i>Atomaria</i> Stephens,1829, ex Kirby MS	<u>Mordellidae</u> "Tombaggar"
<i>A. fuscipes</i> (Gyllenhal,1808) Sweden, Britain	<i>Mordellistena</i> A. Costa, in O. Costa,1854
<u>Nitidulidae</u> "Glansbaggar"	<i>M. parvula</i> (Gyllenhal,18??) <i>inaequalis</i> Mulsant,1???
<i>Meligethes</i> Stephens,1829, ex Kirby MS	* Coastal. Sweden
<i>M. exilis</i> Sturm,1845 * Coastal. S. England	<u>Chrysomeloidea</u> Latreille,1802
<u>Lathridiidae</u> "Mögelbaggar"	<u>Chrysomelidae</u> Latreille,1802 "Bladbaggar"
<i>Corticaria</i> Marsham,1802 * 4 coastal British species	<i>Macropoea</i> Samouelle,1819, ex Hoffmannsegg MS = <i>Haemona</i> Dahl,1823, ex Megerle MS
<i>C. crenulata</i> (Gyllenhal,18??) * Coastal. W. Sweden, S. England	<i>M. mutica</i> Fabricius,1??? * Lives on <i>Ruppia</i> , <i>Zostera</i> etc. A typical brackish (mesohaline) water species. Halobiont. Uppland-Bohuslän, Finland, Danish estuaries, Balticum, Germany, North Sea Shores, France, Br. Isles
<i>C. impressa</i> (Olivier,1790) * Coastal. Uppland, S. England	<i>M. pubipennis</i> Reuter,1??? * A typical brackish water species. Halobiont. Baltic
<u>Coccinellidae</u> "Nyckelpigor"	<i>Chrysolinidae</i> Motschulsky, in Schrenck,1860
<i>Subcoccinella</i> Guérin-Méneville,1842, ex Huber	<i>C. haemoptera</i> (Linnaeus,1758) * on sandy coasts. S. England
<i>S. vigintiquatuorpunctata</i> (Linnaeus,1758) "Härig (24-prickig) nyckelpiga" * Coastal. Skåne-Gästrikland, S. England	<i>C. banksi</i> (J.C. Fabricius,1775) * Coastal. S. England
<i>Micraspis</i> Dejean,1835, ex Chevrolat MS	<u>Psylliodes</u> Berthold, in Latreille,1827
<i>M. sexdecimpunctata</i> (Linnaeus,1758) = <i>Typhaspis</i> Crotch,1874 <i>sexdecimpunctata</i> (Linnaeus,1758) * Coastal. S. England	<i>P. marcidia</i> (Illiger,1807) * Coastal, Halobiont. Sweden, Denmark, Finland, Balticum, Germany, Netherlands, Belgium, France, S. England
<i>Coccinella</i> Linnaeus,1758	<i>P. conannus</i> (Stephens,18???) * Salt-marshes. Britain
<i>C. undecimpunctata</i> Linnaeus,1758 * Halophilic. Sweden, Norway, Finland, Denmark, Balticum, Germany, North Sea Shores, France, Br. Isles	<i>Crepidodera</i> Dejean,1835, ex Chevrolat MS
<i>C. septempunctata</i> Linnaeus,1758 "7-prickig nyckelpiga", "Jungfru Maria Nyckelpiga" * Ubiquitous & often shore occurring Sweden, Britain	<i>C. impressa</i> (Fabricius,17???) * Salt-marshes.
<u>Phalacridae</u>	<i>C. nobilis</i> Linnaeus,1758 * Coastal. Sweden, S. England
<i>Phalacrus</i> G. von Paykull,1800	<i>C. virgata</i> C. de Villers,1789 * Salt-marshes. S. England
<i>P. brunnipes</i> (Brisout,18???) * Coastal & in banks of rivers Britain	<u>Curculionoidea</u> Latreille,1802
<u>Tenebrionoidea</u> Latreille,1802	<u>Curculionidae</u> Latreille,1802 "Vivlar"
<i>Tenebrionidae</i> Latreille,1802 "Svarbbaggar"	<i>Apion</i> Herbst, in Jablonsky,1797 "Spetsvivlar"
<i>Phaleria</i> Latreille,1802	<i>A. atomarium</i> Kirby,1808 * Coastal. S. England
<i>P. cadaverina</i> (J.C. Fabricius,1792) * Sub-maritime. Halobiont. Sweden, Denmark, Germany, Netherlands, Belgium, France, S. England	<i>A. confluens</i> Kirby,18??? * Coastal. S. England
<i>Melanimor</i> Steven,1829	<i>A. hookeri</i> Kirby,1808 * Coastal. S. England
<i>M. tibialis</i> (Fabricius,1781?) S. England	<i>A. schoenherri</i> Boheman,18??? * Coastal. S. England
<i>Opatrium</i> J.C. Fabricius,1775	<i>A. laevicollis</i> Kirby,1808 * Coastal. S. England
<i>O. sabulosum</i> (Linnaeus,1758) * Coastal. Skåne-Västmanland, S. England	<i>A. angustatus</i> Curtis,1838, ?non Dejean,1821 * Sub-maritime. Halobiont. S. England, France
<i>Phylan</i> Dejean,1821	<i>A. ononicola</i> Bach,18??? * Coastal. S. England
<i>P. gibbus</i> (J.C. Fabricius,1775) * Coastal. Halobiont. Sweden, Norway, Denmark, Finland, Germany, Netherlands, Belgium, France, S. England, Scotland	
<i>Cylindrinotus</i> Faldermann,1837	
<i>C. pallidus</i> (Curtis,18???) Britain	
<u>Anthicidae</u> Latreille,1802 "Snabbaggar"	
<i>Anthicus</i> G. von Paykull,1798	
<i>A. angustatus</i> Curtis,1838, ?non Dejean,1821 * Sub-maritime. Halobiont. S. England, France	

- A. limonii* Kirby,1808
 * Halobiont.
 France, ?Belgium
- Otiorhynchus* Germar,1824 "Öronvivlar"
- O. rugifrons* (Gyllenhal,1813)
 * Coastal.
 Sweden, S. England
- O. atroapterus* (C. de Geer,1775)
 * Coastal. Halobiont.
 Sweden, Denmark, Germany, Netherlands, Belgium, S. England, Scotland
- O. lignaeus* (Olivier,1807)
 * Halophilic.
 Sweden, Denmark, Germany
- Bagous* Germar,1817
- B. argillaceus* Gyllenhal, in Shoeherr,1836
 * Halophilic.
 Sweden, Denmark, Germany, England
- Cathormiocerus* Schoenherr,1843
- C. attaphilus* Brisout,18??
 * Coastal.
 S. England
- C. myrmecophilus* (Seidlitz,18???)
 * Coastal.
 S. England
- Polydrusus* Germar,1817 "Glansvivlar"
 = *Polydrosus* Shoeherr,1826
- P. chrysomela* (Olivier,1807)
 * Sub-maritime.
 S. England
- Philopedon* Stephens,1831, non Stephens,1829
 = *Cneorhinus* Shoeherr,1823
- P. plagiatus* (Schaller,18???)
 * Coastal.
 S. England
- Sitona* Germar,1817 "Ärtvivlar"
- S. waterhousei* J. Walton,1846, ex Shoeherr MS
 * Coastal.
 S. England
- Phloeophagia* Aurivillius,1924
- P. spadix* (Herbst, in Jablonsky,1795)
 * Sub-maritime.
 S. England
- Smicronyx* Schoenherr,1843
- S. jungermanniae* (Reich,1797)
 * Coastal.
 S. England
- Tychius* Germar,1817
- T. schneideri* (Herbst, in Jablonsky,1795)
 * Coastal.
 S. England
- Limnobaris* Bedel,1885
- L. talbum* (Linnaeus,1758)
 * Sub-maritime.
 S. England
- Ceuthorhynchidius* J. Du Val,1854
- C. dawsoni* Brisout,18??
 * Coastal, Halobiont.
 S. England, ?France, ?Belgium
- Ceutorhynchus* Germar,1824, ex Schueppel MS
 = *Ceutorhynchus* Gemminger & Harold,1871
- C. terminatus* (Herbst, in Jablonsky,1795)
 * Coastal.
 S. England
- Sirocalus* Heyden, in Heyden, Reitter & Weise,1906
- S. caciis* Hansen,18??
 * Halobiont.
 Denmark, Germany, Netherlands
- Mecinus* Germar,1821
- M. circulatus* (Marsham,1802)
 * Coastal
 S. England