



Agreement on the Eutrophication Monitoring Programme

(Reference Number: 2005-4)¹

1. This agreement updates and supersedes the earlier OSPAR agreement on a nutrient monitoring programme (reference number: 1995-5) which was adopted in 1995.
2. The OSPAR Eutrophication Monitoring Programme is an integral part of the OSPAR Eutrophication Strategy. It provides the basis for enabling Contracting Parties to assess and classify the eutrophication status of their maritime waters under the “Comprehensive Procedure” of the Common Procedure for the Identification of the Eutrophication Status of the OSPAR Maritime Area (“Common Procedure”, reference number: 2005-3).
3. A satisfactory monitoring programme needs clear objectives and should comply with the guiding principles in the OSPAR Strategy for a Joint Assessment and Monitoring Programme (JAMP) (reference number: 2003-22, as amended).
4. Since the monitoring and assessment procedures associated with a eutrophication monitoring programme will address sub-regions of the maritime area with different levels of priority (i.e. problem areas, potential problem areas and non-problem areas with regard to eutrophication) and different characteristics (stratified/non-stratified water bodies, rocky/non-rocky shores etc.), a certain degree of flexibility should be incorporated into the design of the monitoring programme.
5. The minimum requirements for a eutrophication monitoring programme should therefore be as follows:
 - a. Non-problem areas with regard to eutrophication

In non-problem areas with regard to eutrophication the monitoring programme has the function of detecting changes in the eutrophication status or confirming the status of particular areas as non-problem areas. This should be done with respect to assessment parameters and their assessment levels related to the area-specific background concentrations defined in the Comprehensive Procedure. Clearly, monitoring effort should be limited to a limited number of parameters and a limited frequency of measurements, although spatial coverage should not be neglected.
 - b. Problem areas with regard to eutrophication

In problem areas with regard to eutrophication the monitoring programme should focus on long-term trends in nutrient concentration and on a selection of related eutrophication effect parameters, taking into account corresponding long-term trends in nutrient inputs. A larger number of parameters and a higher sampling frequency should be considered than is the case for non-problem areas, so as to satisfy statistical requirements. The spatial coverage should also be more focused than for non-problem areas. Monitoring should continue until the non-problem area status is achieved.
 - c. Potential problem areas with regard to eutrophication

With regard to their unknown status, potential problem areas with regard to eutrophication should be monitored in the same manner as problem areas, for a trial period not exceeding five

¹ Supersedes agreement 1995-5. Source: EUC 2005 Summary Record - EUC 05/13/1, Annex 6.



years. This should enable the area to be reclassified as either a problem area or a non-problem area with regard to eutrophication.

6. In implementing these minimum monitoring requirements Contracting Parties should focus on those eutrophication effects that are sufficiently closely linked to nutrient enrichment to be of value in indicating the eutrophication status of an area. The eutrophication effects should be selected on the basis of the assessment parameters listed in Table 1 of the Common Procedure.

7. If, following the classification of the maritime area in terms of eutrophication status, sub-regions (based for example on hydrographic characteristics) are identified within an area of particular eutrophication status (i.e. a problem area, a potential problem area or a non-problem area with regard to eutrophication), then the minimum monitoring requirements specified under § 5 should be applied to each of the sub-regions.

8. The spatial distribution of the monitoring stations should, prior to the establishment of the eutrophication status of the maritime area using the Common Procedure, be commensurate with the anticipated extent of eutrophication in the area under consideration as well as its hydrographic characteristics. Consequently, each Contracting Party should determine the optimum frequency per year and optimum locations for its monitoring stations. Where appropriate, efforts should be coordinated between Contracting Parties.

9. The minimum requirements for a eutrophication monitoring programme are specified in Tables 1 and 2. Contracting Parties should increase the scope and frequency of monitoring as they consider appropriate.

10. The Eutrophication Monitoring Programme forms part of the OSPAR Coordinated Environmental Monitoring Programme (the CEMP - reference number: 2005-5 (latest update)). Contracting Parties shall report the monitoring results for the parameters listed in Tables 1 and 2 in accordance with the arrangements for the CEMP agreed and updated periodically by OSPAR. ICES is currently the OSPAR data centre for marine environmental monitoring data, and according to the agreements of OSPAR, Contracting Parties are obliged to report their monitoring data to ICES by 1 September in the year following the year of monitoring using the agreed formats and should resolve any data processing issues with the ICES data centre.

11. The programmes and procedures for the assessment or classification by each Contracting Party of the eutrophication status of areas within its marine waters is set out in the OSPAR Eutrophication Strategy (reference number: 2003-21) and in the Comprehensive Procedure of the Common Procedure.




Table 1. Nutrient enrichment¹

	Non-problem areas	Potential problem areas	Problem areas
NH ₄ -N ^{2,4} (μmol l ⁻¹)	+	+	+
NO ₂ -N ^{2,4} (μmol l ⁻¹)	+	+	+
NO ₃ -N ^{2,4} (μmol l ⁻¹)	+	+	+
PO ₄ -P ^{3,4} (μmol l ⁻¹)	+	+	+
SiO ₄ -Si ⁴ (μmol l ⁻¹)	-	+	+
Salinity	+	+	+
Temperature	+	+	+
Frequency ⁵	About every three years during winter	Annually during winter when algal growth is at a minimum and during monitoring of direct and indirect effects	

+ Action required

- Action discretionary

1 All parameters should be monitored in conjunction with area-specific ecosystem features.

2 Winter dissolved inorganic nitrogen (DIN) is the sum of NH₄-N, NO₂-N and NO₃-N.

3 Winter dissolved inorganic phosphate (DIP)

4 Monitoring of winter DIN, DIP and Si should be in conjunction with salinity measurements (see Common Procedure, §§ 4.25 and 4.28).

5 Monitoring should include sufficient samples to confirm that the maximum winter nutrient concentration has been determined.

Table 2. Direct and indirect eutrophication effects¹

	Non-problem areas	Potential problem areas	Problem areas
Phytoplankton chlorophyll <i>a</i> (μg l ⁻¹)	-	+	+
Phytoplankton indicator species (cells l ⁻¹ ; species composition)	-	+ species composition: (genera and nuisance/potentially toxic species)	+ species composition: (genera and nuisance/potentially toxic species) + TOC and POC ²
Macrophytes, including macroalgae and angiosperms ³	-	+ biomass	+ biomass + species composition, coverage, and reduced depth distribution
O ₂ concentration (mg l ⁻¹ ; including % O ₂ saturation)	-	+	+
(zoo) Benthic communities	-	+ biomass and species composition (if time series already exist)	+ biomass, species composition and eutrophication indicator species
Frequency ⁴	-	annually during the algal growing season	

**OSPAR CONVENTION FOR THE PROTECTION OF THE MARINE
ENVIRONMENT OF THE NORTH-EAST ATLANTIC**



- + Action required
- Action discretionary
- 1 All parameters should be monitored in conjunction with area-specific ecosystem features.
- 2 TOC: Total Organic Carbon; POC: Particulate Organic Carbon.
- 3 In shallow areas, primarily in estuaries and coastal waters.
- 4 With adequate frequency and area coverage