

<i>cantabricus</i> Pardos, Higgins & Benito, 1998	A	Nanaloricida, Nanaloricidae) from the Mediterranean Sea. <i>Italian Journal of Zoology</i> 65: 219-226.
<i>capitatus</i> (Zelinka, 1928)	M	
<i>citrinus</i> Zelinka, 1928	M	
<i>druxi</i> d'Hondt, 1973	M	
<i>dujardini</i> Claparède, 1863	A	
<i>elongatus</i> Nyholm, 1947	A	
<i>ferrugineus</i> Zelinka, 1928	M	
<i>gerardi</i> Higgins, 1978	M	
<i>higginsi</i> Huys & Coomans, 1989	A	
<i>hispanicus</i> Pardos, Higgins & Benito, 1998	A	
<i>krishnaswamyi</i> Higgins, 1985	A	
<i>kristensenii</i> Higgins, 1985	A	
<i>levanderi</i> Karling, 1954	A	
<i>riedli</i> Higgins, 1966	M	
<i>setiger</i> Greef, 1869	A	
<i>worthingii</i> Southern, 1914	A	
Family Semnoderidae		
<i>Semnoderes</i>		
<i>armiger</i> Zelinka, 1928	A M	
<i>ponticus</i> Bacescu & Bacescu, 1956	M	
Family Zelinkaderidae		
<i>Zelinkaderes</i>		
<i>submersus</i> (Gerlach, 1969)	A	
Order Homalorhagida		
Family Neocentrophydidae		
<i>Paracentrophytes</i>		
<i>quadridentatus</i> (Zelinka, 1928)	A M	
Family Pycnophyidae		
<i>Kinorhynchus</i>		
<i>giganteus</i> (Zelinka, 1928)	A M	
<i>paraneapolitanus</i> Sheremetevskij, 1974	M	
<i>Pycnophyes</i>		
<i>calmani</i> Southern, 1914	A	
<i>carinatus</i> Zelinka, 1928	M	
<i>communis</i> Zelinka, 1908	A M	
<i>dentatus</i> (Reinhard, 1881)	A M	
<i>flaveolatus</i> Zelinka, 1928	A M	
<i>kielensis</i> Zelinka, 1928	A M	
<i>maximus</i> Reimer, 1963	A	
<i>ponticus</i> (Reinhard, 1881)	M	
<i>robustus</i> Zelinka, 1928	M	
<i>rugosus</i> Zelinka, 1928	M	
<i>zelinkaei</i> Southern, 1914	A	

Loricifera

Compiled by Jacob van der Land and Birger Neuhaus

The marine Loricifera are now known with 11 species from North America, Europe, and the Japanese deep-sea (Higgins & Kristensen 1986, Kristensen 1983, Todaro & Kristensen 1998). Only 2 species have been reported from Europe (Kristensen 1983, Todaro & Kristensen 1998). Loricifera inhabit various kinds of sediment from coarse sand or shelly gravel to red clay with silt and sand. They may be extracted from the substratum either by freshwater shock (Kristensen 1983) or by multiple decantations (Todaro & Kristensen 1998). It is certain that species new to science inhabit the area covered by the ERMS programme.

References

- Higgins, R. P. & R. M. Kristensen 1986: New Loricifera from Southeastern United States coastal waters. *Smithsonian Contributions to Zoology* 438: 1-70.
- Kristensen, P. M. 1983: Loricifera, a new phylum with Aschelminthes characters from the meiobenthos. *Zeitschrift für zoologische Systematik und Evolutionsforschung* 21: 163-180.
- Todaro, M. A. & R. M. Kristensen 1998: A new species and first report of the genus *Nanaloricus* (Loricifera,

Nanaloricida, Nanaloricidae) from the Mediterranean Sea. Italian Journal of Zoology 65: 219-226.

Class Loricifera

Family Nanaloricidae

<i>Nanaloricus</i>		
<i>mysticus</i> Kristensen, 1983	A	

khaitatus Todaro & Kristensen, 1998

M

Nematomorpha and Priapulida

Compiled by Jacob van der Land

The ERMS lists for Priapulida and Nematomorpha were based on Adrianov & Malakhov (1996) and on Malakhov & Adrianov (1995).

References

- Adrianov, A.V. & V.V. Malakhov (1996): Priapilida [sic] (Priapulida): structure, development, phylogeny, and classification. - KMK Scientific Press, Moscow, pp. 266. [in Russian]
- Malakhov, V.V. & A.V. Adrianov (1995): Cephalorhyncha - A new phylum of the animal kingdom. - KMK Scientific Press, Moscow, pp. 202. [in Russian]

Class Nematomorpha

Order Nectonematida

Family Nectonematidae

<i>Nectonema</i>		
<i>agile</i> Verril, 1879	A M	
<i>svensksundi</i> Bock, 1908	A	
<i>munidae</i> Brinkmann, 1930	A	

Class Priapulida

Family Chaetostephanidae

<i>Maccabeus</i>		
<i>tentaculatus</i> Por, 1973	M	

Family Priapulidae

<i>Halicryptus</i>		
<i>spinulosus</i> von Siebold, 1849	A	
<i>Priapulopsis</i>		
<i>bicaudatus</i> (Koren & Danielssen, 1868)	A	
<i>cnidephorus</i> Salvini-Plawen, 1973	M	

Priapulus

<i>abyssorum</i> Menzies, 1959	A M	
<i>caudatus</i> Lamarck, 1816	A	

Family Tubiluchidae

<i>Tubiluchus</i>		
<i>arcticus</i> Adrianov <i>et al.</i> , 1989	A	

NEMATODA

Nematoda (free-living)

Compiled by Guy De Smet, Magda Vincx, Ann Vanreusel, Sandra Vanhove, Jan Vanaverbeke and Maaike Steyaert

The list was compiled by Guy De Smet, a non-specialist, who was assisted by nematologists Magda Vincx, Ann Vanreusel, Sandra Vanhove, Jan Vanaverbeke and Maaike Steyaert. Dr. F. Riemann gave useful advice for the genus *Daptonema*. The classification in the list is based upon Lorenzen (1979, 1994). Information on the geographic distribution in the list is based upon Gerlach and Riemann (1973, 1974), Platt and Warwick (1983, 1988) and Warwick *et al.* (1998).

References

- Lorenzen, S. *Entwurf eines Phylogenetischen Systems der Freilebenden Nematoden*, Kiel, 1979.
 Lorenzen, S. *The Phylogenetic Systematics of Freeliving Nematodes*, The Ray Society, 1994.
 Gerlach, S. A. and Riemann, F. The Bremerhaven Checklist of Aquatic Nematodes. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven*, Supplement 4, (Heft 1 1973, Heft 2 1974).
 Platt, H. M. and Warwick, R. M. Free-living Marine Nematodes (Part I British Enoplids) *Synopsis of the British Fauna* (New series) No. 28, 1983.
 Platt, H. M. and Warwick, R. M. Free-living Marine Nematodes (Part II British Chromadorids) *Synopsis of the British Fauna* (New series) No. 38, 1988.
 Warwick, R. M., Platt, H. M. and Somerfield, P. J. Free-living Marine Nematodes (Part III Monhysterids) *Synopsis of the British Fauna* (New series) No. 53, 1998.

NEMATODA, FREE-LIVING

Acantholaimus

- setosus* Vitiello, 1970
ewensis Platt & Zhang, 1982
polydentatus Gerlach, 1951
longisetosus Allgén, 1933
- Acanthonchus*
gracilis (Ditlevsen, 1918) Wieser, 1955
viviparus Cobb, 1920
arcuatus (Kreis, 1928) Wieser, 1955
pugionatus Vitiello, 1970
- Acanthopharyngoides*
tyrrhenicus Wieser, 1954
- Acanthopharynx*
setosissima Schuurmans Stekhoven, 1943
micrampus Schuurmans Stekhoven, 1942
perarmata Marion, 1870
rigida Schuurmans Stekhoven, 1950
micans (Eberth, 1863) Marion, 1870
affinis Marion, 1870

Actinonema

- longicaudatum* (Steiner, 1918) Wieser, 1954
pachydermatum Cobb, 1920

- fidatum* Vitiello, 1970
celtica Boucher, 1976

Adeuchromadora

- megamphida* Boucher & Bovée, 1972

Adoncholaimus

- thalassophysas* (de Man, 1876) Filipjev, 1918
panicus Cobb, 1930

- derjugini* (Ssaweljev, 1912) Filipjev, 1924

- fuscus* (Bastian, 1865) Filipjev, 1918

- lepidus* (de Man, 1889) Filipjev, 1918

- islandicus* Kreis, 1963

Aegialoalaimus

- punctatus* (Allgén, 1929) Schuurmans Stekhoven, 1931
elegans de Man, 1907
tenuis Kreis, 1928
tenuicaudatus Allgén, 1932
- Alaimella*
cincta Cobb, 1920
- Amphidelus*
puccinelliae Lorenzen, 1966
- Amphimonhystera*
anechma (Southern, 1914) Allgén, 1929
helgolandica Riemann, 1967
paranechma Allgén, 1935
subtilis Lorenzen, 1972
- Anoplostoma*
elegans Kreis, 1929
blanchardi de Man, 1888
exceptum Schulz, 1935
viviparum (Bastian, 1865) Bütschli, 1874
- Anticoma*
strandii Allgén, 1940
stekhoveni Wieser, 1953
- acuminata* (Eberth, 1863) Bastian, 1865
pellucida Bastian, 1865
murmanica Filipjev, 1927
minor Filipjev, 1927
insulaealbae Filipjev, 1927
arctica Steiner, 1916
longicaudata Kreis, 1928
brevisetosa Platonova, 1967
brevicaudata Kreis, 1963
eberthi Bastian, 1865
- Anticomopsis*
typica Micoletzky, 1930
- Anticyathus*
primitivus (Allgén, 1933) Gerlach, 1963
- Anticyclus*
gigas Vitiello, 1969
exilis Cobb, 1920
pachyamphis (Wieser, 1956) Gerlach, 1963
- Antomicron*
pratense Lorenzen, 1966
profundum Vitiello, 1971
elegans (de Man, 1922) De Coninck, 1965
pellucidum Cobb, 1920
- Aponema*
torosa (Lorenzen, 1973) Jensen, 1978
- Araeolaimus*
paucisetosus (Wieser, 1951) Lorenzen, 1981
penelope Moore, 1977
septentrionalis Kreis, 1963
microphthalmus (de Man, 1893) Lorenzen, 1981
macrocirculus Kreis, 1928
longisetosus Wieser, 1954
filipjevi Schuurmans Stekhoven & Adam, 1931
demani (Schuurmans Stekhoven, 1950) Wieser, 1956
bioculatus (de Man, 1876) de Man, 1888
gajevskii Paramonov, 1929
mediterraneus (de Man, 1876) de Man, 1888
ponticus Filipjev, 1922
elegans de Man, 1888
longicauda Allgén, 1929
zosterae (Filipjev, 1918) Chitwood & Chitwood, 1937
tenuicaudatus Allgén, 1935
supralitoralis Wieser, 1960
steineri Filipjev, 1922
- Ascolaimus*
elongatus (Bütschli, 1874) Schuurmans Stekhoven, 1932
- Astomonema*
southwardorum Austen *et al.*, 1993
- Atrochromadora*
parva (de Man, 1893) Wieser, 1954
microlaima (de Man, 1889) Wieser, 1954
- Austranema*

- shirleyae* (Coles, 1965) Inglis, 1969
Axonolaimus
spinosus (Bütschli, 1874) de Man, 1889
helgolandicus Lorenzen, 1972
hexapilus Wieser & Hopper, 1967
paraspinosus Schuurmans Stekhoven & Adam, 1931
filiformis de Man, 1889
drachi Luc & De Coninck, 1959
arcuatus Schuurmans Stekhoven, 1950
orcombensis Warwick, 1970
schuurmansstekhoveni Allgén, 1935
limalis Ssaweljev, 1912
tirrenicus Brunetti, 1941
typicus de Man, 1922
setosus Filipjev, 1918
ponticus Filipjev, 1918
villosus Skwarra, 1922
demani De Coninck & Schuurmans Stekhoven, 1933
Barbonema
setiferum Filipjev, 1927
horridum Gerlach, 1956
Bathyepsilonema
pustulatum Gerlach, 1952
Bathylaimus
parafilicaudatus Allgén, 1935
arcticus Kreis, 1963
septentrionalis (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
denticaudatus Allgén, 1930
zostericola (Allgén, 1933) De Coninck & Schuurmans Stekhoven, 1933
macramphis (Schuurmans Stekhoven & De Coninck, 1933) De Coninck & Schuurmans Stekhoven, 1933
longisetosus (Allgén, 1929) De Coninck & Schuurmans Stekhoven, 1933
filicaudatus (Schuurmans Stekhoven & Adam, 1931) De Coninck & Schuurmans Stekhoven, 1933
dentatus Kreis, 1963
capacosus Hopper, 1962
cobbi Filipjev, 1922
australis Cobb, 1894
inermis (Ditlevsen, 1918) de Man, 1922
paralongisetosus Schuurmans Stekhoven & De Coninck, 1933
stenolaimus Schuurmans Stekhoven & De Coninck, 1933
tenuicaudatus (Allgén, 1933) Allgén, 1947
Bellbolla
teissieri (Luc & De Coninck, 1959) Andrassy, 1973
galanachmorae (Inglis, 1961) Andrassy, 1973
Boreomicrolaimoides
problematicus (Ditlevsen, 1918) Allgén, 1954
Bradybucca
rhopalura (Schuurmans Stekhoven, 1950) Schuurmans Stekhoven, 1956
Calligyrus
gerlachi Lorenzen, 1969
Calyptronema
maxweberi (de Man, 1922) Wieser, 1953
klatti (Allgén, 1953) Wieser, 1953
setifer (Gerlach, 1953) Wieser, 1953
acuminatum (Eberth, 1863) Wieser, 1953
longisetosum (Schuurmans Stekhoven, 1943) Wieser, 1953
eberti (de Man, 1876) Schuurmans Stekhoven, 1950
sabulicola (Filipjev, 1918) Kreis, 1926
Camacolaimoides
praedator (de Man, 1922) De Coninck & Schuurmans Stekhoven, 1933
Camacolaimus
barbatus Warwick, 1970
zostericola (Filipjev, 1918) Allgén, 1932
tardus de Man, 1889
longicauda de Man, 1922
propinquus Allgén, 1929
prytherchi Chitwood, 1935
glauxicola Allgén, 1951
dolichocercus Filipjev, 1922
bathycola Filipjev, 1922
Campylaimus
cylindricus Gerlach, 1956
gerlachi Timm, 1961
mirus Gerlach, 1956
inaequalis Cobb, 1920
lefeverei Gerlach, 1956
Ceramonema
undulatum De Coninck, 1942
filipjevi De Coninck, 1942
pisanum Gerlach, 1953
chitwoodi De Coninck, 1942
reticulatum Chitwood, 1936
yunfengi Platt & Zhang, 1982
salsicum Gerlach, 1956
Cervonema
tenuicaudatum (Wieser, 1954) Schuurmans Stekhoven, 1950
Chaetonema
vicinum Gerlach, 1954
longisetum (Steiner, 1916) Filipjev, 1927
riemanni Platt, 1973
Cheironchus
vorax Cobb, 1917
Chitwoodia
tripapillata Jayasree, 1976
warwicki Jayasree, 1976
menora Gerlach, 1956
falcata Gerlach, 1956
Chloronemella
filiformis Allgén, 1929
Choanolaimus
psammophilus de Man, 1880
Choniolaimus
papillatus Ditlevsen, 1918
panicus Gerlach, 1956
novempapillatus Schuurmans Stekhoven, 1946
crassus Vitiello, 1971
Chromadora
brevipapillata Micoletzky, 1924
macrolaima de Man, 1889
nudicapitata Bastian, 1865
caeca Bastian, 1865
papillata Bastian, 1865
sabelloides Bastian, 1865
bipapillata Micoletzky, 1924
buesumensis Kreis, 1924
kreisi Schuurmans Stekhoven & Adam, 1931
undecimpapillata Wieser, 1959
axi Gerlach, 1951
Chromadorella
membranata (Micoletzky, 1922) Wieser & Hopper, 1967
duopapillata Platt, 1973
acriminiensis (Gerlach, 1953) Lorenzen, 1972
parapoeilosoma (Micoletzky, 1922) Wieser & Hopper, 1967
filiformis (Bastian, 1865) Filipjev, 1918
Chromadorina
inversa Wieser, 1955
germanica (Bütschli, 1874) Wieser, 1954
armata (Allgén, 1933) Wieser, 1954
bergensis (Allgén, 1932) Wieser, 1954
cervix (Wieser, 1951) Wieser, 1954
bioculata (Carus, 1857) Wieser, 1954
demani Inglis, 1962
erythrophthalma (Schneider, 1906) Wieser, 1954
incurvata Wieser, 1956
laeta (de Man, 1876) Micoletzky, 1924
longisetosa (De Coninck & Schuurmans Stekhoven, 1933) Wieser, 1954
ocellata (Paramonov, 1929) Wieser, 1954

- viridis* (Linstow, 1876) Wieser, 1954
supralitoralis Lorenzen, 1969
granulopigmentata (Wieser, 1951) Wieser, 1954
rognoensis (Allgén, 1932) Wieser, 1954
macropunctata (Wieser, 1954) Wieser, 1954
obtusa Filipjev, 1918
nuda Wieser, 1954
micoletzkyi Inglis, 1962
majae Wieser, 1968
Chromadorissa
strandii Kreis, 1928
Chromadorita
tenuis (Schneider, 1906) Filipjev, 1922
obliqua (Gerlach, 1953) Wieser, 1954
leuckarti (de Man, 1876) Filipjev, 1929
norvegica (Allgén, 1946) Wieser, 1954
minor (Allgén, 1927) Wieser, 1954
gracilis (Filipjev, 1922) Wieser, 1954
guidoschneideri (Filipjev, 1929) Wieser, 1954
nana Lorenzen, 1973
brevisetosa Gerlach, 1953
minima (Kreis, 1929) Wieser, 1954
hyaloccephala (Steiner, 1916) Filipjev, 1922
murcodonta (Steiner, 1916) Wieser, 1954
abnormis (Kreis, 1928) Wieser, 1954
pachydemata (Schneider, 1926) Wieser, 1954
demaniana Filipjev, 1922
heterophya (Steiner, 1916) Filipjev, 1922
brachypharynx (Allgén, 1932) Wieser, 1954
Chromaspirina
lunatica Gerlach, 1965
paucispira Schuurmans Stekhoven, 1950
denticulata (Gerlach, 1953) Luc & De Coninck, 1959
multipapillata Jayasree & Warwick, 1977
thieryi De Coninck, 1943
parapontica Luc & De Coninck, 1959
pellita Gerlach, 1954
crinita Gerlach, 1952
amabilis (de Man, 1922) Luc & De Coninck, 1959
inglisi Warwick, 1970
pontica Filipjev, 1918
Cobbia
treffusiaeformis de Man, 1907
caledonia Warwick & Platt, 1973
simplex Allgén, 1929
dentata Gerlach, 1953
triiodonta Filipjev, 1918
Cobbianema
acrocerca Filipjev, 1922
cylindrilaimoides Schuurmans Stekhoven, 1950
Comesoma
profundi Bastian, 1865
vulgare Bastian, 1865
stenocephalum Filipjev, 1918
Coninckia
macrospirifera Zhang, 1983
circularis Gerlach, 1956
Corythostoma
filipjevi (Kreis, 1928) Wieser, 1956
Crassolaimus
conicaudatus Kreis, 1929
bipapillatus Wieser, 1951
Crenopharynx
paralepturus (Schuurmans Stekhoven, 1950) Wieser, 1953
gracilis (Linstow, 1900) Filipjev, 1934
marioni (Southern, 1914) Filipjev, 1934
Cricolaimus
elongatus Southern, 1914
Cryptenoplus
gerlachi Riemann, 1966
Curvolaimus
uteratus Vitiello, 1970
Cyartonema
zosterae (Allgén, 1929)
tenuicauda (Allgén, 1933) Juario, 1972
roervikensis (Allgén, 1946) Juario, 1972
elegans Jayasree & Warwick, 1977
germanicum Juario, 1972
Cyatholaimus
savaljevi Allgén, 1933
nordgaardi Allgén, 1933
microsetosus Wieser, 1954
gracilis (Eberth, 1863) Bastian, 1865
minutus Allgén, 1933
paucispira Schuurmans Stekhoven, 1950
canariensis Steiner, 1921
scarratti Inglis, 1961
simulatus Kreis, 1924
prinzi (Marion, 1870) Inglis, 1961
papilliferus Allgén, 1929
ocellatus Bastian, 1865
Cylicolaimus
edentulus Platonova, 1970
obtusidens Ditlevsen, 1926
jaegerskioeldi Türk, 1903
magnus (Villot, 1875) de Man, 1889
Cynura
uniformis Cobb, 1920
Dagda
bipapillata Southern, 1914
asymmetrica Gerlach, 1953
Daptionema
pseudotortum (Vitiello, 1971) Lorenzen, 1977
kornoeense (Allgén, 1929) Lorenzen, 1977
trabeculosum (Schneider, 1906) Lorenzen, 1977
svalbardense (Gerlach, 1965) Lorenzen, 1977
planiere (Vitiello, 1971) Lorenzen, 1977
paradonsi (Allgén, 1949) Lorenzen, 1977
groenlandicum (Ditlevsen, 1928) Lorenzen, 1977
fistulatum (Wieser & Hopper, 1967) Lorenzen, 1977
prominens (Vitiello, 1971) Lorenzen, 1977
curvispiculum (Gerlach, 1953) Lorenzen, 1977
hirsutum (Vitiello, 1967) Lorenzen, 1977
calceolatum (De Coninck & Schuurmans Stekhoven, 1933) Lorenzen, 1977
procerum (Gerlach, 1951) Lorenzen, 1977
psammoides (Warwick, 1970) Warwick *et al.*, 1998
setifer (Gerlach, 1952) Lorenzen, 1977
donsi (Allgén, 1948) Lorenzen, 1977
nanum (Lorenzen, 1972) Lorenzen, 1977
proprium (Lorenzen, 1972) Lorenzen, 1977
normandicum (de Man, 1890) Lorenzen, 1977
trichospiculum (Allgén, 1933) Lorenzen, 1977
elegans (Kreis, 1929) Lorenzen, 1977
levis Rieger & Ott, 1971
curvatus Gerlach, 1956
biggi (Gerlach, 1965) Lorenzen, 1977
oxyicerca (de Man, 1888) Lorenzen, 1977
furcatum (Juario, 1974) Lorenzen, 1977
acrilabiatum (De Coninck & Schuurmans Stekhoven, 1933) Lorenzen, 1977
invagiferum (Platt, 1973) Lorenzen, 1977
setosum (Bütschli, 1874) Lorenzen, 1977
conicum (Filipjev, 1922) Lorenzen, 1977
adiectus (Schulz, 1932) Wieser, 1956
spirum (Gerlach, 1959) Lorenzen, 1977
rusticum (Kreis, 1929) Lorenzen, 1977
circulum (Vitiello, 1971) Lorenzen, 1977
fallax (Lorenzen, 1972) Lorenzen, 1977
microscopiculum (Gerlach, 1953) Lorenzen, 1977
vicinum (Riemann, 1966) Lorenzen, 1977
albigensis (Riemann, 1966) Hopper, 1968
buetschlii Schuurmans Stekhoven, 1935
paratortum (Vitiello, 1971) Lorenzen, 1977
tenuispiculum (Ditlevsen, 1918) Lorenzen, 1977
longicaudatum (Filipjev, 1922) Lorenzen, 1977
Dasynemella
conica Gerlach, 1956
albaensis Warwick & Platt, 1973

- pselionemoides* (Gerlach, 1953) De Coninck, 1965
Dasynemoides
longicollis Gerlach, 1952
Deltanema
parvum Kreis, 1929
Demonema
rapax Cobb, 1894
Dentatonema
crassum Kreis, 1928
Denticulella
polydonta (Schulz, 1932) Wieser, 1954
stygia (Gerlach, 1952) Wieser, 1954
Deontolaimus
papillatus de Man, 1880
Deontostoma
arcticum (Ssaweljev, 1912) Filipjev, 1916
lobatum (Steiner, 1916) Platonova, 1962
Desmodora
varioannulata (Kreis, 1928) Lorenzen, 1981
tenuidentata Kreis, 1963
poseidoni Steiner, 1916
cincta (Cobb, 1920) Lorenzen, 1981
polychaeta Allgén, 1929
mediterranea Lorenzen, 1981
coniseta (Schuurmans Stekhoven, 1950) Lorenzen, 1981
microchaeta (Allgén, 1929) Lorenzen, 1981
conica Vitiello, 1971
longiseta (Schuurmans Stekhoven, 1950) Lorenzen, 1981
roscoffiensis Luc & De Coninck, 1959
porifera (Wieser, 1951) Lorenzen, 1981
angusticollis Daday, 1901
ditlevenseni (Micoletzky, 1922) Lorenzen, 1981
nybellini Allgén, 1954
microchaetoides Allgén, 1933
intermedia Allgén, 1940
scaldensis de Man, 1889
tenuispiculum (Allgén, 1928) Lorenzen, 1976
tenuicauda Allgén, 1932
bullata Steiner, 1916
sanguinea Southern, 1914
papillata Daday, 1901
hirsuta (Chitwood, 1936) Gerlach, 1963
pontica (Filipjev, 1922) Lorenzen, 1981
sphaerica (Kreis, 1928) Lorenzen, 1981
communis (Bütschli, 1874) de Man, 1889
septentrionalis Kreis, 1963
Desmolaimus
bulbulus Lorenzen, 1969
zeelandicus de Man, 1880
zosterae Allgén, 1933
tristis Allgén, 1935
mirabilis Allgén, 1935
longicaudatus Kreis, 1929
Desmoscolex
petalodes Lorenzen, 1972
proboscis Lorenzen, 1972
labiosus Lorenzen, 1969
amaurus Lorenzen, 1972
pusillus Lorenzen, 1969
nanellus (Allgén, 1939) Lorenzen, 1971
vanoyei De Coninck, 1943
adenotrichus Lorenzen, 1969
tenuiseta Filipjev, 1922
fennicus Lorenzen, 1969
minor Schepotieff, 1907
adriaticus Schepotieff, 1907
longisetosus Timm, 1970
lanuginosus Panceri, 1876
italicus Timm, 1970
cosmopolites Timm, 1970
californiensis (Timm, 1970) Gerlach & Riemann, 1973
borealis Kreis, 1963
laevis Kreis, 1928
falcatus Lorenzen, 1972
balticus Lorenzen, 1971
annulatus Schepotieff, 1907
americanus Chitwood, 1936
minutus Claparède, 1863
chaetogaster Greeff, 1869
Dichromadora
microdonta Kreis, 1929
geophila (de Man, 1876) Kreis, 1929
cephalata (Steiner, 1916) Kreis, 1929
hyalocheile De Coninck & Schuurmans Stekhoven, 1933
gracilis (Kreis, 1929) Wieser, 1954
islandica Kreis, 1963
cucullata Lorenzen, 1973
abnormis Gerlach, 1953
strandii Allgén, 1940
scandula Lorenzen, 1966
punctata Schuurmans Stekhoven, 1950
Didelta
scutellatum Vitiello, 1969
longicaudatum Vitiello, 1969
cascadum Gerlach, 1956
Diodontolaimus
sabulosus Southern, 1914
Diplolaimella
ocellata (Bütschli, 1874) Gerlach, 1957
signifera (Paramonov, 1929) Gerlach & Riemann, 1973
stagnosa Lorenzen, 1966
punicea Timm, 1952
monhysterooides Allgén, 1929
Diplolaimelloides
altherri Meyl, 1954
bruciei Hopper, 1970
islandicus (De Coninck, 1943) Meyl, 1954
oschei Meyl, 1954
deconincki (Gerlach, 1951) Meyl, 1954
Diplopeltis
cirrhatus (Eberth, 1863) Cobb, 1891
Diplopeltoides
ornatus (Gerlach, 1950) Gerlach, 1962
Diplopeltula
striata Gerlach, 1956
bulbosa Vitiello, 1972
lucanica Boucher & Hellouët, 1977
minuta Vitiello, 1972
longiceps Gerlach, 1950
laminata Vitiello, 1972
cassidaignensis Vitiello, 1972
breviceps Gerlach, 1950
asetosa Juario, 1974
incisa (Southern, 1914) Gerlach, 1962
nuda Gerlach, 1956
intermedia Gerlach, 1954
Disconema
sueicum Allgén, 1935
minutum Vitiello, 1969
longicaudatum Vitiello, 1969
longamphida Vitiello, 1969
alaima Filipjev, 1918
Ditlevsenella
danica Filipjev, 1927
inermis Gerlach, 1953
murmanica Filipjev, 1927
Dolicholaimus
marioni de Man, 1888
Dolichosomatum
klatti Allgén, 1951
Doliolaimus
agilis Lorenzen, 1966
Dorylaimopsis
mediterranea Grimaldi-De Zio, 1968
punctata Ditlevsen, 1918

- Dracognomus*
simplex (Gerlach, 1954) Allen & Noffsinger, 1978
- Draconema*
ophicephalum (Claparedé, 1863) Hope & Murphy, 1972
- micoletzkyi* Kreis, 1928
- ponticum* Filipjev, 1918
- solidum* Gerlach, 1952
- spinosum* (Southern, 1914) Kreis, 1938
- armatum* (Giard & Barrois, 1875) Hope & Murphy, 1972
- inarimense* (Panceri, 1876) Hope & Murphy, 1972
- claparedii* (Metschnikoff, 1867) Hope & Murphy, 1972
- macrocephalum* (Schepotieff, 1907) Wieser, 1954
- cephalatum* Cobb, 1913
- simplex* (Gerlach, 1954) Hope & Murphy, 1972
- hibernicum* (Southern, 1914) Kreis, 1938
- arcticum* Kreis, 1963
- campbelli* (Allgén, 1932) Hope & Murphy, 1972
- longirostrum* (Schepotieff, 1907) Hope & Murphy, 1972
- Echinotheristus*
teutonicus Thun & Riemann, 1967
- cimbricus* Thun & Riemann, 1967
- Eleutherolaimus*
parastenosoma Allgén, 1933
- parasabuliculus* (Allgén, 1935) Gerlach, 1958
- longus* Filipjev, 1922
- demanii* (Rouville, 1903) Filipjev, 1922
- inquietosus* Schuurmans Stekhoven, 1935
- cylindricaudatus* (Schuurmans Stekhoven, 1946) Gerlach, 1963
- stenosoma* (de Man, 1907) Filipjev, 1922
- elegans* Schuurmans Stekhoven, 1942
- duplicatus* Schuurmans Stekhoven, 1950
- Elzalia*
mediterranea Vitiello, 1971
- Enchelidiella*
pellucida Allgén, 1954
- Enoploides*
amphioxii Filipjev, 1918
- brunetti* Gerlach, 1953
- tyrrhenicus* Brunetti, 1949
- cephalophorus* (Ditlevsen, 1918) Filipjev, 1927
- labiatus* (Bütschli, 1874) Filipjev, 1918
- caspersi* Riemann, 1966
- typicus* Ssaweljev, 1912
- labrostriatus* (Southern, 1914) Filipjev, 1921
- spiculohamatus* Schulz, 1932
- longispiculosus* Vitiello, 1967
- brunettii* Gerlach, 1953
- hirsutus* Filipjev, 1918
- cirrhatus* Filipjev, 1918
- Enoplolaimus*
filiformis (Allgén, 1935) Wieser, 1953
- propinquus* de Man, 1922
- subterraneus* Gerlach, 1953
- villusus* Gerlach, 1953
- parapropinquus* Allgén, 1949
- robustus* Gerlach, 1954
- zosterae* Schulz, 1932
- denticulatus* Warwick, 1970
- regius* Hopper, 1962
- glabrus* Brunetti, 1949
- conicollis* Gerlach, 1952
- balgensis* Skwarra, 1921
- attenuatus* Brunetti, 1949
- psammae* Gerlach, 1952
- abnormis* Kreis, 1928
- longicaudatus* (Southern, 1914) Filipjev, 1921
- litoralis* Schulz, 1936
- halophilus* Ditlevsen, 1928
- vulgaris* de Man, 1893
- Enoplus*
- groenlandicus* Ditlevsen, 1926
- alatus* Ssaweljev, 1912
- auriculatus* Ssaweljev, 1912
- brachyuris* Ditlevsen, 1923
- constrictus* Ditlevsen, 1926
- crassus* Filipjev, 1916
- flagellicaudatus* Schuurmans Stekhoven, 1943
- littoralis* Filipjev, 1918
- communis* Bastian, 1865
- maeoticus* Filipjev, 1916
- brevis* Bastian, 1865
- quadridentatus* Berlin, 1853
- schulzi* Gerlach, 1952
- obtusocaudatus* Eberth, 1863
- serratus* Ditlevsen, 1926
- inermis* Bastian, 1865
- sphaericus* Kreis, 1928
- meridionalis* Steiner, 1921
- michaelseni* von Linstow, 1896
- paralittoralis* Wieser, 1953
- striatus* Eberth, 1863
- Epacanthion*
mawsoni Warwick, 1977
- buetschlii* (Southern, 1914) Wieser, 1953
- flagellicauda* Gerlach, 1956
- crassum* (Ditlevsen, 1926) Wieser, 1953
- durapelle* (Kreis, 1929) Wieser, 1953
- enoploidiforme* (Gerlach, 1953) Warwick, 1970
- gorgonocephalum* Warwick, 1970
- incurvatum* (Ditlevsen, 1926) Wieser, 1953
- pellucidum* (Ssaweljev, 1912) Wieser, 1959
- murmanicum* (Ssaweljev, 1912) Wieser, 1959
- Epsiloniloma*
cygnoides (Metschnikoff, 1867) Steiner, 1927
- Ethmolaimus*
multipapillatus Paramonov, 1926
- Eubostrichus*
macintyreii Platt & Zhang, 1982
- phalacrurus* Greeff, 1869
- filiformis* Greeff, 1869
- smo* Platt & Zhang, 1982
- Euchromadora*
inflatispiculum Schuurmans Stekhoven, 1943
- striata* (Eberth, 1863) de Man, 1886
- tridentata* Allgén, 1929
- gaulica* Inglis, 1962
- strandii* Allgén, 1934
- vulgaris* (Bastian, 1865) de Man, 1886
- africana* Linstow, 1908
- Eumorpholaimus*
digiticaudatus Schuurmans Stekhoven, 1946
- norvegicus* Allgén, 1946
- sabuliculus* Schulz, 1932
- Eurystomina*
ornata (Eberth, 1863) Marion, 1870
- cassiterides* (Warwick, 1977) Lambshead & Platt, 1979
- terrerala* (de Man, 1907) Filipjev, 1921
- stenolaimoides* Wieser, 1953
- spectabilis* (Marion, 1870) Filipjev, 1921
- gerlachi* Inglis, 1962
- filispiculum* Gerlach, 1954
- filiformis* (de Man, 1888) Filipjev, 1921
- assimilis* (de Man, 1876) de Man, 1888
- retrociliata* Micoletzky, 1930
- pettiti* Inglis, 1962
- Eusynonchus*
hirsutus (Cobb, 1894) Platonova, 1970
- brevisetosus* (Southern, 1914) Platonova, 1970
- Fenestrolaimus*
murmanicus Filipjev, 1927
- insulaealbae* Filipjev, 1927
- profundis* Filipjev, 1927
- vestitus* Gerlach, 1956
- antarcticus* (Mawson, 1956) Lorenzen, 1994

- pselionemoides* (Gerlach, 1953) De Coninck, 1965
Dasygymnoides
longicollis Gerlach, 1952
Deltanema
parvum Kreis, 1929
Demonema
rapax Cobb, 1894
Dentatonema
crassum Kreis, 1928
Denticulella
polydonta (Schulz, 1932) Wieser, 1954
stygia (Gerlach, 1952) Wieser, 1954
Deontolaimus
papillatus de Man, 1880
Deontostoma
arcticum (Sswaljev, 1912) Filipjev, 1916
lobatum (Steiner, 1916) Platonova, 1962
Desmodora
varioannulata (Kreis, 1928) Lorenzen, 1981
tenuidentata Kreis, 1963
poseidonii Steiner, 1916
cincta (Cobb, 1920) Lorenzen, 1981
polychaeta Allgén, 1929
mediterranea Lorenzen, 1981
coniseta (Schuurmans Stekhoven, 1950) Lorenzen, 1981
microchaeta (Allgén, 1929) Lorenzen, 1981
conica Vitiello, 1971
longiseta (Schuurmans Stekhoven, 1950) Lorenzen, 1981
roscophiensis Luc & De Coninck, 1959
porifera (Wieser, 1951) Lorenzen, 1981
angusticollis Daday, 1901
ditlevenseni (Micoletzky, 1922) Lorenzen, 1981
nybellini Allgén, 1954
microchaetoides Allgén, 1933
intermedia Allgén, 1940
scaldensis de Man, 1889
tenuispiculum (Allgén, 1928) Lorenzen, 1976
tenuicauda Allgén, 1932
bullata Steiner, 1916
sanguinea Southern, 1914
papillata Daday, 1901
hirsuta (Chitwood, 1936) Gerlach, 1963
pontica (Filipjev, 1922) Lorenzen, 1981
sphaerica (Kreis, 1928) Lorenzen, 1981
communis (Bütschli, 1874) de Man, 1889
septentrionalis Kreis, 1963
Desmolaimus
bulbus Lorenzen, 1969
zeelandicus de Man, 1880
zosterae Allgén, 1933
tristis Allgén, 1935
mirabilis Allgén, 1935
longicaudatus Kreis, 1929
Desmoscolex
petalodes Lorenzen, 1972
proboscis Lorenzen, 1972
labiosus Lorenzen, 1969
amaurus Lorenzen, 1972
pusillus Lorenzen, 1969
nanellus (Allgén, 1939) Lorenzen, 1971
vanoyei De Coninck, 1943
adenotrichus Lorenzen, 1969
tenuiseta Filipjev, 1922
fennicus Lorenzen, 1969
minor Schepotieff, 1907
adriaticus Schepotieff, 1907
longisetosus Timm, 1970
lanuginosus Panceri, 1876
italicus Timm, 1970
cosmopolites Timm, 1970
californiensis (Timm, 1970) Gerlach & Riemann, 1973
borealis Kreis, 1963
laevis Kreis, 1928
falcatus Lorenzen, 1972
balticus Lorenzen, 1971
annulatus Schepotieff, 1907
americanus Chitwood, 1936
minutus Claparède, 1863
chaetogaster Greeff, 1869
Dichromadora
microdonta Kreis, 1929
geophila (de Man, 1876) Kreis, 1929
cephalata (Steiner, 1916) Kreis, 1929
halocheile De Coninck & Schuurmans Stekhoven, 1933
gracilis (Kreis, 1929) Wieser, 1954
islandica Kreis, 1963
cucullata Lorenzen, 1973
abnormis Gerlach, 1953
strandii Allgén, 1940
scandula Lorenzen, 1966
punctata Schuurmans Stekhoven, 1950
Didelta
scutellatum Vitiello, 1969
longicaudatum Vitiello, 1969
casclatum Gerlach, 1956
Diiodontolaimus
sabulosus Southern, 1914
Diplolaimella
ocellata (Bütschli, 1874) Gerlach, 1957
signifera (Paramonov, 1929) Gerlach & Riemann, 1973
stagnosa Lorenzen, 1966
punicea Timm, 1952
monhysteroides Allgén, 1929
Diplolaimelloides
altherri Meyl, 1954
bruciei Hopper, 1970
islandicus (De Coninck, 1943) Meyl, 1954
oschei Meyl, 1954
deconincki (Gerlach, 1951) Meyl, 1954
Diplopeltis
cirrhatus (Eberth, 1863) Cobb, 1891
Diplopeltoides
ornatus (Gerlach, 1950) Gerlach, 1962
Diplopeltula
striata Gerlach, 1956
bulbosa Vitiello, 1972
lucanica Boucher & Hellouët, 1977
minuta Vitiello, 1972
longiceps Gerlach, 1950
laminata Vitiello, 1972
cassidaigniensis Vitiello, 1972
breviceps Gerlach, 1950
asetosa Juario, 1974
incisa (Southern, 1914) Gerlach, 1962
nuda Gerlach, 1956
intermedia Gerlach, 1954
Disconema
sueicum Allgén, 1935
minutum Vitiello, 1969
longicaudatum Vitiello, 1969
longamphida Vitiello, 1969
alaima Filipjev, 1918
Ditlevsenella
danica Filipjev, 1927
inermis Gerlach, 1953
murmanica Filipjev, 1927
Dolicholaimus
marioni de Man, 1888
Dolichosomatum
klatti Allgén, 1951
Doliolaimus
agilis Lorenzen, 1966
Dorylaimopsis
mediterranea Grimaldi-De Zio, 1968
punctata Ditlevsen, 1918

- Dracognomus*
simplex (Gerlach, 1954) Allen & Noffsinger, 1978
- Draconema*
ophicephalum (Claparedé, 1863) Hope & Murphy, 1972
micoletzkyi Kreis, 1928
ponticum Filipjev, 1918
solidum Gerlach, 1952
spinosum (Southern, 1914) Kreis, 1938
armatum (Giard & Barrois, 1875) Hope & Murphy, 1972
inarimense (Panceri, 1876) Hope & Murphy, 1972
claparedii (Metschnikoff, 1867) Hope & Murphy, 1972
macrocephalum (Schepotieff, 1907) Wieser, 1954
cephalatum Cobb, 1913
simplex (Gerlach, 1954) Hope & Murphy, 1972
hibernicum (Southern, 1914) Kreis, 1938
arcticum Kreis, 1963
campbelli (Allgén, 1932) Hope & Murphy, 1972
longirostrum (Schepotieff, 1907) Hope & Murphy, 1972
- Echinotheristus*
teutonicus Thun & Riemann, 1967
cimbicus Thun & Riemann, 1967
- Eleutherolaimus*
parastenosoma Allgén, 1933
parasabulicolus (Allgén, 1935) Gerlach, 1958
longus Filipjev, 1922
demanii (Rouville, 1903) Filipjev, 1922
inquietosus Schuurmans Stekhoven, 1935
cylindricaudatus (Schuurmans Stekhoven, 1946) Gerlach, 1963
stenosoma (de Man, 1907) Filipjev, 1922
elegans Schuurmans Stekhoven, 1942
duplicatus Schuurmans Stekhoven, 1950
- Elzalia*
mediterranea Vitiello, 1971
- Enchelidiella*
pellucida Allgén, 1954
- Enoploides*
amphioxi Filipjev, 1918
brunetti Gerlach, 1953
tyrrhenicus Brunetti, 1949
cephalophorus (Ditlevsen, 1918) Filipjev, 1927
labiatus (Bütschli, 1874) Filipjev, 1918
caspersi Riemann, 1966
typicus Ssaweljev, 1912
labrostriatus (Southern, 1914) Filipjev, 1921
spiculohamatus Schulz, 1932
longispiculosus Vitiello, 1967
brunettii Gerlach, 1953
hirsutus Filipjev, 1918
cirrhatus Filipjev, 1918
- Enoplolaimus*
filiformis (Allgén, 1935) Wieser, 1953
propinquus de Man, 1922
subterraneus Gerlach, 1953
villosus Gerlach, 1953
parapropinquus Allgén, 1949
robustus Gerlach, 1954
zosterae Schulz, 1932
denticulatus Warwick, 1970
regius Hopper, 1962
glabrus Brunetti, 1949
conicollis Gerlach, 1952
balgensis Skwara, 1921
attenuatus Brunetti, 1949
psammae Gerlach, 1952
abnormis Kreis, 1928
longicaudatus (Southern, 1914) Filipjev, 1921
litoralis Schulz, 1936
halophilus Ditlevsen, 1928
vulgaris de Man, 1893
- Enoplus*
- groenlandicus* Ditlevsen, 1926
alatus Ssaweljev, 1912
auriculatus Ssaweljev, 1912
brachyuris Ditlevsen, 1923
constrictus Ditlevsen, 1926
crassus Filipjev, 1916
flagellicaudatus Schuurmans Stekhoven, 1943
littoralis Filipjev, 1918
communis Bastian, 1865
maeoticus Filipjev, 1916
brevis Bastian, 1865
quadridentatus Berlin, 1853
schulzi Gerlach, 1952
obtusocaudatus Eberth, 1863
serratus Ditlevsen, 1926
inermis Bastian, 1865
sphaericus Kreis, 1928
meridionalis Steiner, 1921
michaelseni von Linstow, 1896
paralittoralis Wieser, 1953
striatus Eberth, 1863
- Epacanthion*
mawsoni Warwick, 1977
buetschlii (Southern, 1914) Wieser, 1953
flagellicauda Gerlach, 1956
crassum (Ditlevsen, 1926) Wieser, 1953
durapelle (Kreis, 1929) Wieser, 1953
enoploidiforme (Gerlach, 1953) Warwick, 1970
gorgonocephalum Warwick, 1970
incurvatum (Ditlevsen, 1926) Wieser, 1953
pellucidum (Ssaweljev, 1912) Wieser, 1959
murmanicum (Ssaweljev, 1912) Wieser, 1959
- Epsiloninema*
cygnoides (Metschnikoff, 1867) Steiner, 1927
- Ethmolaimus*
multipipillatus Paramonov, 1926
- Eubostrichus*
macintyrei Platt & Zhang, 1982
phalacrurus Greeff, 1869
filiformis Greeff, 1869
smo Platt & Zhang, 1982
- Euchromadora*
inflatispiculum Schuurmans Stekhoven, 1943
striata (Eberth, 1863) de Man, 1886
tridentata Allgén, 1929
gaulica Inglis, 1962
strandii Allgén, 1934
vulgaris (Bastian, 1865) de Man, 1886
africana Linstow, 1908
- Eumorpholaimus*
digiticaudatus Schuurmans Stekhoven, 1946
norvegicus Allgén, 1946
sabulicolus Schulz, 1932
- Eurystomina*
ornata (Eberth, 1863) Marion, 1870
cassiterides (Warwick, 1977) Lambshead & Platt, 1979
terricola (de Man, 1907) Filipjev, 1921
stenolaimoides Wieser, 1953
spectabilis (Marion, 1870) Filipjev, 1921
gerlachi Inglis, 1962
filispiculum Gerlach, 1954
filiformis (de Man, 1888) Filipjev, 1921
assimilis (de Man, 1876) de Man, 1888
retrociliata Micoletzky, 1930
pettiti Inglis, 1962
- Eusynonchus*
hirsutus (Cobb, 1894) Platonova, 1970
brevisetosus (Southern, 1914) Platonova, 1970
- Fenestrolaimus*
murmanicus Filipjev, 1927
insulaealbae Filipjev, 1927
profundis Filipjev, 1927
vestitus Gerlach, 1956
antarcticus (Mawson, 1956) Lorenzen, 1994

- Filipjeva*
parameridionalis Vitiello, 1970
arctica Ditlevsen, 1928
mediterranea Schuurmans Stekhoven, 1950
- Filipjevia*
macrolabiata Kreis, 1928
- Filoncholaimus*
filicaudatus (Ditlevsen, 1926) Filipjev, 1927
- Fusonema*
multilabiatum Kreis, 1928
- Gairleanema*
anagremiae Warwick & Platt, 1973
- Gammannema*
mediterraneum Vitiello, 1970
conicauda Gerlach, 1953
rapax (Ssaweljev, 1912) Gerlach, 1964
- Gammarinema*
ligiae Gerlach, 1967
gammari Kinne & Gerlach, 1953
- Gerlachius*
novisetosus Platt & Zhang, 1982
- Gonianchus*
cumbaensis Benwell, 1981
inaequalis Warwick & Platt, 1973
longicaudatus (Ward, 1972)
- Graphonema*
scampae (Coles, 1965) Platt & Warwick, 1988
northumbriae Warwick & Coles, 1975
- Greeffella*
oxycaudata (Greeff, 1869) Cobb, 1922
beatlei Lorenzen, 1969
comosa Lorenzen, 1969
moppa Schrage, 1972
- Halalaimus*
monstrocaudatus Vitiello, 1970
cirrhatus Gerlach, 1953
delamarei Vitiello, 1970
diplocephalus Filipjev, 1927
horridus Gerlach, 1956
longicollis Allgén, 1932
pachyderma (Filipjev, 1927) Schuurmans Stekhoven, 1935
zenkevitshi Filipjev, 1927
amphistrius Vitiello, 1970
filicorpus Vitiello, 1970
papillifer Gerlach, 1956
lutarus Vitiello, 1970
capitulatus Boucher, 1978
borealis (Steiner, 1916) Lorenzen, 1981
striatus Gerlach, 1956
gracilis de Man, 1888
longicaudatus (Filipjev, 1927) Schneider, 1939
amphidellus Vitiello, 1970
pachyodoroides Vitiello, 1970
ponticus Filipjev, 1922
turbidus Vitiello, 1970
terrestris Gerlach, 1959
isaitshikovi (Filipjev, 1927) Schuurmans Stekhoven, 1935
leptosoma (Southern, 1914) Schuurmans Stekhoven, 1935
- Halanonchus*
papillatus Groza-Rojancovski, 1972
cornutus Vitiello, 1971
zosterae (Allgén, 1933) Gerlach, 1964
- Halaphanolaimus*
longisetosus Allgén, 1928
minutus Schuurmans Stekhoven, 1942
pellucidus Southern, 1914
- Halichoanolaimus*
norvegicus Allgén, 1940
minor Ssaweljev, 1912
consimilis Allgén, 1933
robustus (Bastian, 1865) de Man, 1886
lanceolatus Vitiello, 1970
- dolichurus* Ssaweljev, 1912
- Halinema*
norwegianum Allgén, 1933
- Haliplectus*
bibulbosus (Schulz, 1935) Gerlach, 1957
bickneri Chitwood, 1956
wheeleri Coles, 1965
dorsalis Chitwood, 1956
schulzi (De Coninck, 1943) Gerlach, 1957
- Hapalomus*
terrestris Lorenzen, 1969
- Heterocyatholaimus*
macrolaimus Allgén, 1935
- Hoffmaenneria*
niddensis (Skwarra, 1921) Schneider, 1940
- Hopperia*
massiliensis Vitiello, 1969
- Hypodontolaimus*
inaequalis (Bastian, 1865) de Man, 1886
schuurmansstekhoveni Gerlach, 1951
balticus (Schneider, 1906) Filipjev, 1918
abyssalis Allgén, 1933
heymonsii (Steiner, 1921) Filipjev, 1930
longiseta (Allgén, 1933) Wieser, 1954
mediterraneus Brunetti, 1949
sivertseni Allgén, 1951
setosus (Bütschli, 1874) Wieser, 1954
colesi Inglis, 1962
- Innocuonema*
tentabunda (de Man, 1890) Inglis, 1969
- Ionema*
cobbi (Steiner, 1916) Micoletzky, 1924
coecum Kreis, 1963
boreale Kreis, 1963
- Ixonema*
sordidum Lorenzen, 1971
- Klugea*
pigmentata (Ditlevsen, 1926) Filipjev, 1927
trilabia Filipjev, 1927
filiformis (Allgén, 1929) Filipjev, 1934
polaris (Steiner, 1916) Filipjev, 1927
stenolaima (Wieser, 1954) Wieser, 1954
- Krampia*
opaca (Allgén, 1930) Kreis, 1934
longicollis Allgén, 1939
acropora Ditlevsen, 1921
- Kraspedonema*
octogonata Gerlach, 1954
- Laimella*
minuta Vitiello, 1970
vera Vitiello, 1971
longicauda Cobb, 1920
- Latronema*
aberrans (Allgén, 1934) Gerlach, 1964
orcinum (Gerlach, 1952) Wieser, 1954
- Lauratonema*
adriaticum Gerlach, 1953
reductum Gerlach, 1953
- Lauratonemoides*
originalis (Gerlach, 1956) De Coninck, 1965
- Laxus*
contortus Cobb, 1894
- Ledovitria*
obtusidens (Schuurmans Stekhoven, 1956) Vitiello, 1970
longiseta (Micoletzky, 1930) Wieser, 1953
hirsuta Filipjev, 1927
- Leptolaimoides*
tubulosus Vitiello, 1971
thermastris (Lorenzen, 1966) Vitiello, 1971
- Leptolaimus*
macer Lorenzen, 1972
turridus Timm, 1963
longiseta (Allgén, 1934) De Coninck, 1965
puccinelliae Gerlach, 1959

- leptaleus* Lorenzen, 1972
membranatus (Wieser, 1951) De Coninck, 1965
minutus Vitiello, 1971
mixtus Lorenzen, 1972
nobilis Gerlach, 1956
papilliger de Man, 1876
punicosus Vitiello, 1971
steineri (Filipjev, 1922) De Coninck, 1965
scotlandicus Jayasree & Warwick, 1977
limiculus Lorenzen, 1972
elegans (Schuurmans Stekhoven & De Coninck, 1933)
 Gerlach, 1958
ampullaceus Warwick, 1970
trichodes (Kreis, 1929) De Coninck, 1965
venustus Lorenzen, 1972
alatus Vitiello, 1971
acicula Lorenzen, 1966
ditlevenseni (Steiner, 1916) De Coninck, 1965
setiger Schuurmans Stekhoven & De Coninck, 1933
timmi Vitiello, 1971
cupulatus Lorenzen, 1972
tenuis Gerlach, 1956
- Leptonemella*
gorgo Gerlach, 1950
parabullata (Allgén, 1929) Gerlach, 1963
aphanothecae Gerlach, 1950
froeiensis (Allgén, 1946) Gerlach, 1963
- Leptosomatides*
steineri Filipjev, 1922
euxinus Filipjev, 1918
- Leptosomatina*
conicaudata Vitiello, 1970
appendixocaudata Allgén, 1957
- Leptosomatum*
bacillatum (Eberth, 1863) Bastian, 1865
arcticum Filipjev, 1916
tetraphthalmum Ssweljev, 1912
sabangense Steiner, 1915
elongatum Bastian, 1865
longisetosum Schuurmans Stekhoven, 1943
filipjevi Schuurmans Stekhoven, 1950
breviceps Platonova, 1967
gracile Bastian, 1865
subulatum (Eberth, 1863) Bastian, 1865
punctatum (Eberth, 1863) Bastian, 1865
bathybium Allgén, 1954
caecum Ditlevsen, 1923
- Leptosomella*
acrocerca Filipjev, 1927
- Lineolia*
sieboldii (Kölliker, 1845) Gerlach & Riemann, 1974
- Linhomoeus*
paralineatus Allgén, 1933
timmi Inglis, 1963
elongatus Bastian, 1865
ponticus Filipjev, 1922
pilosus (Schuurmans Stekhoven, 1950) Wieser, 1956
paralongicaudatus (Allgén, 1934) Wieser, 1956
macramphis Schuurmans Stekhoven, 1942
hirsutus Bastian, 1865
kornoeensis Allgén, 1929
gracilisetosus (Allgén, 1946) Wieser, 1956
brevisetosus Schuurmans Stekhoven, 1946
- Litinium*
parmatum Wieser, 1954
bananum Gerlach, 1956
simplex (Bird, 1995) Lorenzen, 1981
- Longicyatholaimus*
cervoides Vitiello, 1970
longicaudatus (de Man, 1876) Micoletzky, 1924
falcatus Vitiello, 1971
filicaudatus Schuurmans Stekhoven, 1950
capsulatus Vitiello, 1970
- Manunema*
proboscidis Gerlach, 1957
- annulatum* (Vitiello & De Coninck, 1968) Riemann et al., 1971
- Margonema*
longisetosum Allgén, 1935
- Marylynna*
bellula (Vitiello, 1970) Hopper, 1977
choanolaimoides (Schuurmans Stekhoven, 1942)
 Hopper, 1977
effilata (Schuurmans Stekhoven, 1946) Hopper, 1977
stekhoveni (Wieser, 1954) Hopper, 1977
complexa (Warwick, 1971) Hopper, 1977
- Megadesmolaimus*
incisus Wieser, 1954
- Megeuryostomina*
combesi Luc & De Coninck, 1959
- Mesacanthion*
audax (Ditlevsen, 1918) Filipjev, 1927
longispiculum Gerlach, 1954
tenuicaudatum (Ssweljev, 1912) De Coninck & Schuurmans Stekhoven, 1933
ditlevenseni (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
conicum (Filipjev, 1918) Filipjev, 1927
banale (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
lucifer (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
agubernatus Vitiello, 1971
africaniforme Warwick, 1970
infantilis (Ditlevsen, 1930) De Coninck & Schuurmans Stekhoven, 1933
breviseta (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
hirsutum Gerlach, 1953
diplechma (Southern, 1914) Filipjev, 1927
virile (Ditlevsen, 1930) De Coninck & Schuurmans Stekhoven, 1933
majus (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
- Mesacanthoides*
latignathus (Ditlevsen, 1918) Wieser, 1953
caputmedusae (Ditlevsen, 1918) Wieser, 1959
- Metachromadora*
cystoseirae Gerlach & Riemann, 1973
quadribulba Gerlach, 1956
macrourera Filipjev, 1918
remanei Gerlach, 1951
vivipara (de Man, 1907) Allgén, 1928
suecica (Allgén, 1929) Schulz, 1938
scotlandica Warwick & Platt, 1973
- Metacyatholaimus*
cylindribucca (Schuurmans Stekhoven, 1950) Wieser, 1954
hirschi Schuurmans Stekhoven, 1942
- Metacylicolaimus*
effilatus Schuurmans Stekhoven, 1946
filicaudatus (Ditlevsen, 1926) Schuurmans Stekhoven, 1946
flagellicaudatus Schuurmans Stekhoven, 1946
- Metadasynemella*
macrophalla De Coninck, 1942
cassidinensis Vitiello & Haspeslagh, 1972
falciphalla Vitiello & Haspeslagh, 1972
- Metadesmodora*
amphidiscata Schuurmans Stekhoven, 1942
- Metadesmolaimus*
aduncus Lorenzen, 1972
gelana (Warwick & Platt, 1973)
aversivulva Gerlach, 1953
coronatus Schuurmans Stekhoven, 1950
heteroclitus Lorenzen, 1972
pandus Lorenzen, 1972
varians Lorenzen, 1972
labiosetosus Schuurmans Stekhoven, 1935
tersus (Gerlach, 1956) Lorenzen, 1972

- hamatus* (Gerlach, 1956) Lorenzen, 1972
Metalaimus
gracilis Kreis, 1928
Metalinhomoeus
zosterae Filipjev, 1918
effilatus Schuurmans Stekhoven, 1942
paratypicus Vitiello, 1969
musaecauda Lorenzen, 1966
microsetosus Allgén, 1953
maroeyensis Allgén, 1946
gracilis (Kreis, 1929) Wieser, 1956
flagellicaudatus Schuurmans Stekhoven, 1935
tenuicaudatus Allgén, 1935
parvasetosus Vitiello, 1969
pareffilatus Wieser, 1956
filicaudatus (Allgén, 1930)
tristis (Allgén, 1933)
breviseta Schuurmans Stekhoven, 1950
biratus Vitiello, 1969
typicus de Man, 1907
longiseta Kreis, 1929
filiformis (de Man, 1907) Schuurmans Stekhoven, 1935
Metaparoncholaimus
campylocercus (de Man, 1876) De Coninck & Schuurmans Stekhoven, 1933
Metaraeolaimoides
oxystoma De Coninck, 1936
Metepsilonema
hagmeieri (Stauffer, 1924) Steiner, 1927
Metoncholaimus
pristurus (Zur Strassen, 1894) Filipjev, 1918
albidus (Bastian, 1865) Filipjev, 1918
demani (Zur Strassen, 1894) Filipjev, 1918
anthophorus (Ssaweljev, 1912) Filipjev, 1918
scanicus (Allgén, 1935) Warwick, 1977
Meylia
alata Gerlach, 1956
lissa Gerlach, 1956
spinosa Gerlach, 1956
Micoletzyia
mucronata Vitiello, 1970
magna Vitiello, 1970
parelegans Allgén, 1954
elegans Ditlevsen, 1926
Microlaimus
criminalis Rieger & Ott, 1971
ostracion Schuurmans Stekhoven, 1935
cyatholaimoides de Man, 1922
acuticaudatus De Coninck & Schuurmans Stekhoven, 1933
crassiceps Gerlach, 1953
compridus Gerlach, 1956
citrus Gerlach, 1959
borealis Steiner, 1916
arenicola Schulz, 1938
allgeni Gerlach, 1950
zosterae Allgén, 1930
spirifer Warwick, 1970
dentatus Allgén, 1935
marinus (Schulz, 1932) Schuurmans Stekhoven & De Coninck, 1933
cuanensis Platt, 1973
conothelis (Lorenzen, 1973) Jensen, 1978
acinaces Warwick & Platt, 1973
robustidens Schuurmans Stekhoven & De Coninck, 1933
teutonicus Riemann, 1966
parahonestus Gerlach, 1950
pygmaeus Meyl, 1954
honestus de Man, 1922
undulatus Gerlach, 1953
turgofrons Lorenzen, 1972
texianus Chitwood, 1951
monstrosus Gerlach, 1953
tenuicollis Gerlach, 1952
donsi Allgén, 1935
punctulatus Gerlach, 1950
paraborealis Allgén, 1940
microseta Gerlach, 1953
inermis Ditlevsen, 1923
tenuispiculum de Man, 1922
honestoides Meyl, 1954
lazonus Vitiello, 1971
macrocirculus Gerlach, 1950
globiceps de Man, 1880
Minolaimus
cervoides Vitiello, 1970
lineatus Vitiello, 1970
Molgolaimus
demani Jensen, 1978
Monhyphista
paraambigua Allgén, 1933
macquariensis Allgén, 1929
filiformis Bastian, 1865
disjuncta Bastian, 1865
macrolabiata Kreis, 1928
microphthalmia de Man, 1880
minuta Filipjev, 1929
natans (Bastian, 1865) Bütschli, 1874
paramacramphis Meyl, 1954
parasimplex De Coninck, 1943
paradisjuncta De Coninck, 1943
parelegantula De Coninck, 1943
refringens Schuurmans Stekhoven, 1935
rotundicapitata Filipjev, 1922
sabulicola Allgén, 1929
socialis Bütschli, 1874
multisetosa Meyl, 1955
attenuata Filipjev, 1922
collaris Filipjev, 1922
fasciculata Skwarra, 1921
filicaudata Allgén, 1929
anophthalma Lorenzen, 1969
gracilicauda Schuurmans Stekhoven, 1950
heterolaima Schuurmans Stekhoven, 1950
islandica De Coninck, 1943
izhorica Filipjev, 1929
paraambiguoidea Allgén, 1932
luisae Schuurmans Stekhoven, 1935
macramphis Filipjev, 1929
barentsi Steiner, 1916
capitata Schuurmans Stekhoven, 1942
cartaromanae Meyl, 1954
ampullocauda Paramonov, 1926
droebachiensis Allgén, 1931
parva (Bastian, 1865) Bütschli, 1874
cephalophora de Man, 1876
tenuis Schuurmans Stekhoven, 1950
longicapitata Filipjev, 1922
trichura Allgén, 1930
vulgaris de Man, 1880
similis Bütschli, 1873
elegantula Schuurmans Stekhoven, 1935
Monoposthia
mirabilis Schulz, 1932
costata (Bastian, 1865) de Man, 1889
mielcki Steiner, 1916
arctica Allgén, 1954
Morlaixia
obscura Vincx & Gourbault, 1988
Nannolaimoides
effilatus (Boucher, 1976) Platt & Warwick, 1988
Nannolaimus
volutus Gerlach, 1956
fusus Gerlach, 1956
Nemanema
rectoresectum Kreis, 1928
sabulicola (Allgén, 1939) Wieser, 1953
cylindraticeaudatum (de Man, 1922) Filipjev, 1927

- rotundicaudatum* (Ditlevsen, 1926) Filipjev, 1946
filiforme (Filipjev, 1918) Kreis, 1926
Nemella
ocellata Cobb, 1920
Neochromadora
izhorica (Filipjev, I., 1929) Schuurmans Stekhoven, 1935
paractecta Blome, 1974
trichophora (Steiner, 1921) Gerlach, 1951
sabulicola (Filipjev, 1918) Wieser, 1954
poeilosomoides (Filipjev, 1918) Micoletzky, 1924
munita Lorenzen, 1971
craspedota (Steiner, 1916) Micoletzky, 1924
coudenhovei Wieser, 1956
brevisetosa Wieser, 1954
amembranata Wieser, 1954
poeilosoma (de Man, 1893) Micoletzky, 1924
angelica Riemann, 1976
paramunita Boucher, 1976
tecta Gerlach, 1951
Neotonchus
vittius (Warwick, 1971)
corcundus (Gerlach, 1956) Wieser & Hopper, 1966
cupulatus Vitiello, 1970
meeki Warwick, 1971
pseudocorcundus Vitiello, 1971
boucheri Platt, 1982
warwicki (Platt, 1982)
interruptus (Warwick, 1971)
votadinii (Warwick, 1971)
cuaneensis (Platt, 1982)
Nijhoffia
longicaudata Allgén, 1935
Notochaetosoma
killieri Warwick, 1977
Nudora
thorakista (Schulz, 1935) Gerlach & Riemann, 1973
bipapillata Platt, 1973
gourbaultae Vanreusel & Vincx, 1989
steineri (Steiner, 1921) Wieser, 1954
Octonchus
marinus (Schulz, 1932) Clark, 1961
Odontophora
rectangula Lorenzen, 1972
marina Bütschli, 1874
armata (Ditlevsen, 1918) Allgén, 1929
parangustilaima Wieser, 1956
parasetosa (Allgén, 1929) Allgén, 1929
paratenuicaudata Allgén, 1942
phalarata Lorenzen, 1972
angustilaimus (Filipjev, 1918) Allgén, 1929
longicaudata Schuurmans Stekhoven & De Coninck, 1933
fatica Vitiello, 1971
tenuicaudata Allgén, 1935
ornata Lorenzen, 1972
wieseri Luc & De Coninck, 1959
villoti Luc & De Coninck, 1959
longisetosa (Allgén, 1928) Allgén, 1929
exharena Warwick & Platt, 1973
setosa (Allgén, 1929) Allgén, 1929
Odontophoroides
paramonhystera Lambshead, 1982
Omicronema
truncatum Schuurmans Stekhoven, 1950
nidrosiense Allgén, 1933
Onchiium
robustum Gerlach, 1965
ocellatum Cobb, 1920
conicaudatum (Allgén, 1935) Wieser, 1956
parocellum (Allgén, 1940) Gerlach & Riemann, 1973
Oncholaimellus
mediterraneus Schuurmans Stekhoven, 1942
calvadosicus de Man, 1890
Oncholaimus
oxyuris Ditlevsen, 1911
marinus Schulz, 1932
vesicarius (Wieser, 1959) Rachor, 1969
onchouris Ditlevsen, 1928
groenlandicus Ditlevsen, 1926
medius Kreis, 1928
brachycercus de Man, 1889
steineri Ditlevsen, 1928
skawensis Ditlevsen, 1921
donsi (Allgén, 1947) Wieser, 1953
attenuatus Dujardin, 1845
viridis Bastian, 1865
dujardinii de Man, 1876
lanceolatus Vitiello, 1970
septentrionalis Filipjev, 1927
brevicaudatus Filipjev, 1918
aegypticus Steiner, 1921
spissus Allgén, 1935
serratus Allgén, 1935
campyloceroides De Coninck & Schuurmans Stekhoven, 1933
scanicus (Allgén, 1935) Wieser, 1953
conicauda Filipjev, 1929
curvicauda Allgén, 1957
steinboecki Ditlevsen, 1928
paroxyuris Schuurmans Stekhoven, 1942
Onyx
perfectus Cobb, 1891
sagittarius Gerlach, 1950
Oxyonchus
polaris (Filipjev, 1927) De Coninck & Schuurmans Stekhoven, 1933
pachylabiatus Schuurmans Stekhoven, 1946
hamatus (Steiner, 1916) Filipjev, 1927
acantholaimus (Ssaweljev, 1912) Filipjev, 1927
dentatus (Ditlevsen, 1918) Filipjev, 1927
dubius (Filipjev, 1918) De Coninck & Schuurmans Stekhoven, 1933
Oxystomina
flicauda (Kreis, 1929) Wieser, 1953
nidroensiis Allgén, 1933
alpha Chitwood, 1937
islandica (De Coninck, 1943) Wieser, 1953
novozemelica (Filipjev, 1927) Wieser, 1953
nuda (Filipjev, 1927) Wieser, 1953
pulchella Vitiello, 1970
unguiculata Schuurmans Stekhoven, 1935
brevicaudata Gerlach & Riemann, 1974
alpatovi (Filipjev, 1927) Wieser, 1953
elongata (Bütschli, 1874) Filipjev, 1921
oxycaudata (Ditlevsen, 1926) Filipjev, 1921
asetosa (Southern, 1914) Filipjev, 1921
tenuis Thorne, 1939
clavicauda (Filipjev, 1918) Filipjev, 1921
Pandolaimus
kornoensis (Allgén, 1929) Jensen, 1976
viscosa (Allgén, 1935) Jensen, 1976
latilaimus (Allgén, 1929) Schuurmans Stekhoven, 1935
Paracanthonchus
multitubifer Timm, 1961
spectabilis Allgén, 1931
macrodon (Ditlevsen, 1918) Micoletzky, 1924
breviseta (Schuurmans Stekhoven, 1950) Hope & Murphy, 1972
longicaudatus Warwick, 1971
thaumasius (Schulz, 1932)
platti Vadhyar, 1980
heterodontus (Schulz, 1932)
caecus Micoletzky, 1924
longus Allgén, 1934
elongatus (de Man, 1907) Micoletzky, 1924
steueri (Micoletzky, 1922) Micoletzky, 1924
sunesoni (Allgén, 1942) Wieser, 1954
tyrrhenicus (Brunetti, 1949) Gerlach, 1953

- uniformis* (Schuurmans Stekhoven, 1950) Wieser, 1954
paradoxus Allgén, 1929
nannodontus (Schulz, 1932) Wieser, 1954
microdon (Ditlevsen, 1918) Wieser, 1954
micoletzkyi Schuurmans Stekhoven, 1943
major (Kreis, 1928) Wieser, 1954
filipjevi Micoletzky, 1924
stekhoveni Wieser, 1954
bipapillatus Kreis, 1928
angulatus (Schuurmans Stekhoven, 1950) Wieser, 1954
inglisi Coles, 1965
barka Inglis, 1962
Paracomesoma
dubium (Filipjev, 1918) Schuurmans Stekhoven, 1950
Paracyatholaimoides
labiosetosus Riemann, 1966
multispiralis Gerlach, 1953
Paracyatholaimus
proximus (Bütschli, 1874) Micoletzky, 1924
arcticus Kreis, 1963
dubiosus (Bütschli, 1874) Micoletzky, 1921
oistosticuloides (Allgén, 1935) Wieser, 1954
parasaveljevi Allgén, 1935
pentodon Riemann, 1966
separatus Wieser, 1954
occultus Gerlach, 1956
intermedius (de Man, 1880) Micoletzky, 1922
Paracyclicolaimus
brevisetosus Platonova, 1970
Paradesmodora
cephalata Schuurmans Stekhoven, 1950
Paradracanema
spinosum (Southern, 1914) Allen & Noffsinger, 1978
Paraeagialoalaimus
filiocaudatus Allgén, 1934
Paralinhomoeus
strandibrevicaudatus Allgén, 1934
linurus (Ssweljev, 1912)
viscosus (Allgén, 1929) Allgén, 1934
trichurus (Allgén, 1933)
lineatus (Ditlevsen, 1918) Allgén, 1959
tenuicaudatus (Bütschli, 1874) de Man, 1907
staurenensis Allgén, 1940
uniovarium Warwick, 1970
conicaudatus (Allgén, 1930)
lepturus (de Man, 1907) de Man, 1907
brevibucca Schuurmans Stekhoven, 1950
intermedius (Allgén, 1929) Wieser, 1956
amphilabiatus Schuurmans Stekhoven, 1942
bocki Schuurmans Stekhoven, 1946
brevicaudatus Schuurmans Stekhoven, 1950
caxinus Vitiello, 1969
clavicaudatus Schuurmans Stekhoven, 1950
filiformis (Filipjev, 1918)
ilenensis (Allgén, 1933) Gerlach, 1963
anteporus Vitiello, 1969
deconincki Groza-Rojancovski, 1972
Paralongicyatholaimus
zosterae (Allgén, 1933) Lorenzen, 1972
mastigodes Schuurmans Stekhoven, 1950
minutus Warwick, 1971
macramphis Lorenzen, 1972
Paramesacanthion
marei Warwick, 1970
truncum Vitiello, 1971
tricuspidis (Schuurmans Stekhoven, 1950) Wieser, 1953
klugei (Filipjev, 1927) Wieser, 1953
hirsutum Warwick, 1970
catellum Boucher, 1970
oxycephalum (Ditlevsen, 1926) Wieser, 1953
Paramicrolaimus
acanthus Jayasree & Warwick, 1977
Paramonohystera
- micramphis* Schuurmans Stekhoven, 1950
levicula (Lorenzen, 1973) Lorenzen, 1977
paranormandica (Micoletzky, 1922) Wieser, 1956
pellucida (Cobb, 1920) Wieser, 1956
riemanni (Platt, 1973)
megacephala (Steiner, 1916) Filipjev, 1918
elliptica (Filipjev, 1918) Wieser, 1956
Parapinnanema
harveyi Warwick & Coles, 1975
Pararaeolaimus
nudus (Gerlach, 1951) Timm, 1961
Parasaveljevia
clavicauda (Filipjev, 1927) Wieser, 1953
Parasphaerolaimus
paradoxus (Ditlevsen, 1918) Wieser, 1956
Paraterschellingia
brevicaudata (Kreis, 1924) Schuurmans Stekhoven, 1935
fusiformis Gerlach, 1951
Paratallassoalaimus
sabulicola Allgén, 1929
Paratricoma
papillifer Gerlach, 1956
pratensis Lorenzen, 1969
Paratuerkiana
comes (Tsalolikhin, 1972) Platonova, 1970
Pareudesmoscolex
laciniosus Lorenzen, 1969
papillosus (Schulz, 1935) Weischer, 1962
pratensis Lorenzen, 1969
Pareurystomina
armorica Luc & De Coninck, 1959
acuminata (de Man, 1889) Gerlach, 1952
filicaudata Allgén, 1934
flagellicaudata Schuurmans Stekhoven, 1946
parapagetinus Vitiello, 1970
tenuicauda Schuurmans Stekhoven, 1950
biserialis Schuurmans Stekhoven, 1946
scilloniensis Warwick, 1977
typica Micoletzky, 1930
tenuissima (Filipjev, 1927) Filipjev, 1946
Parodontophora
quadristicha (Schuurmans Stekhoven, 1950) Timm, 1963
breviseta (Schuurmans Stekhoven, 1950) Timm, 1963
Paroxystomina
asymmetrica Micoletzky, 1924
Pelagonema
omalum Vitiello, 1970
obtusicauda Filipjev, 1918
tenue Kreis, 1928
norvegicum Allgén, 1939
propinquum Allgén, 1929
simplex Cobb, 1894
angusticavatum Kreis, 1928
Perilinhomoeus
longisetosus Schuurmans Stekhoven, 1950
Phanoderm
cocksi Bastian, 1865
etha Gadea, 1960
laticolle (Marion, 1870) Marion, 1875
conicaudatum (Steiner, 1916) Steiner, 1921
annulocaudatum Allgén, 1939
parasiticum Ditlevsen, 1926
islandicum Ditlevsen, 1926
longisetum Allgén, 1939
tuberculatum (Eberth, 1863) Bastian, 1865
campbelli Allgén, 1928
brachyductum Wieser, 1953
rigidum Schuurmans Stekhoven, 1946
steineri Ditlevsen, 1918
nasutum Schuurmans Stekhoven, 1950
albidum Bastian, 1865
robustum Allgén, 1957
setigerum (Marion, 1870) Filipjev, 1918

- Phanodermella*
flagellicaudata Vitiello, 1970
- Phanodermopsis*
caudata Filipjev, 1927
reducta Filipjev, 1927
longicauda (Schuurmans Stekhoven, 1950) Filipjev, 1927
groenlandica Ditlevsen, 1926
- Pierrickia*
decasetaosa Vitiello, 1970
aequalis (Gerlach, 1956) Vitiello, 1970
- Platycoma*
cephalata Cobb, 1894
- Platycopomopsis*
nuchapilosa Vitiello, 1970
mesjatezvi Filipjev, 1927
effilata Schuurmans Stekhoven, 1946
cobbii Ditlevsen, 1926
- Polygastrophora*
alexandriae (Schuurmans Stekhoven, 1943) Wieser, 1953
hexabulba (Filipjev, 1918) Kreis, 1926
maior Schulz, 1932
attenuata de Man, 1922
- Polysigma*
fuscum Gerlach, 1956
- Pomponema*
loticum Lorenzen, 1972
tesselatum Wieser & Hopper, 1967
debile Lorenzen, 1972
cotylophorum (Steiner, 1916) Lorenzen, 1972
elegans Lorenzen, 1972
reductum Warwick, 1970
sedecima Platt, 1973
tautraense (Allgén, 1933) Lorenzen, 1972
lineatum (Gerlach, 1953) Wieser, 1959
cyatholaimoides (Allgén, 1929) Hope & Murphy, 1972
multipapillatum (Filipjev, 1922) Wieser, 1954
compactum Lorenzen, 1972
clavicaudatum (Schuurmans Stekhoven, 1935) Lorenzen, 1972
astrodes Lorenzen, 1972
ammophilum Lorenzen, 1972
- Pontonema*
ditlevensi (Kreis, 1934) Cobb & Steiner, 1934
vulgare (Bastian, 1865) Filipjev, 1916
macrolaimus (Southern, 1914) Filipjev, 1921
simile (Southern, 1914) Filipjev, 1921
brevicaudatoidea (Menzel, 1920) Gerlach & Riemann, 1974
parpapilliferum (Micoletzky, 1924) Cobb & Steiner, 1934
arcticum (Kreis, 1928) Cobb & Steiner, 1934
parocellatum Wieser, 1954
teissieri (Vitiello, 1967) Lorenzen, 1981
filicaudatus (Kreis, 1928) Lorenzen, 1981
zernovi (Filipjev, 1916) Cobb & Steiner, 1934
donsi (Allgén, 1932) Cobb & Steiner, 1934
papilliferum (Filipjev, 1916) Cobb & Steiner, 1934
balticum (Schulz, 1932) Cobb & Steiner, 1934
- Praeacanthonchus*
inglisi (Coles, 1965) Platt & Warwick, 1988
opheliae (Warwick, 1970) Gerlach & Riemann, 1973
kreisi (Allgén, 1929) Gerlach & Riemann, 1973
punctatum (Bastian, 1865) Micoletzky, 1924
- Procamacolaimus*
acer Gerlach, 1954
dolichostylum Gerlach, 1953
- Prochaetosoma*
primitivum (Steiner, 1916) Micoletzky, 1922
- Prochromadora*
exigua (Ditlevsen, 1928) Wieser, 1954
megodonta Filipjev, 1922
orleji (de Man, 1880) Filipjev, 1922
- magna* (Schulz, 1935) Wieser, 1954
spitzbergensis Gerlach, 1965
- Prochromadorella*
septempapillata Platt, 1973
neapolitana (de Man, 1876) Micoletzky, 1924
macrocellata Wieser, 1951
ditlevensi (de Man, 1922) Wieser, 1954
acridentata (Schulz, 1932) Wieser, 1954
actuaria Vitiello, 1971
ambigua (Ditlevsen, 1928) Wieser, 1954
brachyura Schuurmans Stekhoven, 1950
norwegica (Allgén, 1932) Wieser, 1951
attenuata (Gerlach, 1952) Lorenzen, 1972
maculata (Ditlevsen, 1918) Wieser, 1954
mediterranea (Micoletzky, 1922) Micoletzky, 1924
paramucrodonta (Allgén, 1929) Wieser, 1951
tenuicauda Gerlach, 1953
subterranea Gerlach, 1953
quinquepapillata (Schuurmans Stekhoven, 1935) Wieser, 1954
obtusidens (Schuurmans Stekhoven & Adam, 1931) Wieser, 1951
longicaudata (Kreis, 1929) Lorenzen, 1972
- Prooncholaimus*
eberthi (Filipjev, 1918) Micoletzky, 1924
megastoma (Eberth, 1863) Micoletzky, 1924
banyulensis Inglis, 1962
- Prosphaerolaimus*
eurypharynx Filipjev, 1918
- Protochromadora*
scampae (Coles, 1965) Inglis, 1969
mediterranea (Allgén, 1942) Inglis, 1969
- Pselionema*
simplex De Coninck, 1942
simile De Coninck, 1942
richardi De Coninck, 1942
minutum Vitiello & Haspeslagh, 1972
parasimplex Vitiello, 1971
annulatum (Filipjev, 1922) Cobb, 1933
deconincki Vitiello & Haspeslagh, 1972
longissimum Gerlach, 1953
longiseta Ward, 1974
- Pseudocella*
citronicauda (Wieser, 1954) Platonova, 1962
saveljevi (Filipjev, 1927) Platonova, 1962
gracilis Platonova, 1967
coeca (Saweljev, 1912) Filipjev, 1927
trichodes (Leuckart, 1849) Filipjev, 1927
cavernicola (Wieser, 1954) Platonova, 1962
elegans (Ditlevsen, 1926) Wieser, 1953
- Pseudolella*
norvegica Allgén, 1947
- Pseudometachromadora*
longilaima (Schuurmans Stekhoven, 1950) Timm, 1952
papillata (Schuurmans Stekhoven, 1950) Timm, 1952
- Pseudoncholaimus*
saveljevi (Filipjev, 1927) Kreis, 1934
- Pseudonchonus*
deconincki Warwick, 1969
northumbriensis Warwick, 1969
norwegianus Allgén, 1933
donsi Allgén, 1948
gerlachi Warwick, 1969
longus Allgén, 1949
- Pseudosteineria*
horrida (Steiner, 1916) Wieser, 1956
- Pterygonema*
platti Zhang, 1983
cambricensis Ward, 1973
- Ptycholaimellus*
monodon (Schuurmans Stekhoven, 1942) Hopper, 1969
ponticus (Filipjev, 1922) Gerlach, 1955
- Punctodora*

- ratzeburgensis* (Linstow, 1876) Filipjev, 1929
- Quadricoma*
frontalis (Gerlach, 1952) Timm, 1970
greeffii (Reinhard, 1881) Filipjev, 1922
loricata Filipjev, 1922
media (Reinhard, 1881) Filipjev, 1922
steineri Filipjev, 1922
suecica (Allgén, 1930) Timm, 1970
vittata (Lorenzen, 1969) Timm, 1970
trigintatres Timm, 1970
tenuis (Steiner, 1916) Filipjev, 1922
eurycricus (Filipjev, 1922) Timm, 1970
crassicauda Timm, 1970
cobbi (Steiner, 1916) Filipjev, 1922
scanica (Allgén, 1935) Timm, 1970
- Rhabdiitis*
marina Bastian, 1865
ehrenbaumi Schuurmans Stekhoven, 1935
- Rhabdocoma*
brevicauda Schuurmans Stekhoven, 1950
macrura Cobb, 1920
riemanni Jayasree & Warwick, 1977
- Rhabdodemania*
minor (Southern, 1914) Baylis & Daubney, 1926
major (Southern, 1914) Baylis & Daubney, 1926
mediterranea Boucher, 1971
obtusicauda (Allgén, 1954) Hope & Murphy, 1972
imer Warwick & Platt, 1973
pontica Platonova, 1965
laticauda (Ditlevsen, 1926) Baylis & Daubney, 1926
gracilis (Ditlevsen, 1918) Baylis & Daubney, 1926
coronata Gerlach, 1952
brevicaudata (Schuurmans Stekhoven, 1942)
 Lorenzen, 1981
- Rhadinema*
flexile Cobb, 1920
- Rhips*
paraornata Platt & Zhang, 1982
- Rhynchonema*
longituba Gerlach, 1953
deconincki Vitiello, 1967
lyngei (Allgén, 1940) Gerlach, 1953
scutatum Lorenzen, 1972
brevituba Gerlach, 1953
- Richtersia*
inaequalis Riemann, 1966
norvegica Allgén, 1940
elongata Schuurmans Stekhoven, 1950
demani Schuurmans Stekhoven, 1935
tenuis (Kreis, 1929) Schuurmans Stekhoven, 1935
collaris Steiner, 1916
- Sabatieria*
conicauda Vitiello, 1970
pulchra (Schneider, 1906) Riemann, 1970
stekhoveni Vitiello, 1970
strigosa Lorenzen, 1972
demani Schuurmans Stekhoven, 1935
dodecaspillata (Kreis, 1929) Wieser, 1954
granifer Wieser, 1954
elongata Jayasree & Warwick, 1977
vasicola Vitiello, 1970
lyonessa Warwick, 1977
pisinna Vitiello, 1970
armata Gerlach, 1952
macramphis Lorenzen, 1972
longispinosa Lorenzen, 1972
longisetosa (Kreis, 1929) Wieser, 1954
longicaudata Filipjev, 1922
breviseta Schuurmans Stekhoven, 1935
ornata (Ditlevsen, 1918) Filipjev, 1922
praedatrix de Man, 1907
abyssalis (Filipjev, 1918) Filipjev, 1922
celtica Southern, 1914
- Sarsonia*
leptosoma (de Man, 1893) Gerlach, 1967
- Saveljevia*
curvidens Filipjev, 1927
xiphonchus Filipjev, 1927
spissignatha (Allgén, 1940) Wieser, 1953
kolaensis Filipjev, 1927
cornuta Gerlach, 1956
- Scaptrella*
brevicaudata Gerlach, 1953
tenuicaudata Gerlach, 1956
- Setoplectus*
gerlachi Vitiello, 1971
- Setosabatieria*
hilarula (de Man, 1922) Platt, 1985
- Sigmophoranema*
monstrorum (Gerlach, 1956) Hope & Murphy, 1972
litorum (Schulz, 1938) Gerlach, 1951
rufum (Cobb, 1933) Hope & Murphy, 1972
- Siphonolaimus*
niger de Man, 1893
cylindricaudatus Schuurmans Stekhoven, 1942
cobbi Riemann, 1966
elongatus (Schuurmans Stekhoven, 1950) Vitiello, 1971
roervikensis Allgén, 1946
nigricans (Cobb, 1894) Filipjev, 1918
obtusicaudatus Allgén, 1930
weismanni (Zur Strassen, 1904) Steiner, 1916
ewensis Warwick & Platt, 1973
anticomoides (Allgén, 1933) Filipjev, 1918
- Solenolaimus*
obtusus Cobb, 1894
- Southerniella*
allometrica Vitiello, 1971
brevicauda (Schuurmans Stekhoven, 1942) Wieser, 1956
conicauda (Schuurmans Stekhoven, 1950) Wieser, 1956
zostericola (Allgén, 1929) Wieser, 1956
stenolaima (Steiner, 1916) Wieser, 1956
- Sphaerocephalum*
crassicauda Filipjev, 1918
chabaudi Inglis, 1962
bulbiferum Schuurmans Stekhoven, 1943
- Sphaerolaimus*
brevicollis Ssweljev, 1912
abyssorum Allgén, 1933
dispar Filipjev, 1918
glaphyrus Vitiello, 1971
latilaimus Allgén, 1958
macrocirculoides Wieser, 1954
paragracilis Vitiello, 1971
cuneatus Paramonov, 1929
minutus Vitiello, 1971
papillatus Kreis, 1929
balticus Schneider, 1906
islandicus Ditlevsen, 1926
paradoxus (Ditlevsen, 1918) Wieser, 1956
macrocirculus Filipjev, 1918
gracilis de Man, 1876
hirsutus Bastian, 1865
ostreae Filipjev, 1918
uncinatus Freudenhammer, 1970
polaris (Filipjev, 1946) Wieser, 1956
makrolasius Schulz, 1932
profundus Vitiello, 1971
striatus Allgén, 1935
tenuis (Kreis, 1924) Schuurmans Stekhoven, 1935
maeoticus Filipjev, 1922
pacificus Allgén, 1947
- Spiliphera*
dolichura de Man, 1893
hirsuta Gerlach, 1956
gracilicauda de Man, 1893
spinicauda Vitiello, 1970
minuta Vitiello, 1972

- elegans* Bastian, 1865
Spilophorella
papillata Kreis, 1929
paradoxa (de Man, 1888) Filipjev, 1917
euxina Filipjev, 1918
candida Gerlach, 1951
Spiramphinema
microcephalon (Schuurmans Stekhoven, 1942)
Wieser, 1956
Spirinia
laevis (Bastian, 1865) Gerlach, 1963
flagellata (Vitiello, 1971) Lorenzen, 1981
schneideri (Villot, 1875) Gerlach, 1963
parasitifera (Bastian, 1865) Gerlach, 1963
granulata (Allgén, 1929) Gerlach, 1963
gerlachi (Luc & De Coninck, 1959) Gerlach, 1963
sabulicola (Filipjev, 1918) Gerlach, 1963
Steineria
longicaudata Vitiello, 1971
polychaetoides Gerlach, 1951
pontica Groza-Rojancovski, 1972
pilosa (Cobb, 1914) Micoletzky, 1921
Steineridora
loricata (Steiner, 1916) Inglis, 1969
adriatica (Daday, 1901) Inglis, 1969
Stephanolaimus
paraflevensis Gerlach, 1953
jayasreei Platt, 1983
filicaudatus Schuurmans Stekhoven, 1946
spartinae Lorenzen, 1969
flevensis Schuurmans Stekhoven, 1935
Symplocostoma
reisingeri Ditlevsen, 1928
medium Filipjev, 1927
tenunicolle (Eberth, 1863) Wieser, 1953
leptolaimus Filipjev, 1927
murmanicum Ssaweljev, 1912
majus Filipjev, 1927
karensis Filipjev, 1927
acutum (Cobb, 1920) Wieser, 1953
ponticum Filipjev, 1918
papillatum Kreis, 1928
Synodontium
impar (Ssaweljev, 1912) Wieser, 1956
Synonchiella
micramphis (Schuurmans Stekhoven, 1950) Gerlach, 1964
riemanni Warwick, 1970
minuta Vitiello, 1970
Synonchium
depressum Gerlach, 1954
obtusum Cobb, 1920
Synonchus
armatus (Ditlevsen, 1923) Filipjev, 1927
intermedius Allgén, 1939
longisetosus (Southern, 1914) Filipjev, 1927
fasciculatus Cobb, 1894
acuticaudatus (Jägerskjöld, 1901) Filipjev, 1916
dubius (Wieser, 1954) Gerlach & Riemann, 1974
roscoffianus (Villot, 1875) Filipjev, 1927
murmanicus Filipjev, 1927
Synonema
triadenophorum (Steiner, 1921) Wieser, 1954
obtusicaudatum (de Man, 1889) Gerlach, 1963
Syringolaimus
filicaudatus Vitiello, 1970
striatocaudatus de Man, 1888
brevicaudatus Micoletzky, 1922
caspersi Gerlach, 1951
Syringonomus
typicus Hope & Murphy, 1969
Tarvaiia
angusta Gerlach, 1953
donsi Allgén, 1934
Terschellingia
communis de Man, 1888
longicaudata de Man, 1907
gourbaultae Austen, 1989
pontica Filipjev, 1918
similis Allgén, 1933
parva Vitiello, 1969
capitata Vitiello, 1969
Thalassironus
britannicus de Man, 1889
Thalassoalaimus
egregius Steiner, 1916
septentrionalis Filipjev, 1927
rotundicaudatus Filipjev, 1927
longicaudatus Vitiello, 1970
brevicaudatus Vitiello, 1970
tenuis Allgén, 1935
oxycauda Filipjev, 1927
paramontemari Vitiello, 1970
mediterraneus Vitiello, 1970
pirum Lorenzen, 1969
filiformis Allgén, 1933
aceratus Vitiello, 1970
tardus de Man, 1893
Thassomonhystra
venusta (Lorenzen, 1979) Jacobs, 1987
parva (Bastian, 1865) Jacobs, 1987
Theristus
maior Gerlach, 1956
meyli Riemann, 1966
pellucidus Allgén, 1939
monstrosus Gerlach, 1954
norwegicus (Allgén, 1933) Wieser, 1956
pyronasi Gerlach, 1965
velox (Bastian, 1865)
obtusicephalus Schuurmans Stekhoven, 1950
lineatus Gerlach, 1965
scanicus Allgén, 1949
flevensis Schuurmans Stekhoven, 1935
pictus Gerlach, 1951
littoralis Filipjev, 1922
leptosoma Allgén, 1950
heterospiculoides Gerlach, 1953
inermis Gerlach, 1953
anisotrichus Lorenzen, 1972
tenuicaudatus Allgén, 1951
strandi Allgén, 1934
roscoffiensis Vitiello, 1967
ensifer Gerlach, 1951
gracilis (de Man, 1876) Filipjev, 1918
parasetosus (Allgén, 1928)
latissimus Filipjev, 1922
heterospiculum (Allgén, 1932) Gerlach, 1953
interstitialis Warwick, 1970
longus Platt, 1973
otoplanobius Gerlach, 1951
pertenuis Schuurmans Stekhoven, 1935
sabulicola (Filipjev, 1918) Wieser, 1956
stichotricha Schuurmans Stekhoven, 1950
bipunctatus (Schneider, 1906) Filipjev, 1929
complexus Jaysree & Warwick, 1977
longisetosus (De Coninck & Schuurmans Stekhoven, 1933) Lorenzen, 1977
aculeatus Schulz, 1935
blandicor Rachor, 1971
bastiani Gerlach & Riemann, 1973
acer Bastian, 1865
acriformis (de Man, 1922) Gerlach & Riemann, 1973
denticulatus Warwick, 1970
Thoracostoma
steineri Micoletzky, 1922
zolae Marion, 1870
penduli Allgén, 1954
multipapillosum Allgén, 1957
amphacanthus (Grube, 1855) Bütschli, 1874
brachylobatum Allgén, 1954

- sivertensi* Allgén, 1957
filiformis (Eberth, 1863) Filipjev, 1921
hispanicum Platonova, 1970
coronatum (Eberth, 1863) Marion, 1870
- Thoracostomopsis*
galeata Filipjev, 1927
ditlevseni Filipjev, 1927
barbata Ditlevsen, 1918
doveae Warwick, 1970
longissima Filipjev, 1927
- Trefusia*
litoralis (Allgén, 1932) De Coninck & Schuurmans Stekhoven, 1933
silicauda Allgén, 1933
zostericola Allgén, 1933
helgolandica Riemann, 1966
longicauda de Man, 1893
pseudilitoralis Vitiello, 1970
filum Schuurmans Stekhoven, 1942
- Trichethimolaimus*
hirsutus (Gerlach, 1956) Platt, 1982
- Trichotheristus*
mirabilis (Schuurmans Stekhoven & De Coninck, 1933) Wieser, 1956
psammoides (Warwick, 1970)
- Tricoma*
brevirostris (Southern, 1914) Steiner, 1916
longirostris (Southern, 1914) Steiner, 1916
suecica Allgén, 1930
strandii (Allgén, 1939) Gerlach, 1958
steineri de Man, 1922
septuaginta Schuurmans Stekhoven, 1942
profunda (Schepotieff, 1907) Stauffer, 1924
pellucida Allgén, 1935
maxima (Schepotieff, 1907) Allgén, 1942
islandica Kreis, 1963
bacescui (Paladian & Andriescu, 1963) Timm, 1970
bergensis (Schepotieff, 1907) Filipjev, 1922
platycephala Filipjev, 1922
elongata (Panceri, 1876) Stauffer, 1924
polydesma (Southern, 1914) Steiner, 1916
litoralis (Allgén, 1934) Timm, 1970
norvegica (Schepotieff, 1907) Stauffer, 1924
nematoides (Greeff, 1869) Stauffer, 1924
adelpha (Greeff, 1869) Stauffer, 1924
cincta Cobb, 1894
- Trileptium*
subterraneum (Gerlach, 1952) Cobb, 1933
parisetum Warwick & Platt, 1973
- Triodontolaimus*
acus (Villot, 1875) de Man, 1893
- Tripyla*
cornuta Skwarra, 1921
setifera Bütschli, 1873
glomerans Bastian, 1865
- Tripyloides*
gracilis (Ditlevsen, 1918) Filipjev, 1927
omblaica Micoletzky, 1924
marinus (Bütschli, 1874) de Man, 1886
- Trissonchulus*
benepapillosus (Schulz, 1935) Gerlach & Riemann, 1974
oceanus Cobb, 1920
obtusus (Schuurmans Stekhoven, 1935) Inglis, 1961
- Trochamus*
carinatus Boucher & Bovée, 1972
- Tubolaimoides*
tenuicaudatus (Allgén, 1934) Gerlach, 1963
- Valvaelaimus*
euximus (Filipjev, 1918) Lorenzen, 1977
- Vasculonema*
caeruleum (Eberth, 1863) Kreis, 1928
conicaudatum Kreis, 1928
- Viscosia*
minudonta Vitiello, 1970
- separabilis* (Wieser, 1953) Lorenzen, 1981
stenolaima Filipjev, 1927
longidentatus (Schuurmans Stekhoven, 1931) Lorenzen, 1981
brevidentatus (Vitiello, 1967) Lorenzen, 1981
elegans (Kreis, 1924) Lorenzen, 1981
parva Kreis, 1929
palmae Schuurmans Stekhoven, 1942
profundus (Vitiello, 1970) Lorenzen, 1981
langrunensis (de Man, 1890) Filipjev, 1918
brachylaima Filipjev, 1927
elongata Filipjev, 1922
rusticus (Kreis, 1929) Lorenzen, 1981
papillosa (Eberth, 1863) Kreis, 1934
viscosa (Bastian, 1865) de Man, 1890
minor Filipjev, 1918
glabra (Bastian, 1865) de Man, 1890
abyssorum (Allgén, 1933) Wieser, 1953
filipjevi Paramonov, 1929
macramphida Chitwood, 1951
macrorhopalocerca Kreis, 1932
megalaima (Ditlevsen, 1928) Hope & Murphy, 1972
cobbi Filipjev, 1918
diodon (Wieser, 1951) Lorenzen, 1981
- Wieseria*
glandulosa (Kreis, 1929) Gerlach, 1956
clavata Gerlach, 1956
pica Gerlach, 1956
hispida Vitiello, 1972
inaequalis Gerlach, 1956
leptura Vitiello, 1972
longiseta (Allgén, 1947) Gerlach, 1956
- Xennella*
suecica Allgén, 1935
- Xyla*
striata Cobb, 1920
longicaudata Ward, 1972
barbata Ward, 1972
riemannii Boucher & Hellouët, 1977

Nematoda (parasitic)

Compiled by David I. Gibson

The list of parasitic nematode species was derived from the Host-Parasite Data-base and Catalogue maintained by the Parasitic Worms Division, Department of Zoology, The Natural History Museum. Some attempt has been made to restrict the list to valid taxa and to use that latest nomenclature. However, the many, usually unrecognisable and often larval, names listed under *Ascaris* have been listed separately.

The list was kindly commented upon by Dr F. Moravec and Dr H.-P. Fagerholm.

NEMATODA, PARASITIC

- Acanthocheilus*
rotundatus (Rudolphi, 1819)
- Agamascaris*
septentrionalis Kreis, 1952
- Agamoneema*
 sp.
- Amidostomum*
acutum (Lundahl, 1848)
- Anguillicola*
crassus Kuwahara, Niimi & Itagaki, 1974
- Anisakis*
pegreffii Campana-Rouget & Biocca, 1955
physeteris (Baylis, 1923)
schupakovi Mosgovoi, 1951
simplex (Rudolphi, 1809)
simplex C of Nascetti *et al.* (1986)
typica (Diesing, 1860)
ziphidarum Paggi *et al.*, 1988

- Ascarophis*
arctica Poljansky, 1952
capelanus Nikolaeva & Naidenova, 1964
crasicollis Dollfus & Campana-Rouget, 1956
filiformis Poljansky, 1952
morrhuae van Beneden, 1871
nullus Naidenova & Nikolaeva, 1968
- Bulbocephalus*
pseudopenei Vassiliades & Diaw, 1978
- Camallanus*
lacustris (Zoega, 1776)
truncatus Rudolphi, 1814
- Capillaria*
anatis (Schrank, 1790)
gracilis (Bellingham, 1840)
wickinsi Ogden, 1965
- Contracaecum*
filiforme (Stossich, 1904)
mulli (Wedl, 1855)
osculatum (Rudolphi, 1802)
papilligerum (Creplin, 1846)
pedum (Deslongchamps, 1824)
rudolphii Hartwich, 1964
septentrionale Kreis, 1955
- Cosmoccephalus*
obvelatus (Creplin, 1825)
- Crassicauda*
anthonyi Chabaud, 1962
boopis Baylis, 1920
carbonelli Raga *et al.*, 1988
grampicola Johnston & Mawson, 1941
- Cristitectus*
congeri Petter, 1970
- Cucullanus*
attenuata (Molin, 1858)
bioccai Orecchia & Paggi, 1987
campanae Lebre & Petter, 1984
carbonelli Campos, Carbonell & Rodriguez-Babio, 1993
cirratus Müller, 1777
dodsworthi Barreto, 1922
dogieli Krotas, 1959
elongatus Smedley, 1933
gendrei Campana-Rouget, 1957
heterochrous Rudolphi, 1802
hians (Dujardin, 1845)
longicollis (Stossich, 1899)
longispiculum De Oliveira Rodrigues *et al.*, 1973
lophii Campana-Rouget & Chabaud, 1956
micropapillatus Tornquist, 1931
mugili Belous, 1965
murenophidis Campana-Rouget, 1957
orthagorisci (Rudolphi, 1819)
parvus Tornquist, 1931
praecinctus (Dujardin, 1845)
pulcherrimus Barreto, 1918
rotundata (Molin, 1859)
truttae Fabricius, 1794
- Cystidicola*
farionis Fischer, 1798
- Cystoopsis*
scomber Zlatev, 1936
- Desmidocercella*
sp.
- Dichelyne*
abbreviatus (Rudolphi, 1819)
adriaticus (Tornquist, 1931)
elongatus (Tornquist, 1931)
mauritanicus (Gendre, 1928)
minutus (Rudolphi, 1819)
pomadasys Vassiliades & Petter, 1981
tripapillatus (Gendre, 1927)
- Dipetalonema*
spirocauda (Leidy, 1858)
- Echinocephalus*
spinosisimus von Linstow, 1905
uncinatus Molin, 1858
- Echinomermella*
matsi Jones & Hagen, 1987
- Eustrongylides*
sp.
- Fellicola*
longispiculus Petter & Køie, 1993
- Goezia*
annulata (Molin, 1859)
ascaroides (Goeze, 1782)
kollari (Molin, 1859)
tricirrata Osmanov, 1940
- Halocercus*
delphinii (Baylis & Daubney, 1925)
kleinenbergi Delamure, 1951
lagenorhynchi Baylis & Daubney, 1925
ponticus Delamure, 1946
taurica Delyamure, 1942
- Heliconema*
heliconema Travassos, 1919
- Heligmus*
longicirrus Dujardin, 1845
- Hysterothylacium*
aduncum (Rudolphi, 1802)
arnoglossi Petter & Maillard, 1988
auctum (Rudolphi, 1802)
bifidalatum Petter & Maillard, 1988
collare (Cobb, 1929)
cornutum (Stossich, 1904)
corrugatum Deardorff & Overstreet, 1981
cyclopteri (Kreis, 1952)
fabri (Rudolphi, 1819)
increscens (Molin, 1858)
incurvum (Rudolphi, 1819)
legendrei (Dollfus, 1933)
reliquens (Norris & Overstreet, 1975)
rhacodes (Deardorff & Overstreet, 1978)
rigidum (Rudolphi, 1809)
- Ichthyaniakis*
monodi Gendre, 1928
- Ichthyascaris*
mediterraneus (Lebre & Petter, 1983)
- Ichthyobronema*
hamulatum (Moulton, 1931)
- Ichthyofilaria*
bergensis (Wulker, 1930)
- Johnstonmawsonia*
murenophidis (Campana-Rouget, 1955)
- Kathlania*
leptura (Rudolphi, 1819)
- Mawsonascaris*
laymani (Mozgovoy, 1950)
pastinaceae (Rudolphi, 1819)
- Onchophora*
melanocephala (Rudolphi, 1819)
- Paracapillaria*
gibsoni Moravec, 1987
- Paracuaria*
adunca (Creplin, 1846)
tridentata (Linstow, 1877)
- Paranisakiopsis*
coelorhynchi Yamaguti, 1941
- Paranisakis*
monodi (Gendre, 1928)
squatinae Baylis, 1923
- Parascarophis*
sphyraeae Campana-Rouget, 1955
- Philometra*
globiceps (Rudolphi, 1819)
obturans (Pennant, 1886)
onchorhynchi Kuitinen-Ekbaum, 1933
ovata (Zeder, 1803)
rischta Skrjabin, 1917
rubra (Leidy, 1856)