
Lipid Composition Modifications in the Blue Mussels (*Mytilus edulis* L.) from the White Sea

Natalia N. Fokina, Tatiana R. Ruokolainen and
Nina N. Nemova

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67811>

Abstract

Studying biochemical indicators in response to various environmental factors allows revealing the metabolic adaptive strategy of the organism's tolerance and survival under a variety of environmental impacts. This review analyses both the authors' own data and the available literature on the problem of biochemical adaptations of the lipid composition in marine bivalves, particularly blue mussels, *Mytilus edulis* L., to various environmental impacts. Modifications in the composition of lipids and their fatty acids in blue mussels caused by short-term (under laboratory conditions) and chronic (field monitoring) exposure to natural and human factors indicate that homeostasis is maintained in cell membranes and the organism's energy requirements and facilitate the adaptation and tolerance of the mussels to environmental disturbances. The lipid and fatty acid composition indices in White Sea intertidal mussels which reflect their chronic exposure to a wide variety of environmental factors are discussed and compared to data on changes in the lipid composition of blue mussels exposed to some environmental factors (salinity, anoxia, metals) in aquarium experiments. The lipid profile plays an important role in the adaptation of blue mussels to new conditions in the habitat, and it can be used as a biochemical marker for indicating the organism's physiological state.

Keywords: lipids, fatty acids, biochemical adaptation, environmental factors, *Mytilus edulis*

1. Introduction

Biochemical processes underlie the development of cell metabolic responses to environmental impacts and allow an organism to adapt and survive in a changing environment [1]. Metabolic modifications up to the level of physiological and morphological disorders are reflected in the

changes of various biochemical indicators, which allow determining the adaptive strategy of an organism's tolerance and survival under both natural and human impacts. Lipid molecules, which are involved in all the essential physiological-biochemical processes [2], play a major role in the organism's adaptive responses to various factors in the environment [1]. The primary response to stress is modification of the physical state of cell membranes (mainly fluidity), which triggers a change of their lipid and fatty acid composition [3, 4]. The main lipid components of biological membranes are phospholipids and cholesterol. The ratio of phospholipids and cholesterol is considered as an indicator of membrane fluidity. Cholesterol is known to increase the order of the phospholipid fatty acid chains in membranes [5]. Membrane phospholipids in different molecular species and molecular shapes as well as their interaction with cholesterol and membrane proteins determine membrane fluidity and subsequently regulate the activity of membrane-bound enzymes and the functioning of ion channels, pumps and receptors [2, 6, 7]. Besides their effects on membrane fluidity, membrane phospholipids are also a source of bioactive compounds and messengers [8]. In particular, eicosapentaenoic acid (EPA, 20:5n-3) and arachidonic acid (AA, 20:4n-6) are released from phosphatidylcholine (PC), phosphatidylethanolamine (PE) and phosphatidylinositol (PI) by phospholipase A2 and serve as precursors of short-lived hormone-like substances called eicosanoids (prostaglandins, thromboxanes, leukotrienes, etc.). The bioactive molecules have a wide range of physiological actions, including immune response, inflammatory response, neural function, reproduction and enhancement of an organism's adaptation to environmental stress [2, 8]. PI is also a source of such messengers as diacylglycerols and inositol phosphates (namely, inositol trisphosphate and others). These messengers, as well as the phospholipid phosphatidylserine (PS), are involved in regulating the activity of protein kinase C, which controls many cell functions, such as differentiation, proliferation, metabolism and apoptosis [2, 8, 9]. Moreover, fatty acids are the most labile components of lipid molecules, quickly and accurately reflecting environmental impacts and activating an organism's adaptive abilities. For example, a well-known biochemical response of poikilothermic organisms to low temperature is increased fatty acid unsaturation of both membrane and storage lipids [7, 10, 11]. In addition to membrane lipids, an important role in the adaptive response of organisms to various environmental factors belongs to high-energy storage lipids, chiefly triacylglycerols and their fatty acids [12–15], which cover the energy costs needed for maintaining homeostasis under the new environmental conditions. Since long-chain polyunsaturated fatty acids, particularly such essential fatty acids as EPA and docosahexaenoic acid (DHA, 22:6n-3), cannot be de novo synthesized in marine mussels [16, 17], their incorporation and elimination in membrane and storage lipids are strongly regulated [18, 19]. Thus, lipid and fatty acid composition as a key component of various metabolic pathways that are linked to processes important for survival and tolerance reflects the adaptive response of an organism to environmental effects. It is assumed that lipid composition may be used as a biochemical marker for indicating the organism's physiological state in environmental assessments and biomonitoring.

The blue mussels, *Mytilus edulis* L., are used worldwide as marine sentinel organisms in biomonitoring programmes due to their longevity, sessile nature, global distribution and ability to bioaccumulate high concentrations of pollutants [20–22]. In the White Sea, *M. edulis* L. is the dominant species of coastal (intertidal) ecosystems. Numerous studies on White Sea mussels' response to various environmental effects have identified adaptive mechanisms on molecular, biochemical, cellular, physiological and behavioural levels of biological organization [23–38].

This paper summarizes the results of research on lipid composition effects in White Sea blue mussels, *M. edulis* L. (1758), in response to environmental factors such as temperature, salinity, short-term anoxia, change of nutrition source, metals and oil pollution. Phenotype-specific features of the lipid composition in White Sea blue mussels from different habitat conditions (intertidal zone and aquaculture), as well as compensatory modifications of the lipid composition in intertidal mussels under chronic stress in the natural habitat and under short-term exposure to stress in laboratory experiments, are discussed.

2. Environmental factors

2.1. Temperature

Being a major environmental factor, influences all aspects of life of an organism, especially poikilotherms [1]. The literature offers quite detailed descriptions of the contribution of the lipid composition to thermal adaptations in bivalves [11, 39–43]. Often, the response involving the lipid composition depends on the duration of exposure to ambient temperatures. Moreover, organ-specific distribution of lipids in Bivalvia causes differences in lipid composition response to temperature effects, depending on the studied organ. Thus, gills of bivalves, which are the location of primary contact with the environment, contain high concentrations of membrane lipids, chiefly cholesterol [42]. It is well-known cholesterol is necessary for membrane stabilizing and maintaining the permeability of membranes [2, 5]. It was demonstrated that when exposed to rapid (several hours) temperature fluctuations gills of bivalves experience modifications in cholesterol levels, whereas prolonged (several weeks) temperature impacts induce changes in the amount of phospholipids enriched in polyunsaturated fatty acids [41, 43]. Whereas the gill lipid composition response in White Sea mussels in the temperature experiment was the opposite: a significant rise of the cholesterol concentration in response to prolonged (14 days) impact of both low and high temperatures, while short-term temperature stress (1 day) influenced the content of phospholipids and their fatty acids [36]. Since the synthesis of polyunsaturated fatty acids in bivalves is limited as well as an involvement of essential fatty acids in/from membrane lipids is strictly regulated [18, 19], presumably, the mussels' adaptive strategy is to use a less energy-intensive mechanism for maintaining optimal membrane fluidity by redistributing polyunsaturated fatty acids among storage and membrane lipid fractions (in the absence of additional cholesterol synthesis) that bivalves employ to adapt to rapid temperature changes [35, 36]. The role of minor membrane phospholipids (namely, phosphatidylserine and sphingomyelin) in the acclimation of mussels to elevated ambient temperature was also demonstrated in gills. They are believed to facilitate adaptive modifications of the fluidity and permeability of cell membranes in response to elevation of the ambient temperature [35, 36]. Let us remark that a similar effect involving these phospholipids was observed in gills of mussels acclimating to variable seawater salinity [44]. Elevated ambient temperature is known to produce a destabilizing effect on cell membranes in poikilotherms [3]. Apparently, seawater salinity variations, primarily reduction of salinity, cause an analogous response of the membrane physicochemical properties which, in turn, initiates compensatory modifications of the composition of membrane

lipids and their fatty acids similar to those observed in response to elevated temperature. Digestive glands in bivalves contain higher concentrations of triacylglycerols (TAGs) required for energy metabolism during their acclimation to new environmental conditions as well as for reproductive processes. In particular, it was shown that bivalves from thermally different habitats differed in TAG metabolism in the digestive glands during overwintering. Blue mussels, *M. edulis*, adapted to harsh winters, accumulate TAGs enriched in 20:5n-3 fatty acids (EPA) in the digestive glands, while the oyster, *Crassostrea virginica*, which generally occurs in warmer habitats, on the contrary, did not reserve TAG before overwintering [42].

2.2. Salinity

Is one of the key abiotic factors in the marine environment. Marine and freshwater molluscs were shown to be alike in the content of total and neutral lipids, but the levels of individual fractions of phospholipids such as PC, PI, PS and PE, as well as PE plasmalogen form, were found to be dependent on the ambient salinity [45]. In gills of bivalves, salinity stress induces an increase in negatively charged phospholipids, chiefly PI and cardiolipin. This may supposedly be one of the intracellular mechanisms to bind excessive cations lest they degrade the cell's enzymatic systems [46]. Exposure of White Sea mussels to reduced seawater salinity (15 psu) in aquarium experiments (25 psu as a control) resulted in an increase in the concentration of phospholipids, mainly PC and PS, in gills [44], alongside a reduction in the levels of cholesterol and storage lipids (triacylglycerols and cholesterol esters) in gills and digestive glands [34, 47]. The reduced levels of cholesterol and storage lipids indicate an adverse effect of low salinity on the physiological state of the mussels and their metabolism. We know that when salinity is drawn down from 25 psu (normal values of salinity in the White Sea) to 14 psu, the functional activity of *M. edulis* is suppressed [23, 48], and the mussels' tolerance of low salinity is ensured by cellular volume regulation using organic and inorganic osmolytes [32, 49, 50]. Thus, the acclimation of molluscs to reduced salinity apparently implies that storage lipids are utilized not only as sources of metabolic energy but also as substrates for the synthesis of organic osmolytes [34]. Moreover, it was shown that the action of increased salinity (from 25 psu to 35 and 45 psu) leads to various responses of the lipid composition of White Sea mussels, *M. edulis*, depending on the studied organ. Although mantle edge and gills are sites for primary contact with external environment, the effect of increased seawater salinity on both intertidal and cultured mussels caused organ-specific reactions in the cholesterol level: the level rose in the mantle edge [51] but declined in gills [44]. Apparently, the different cholesterol content in gills and mantle edge of the mussels reflects differences in membrane fluidity and ion permeability in response to increased salinity effect. Some authors have pointed out the lack of distinctions between marine and freshwater bivalves in the fatty acid composition [45, 52, 53], although some papers have reported elevated concentrations of C20 and C22 unsaturated fatty acids, predominantly 20:5n-3, 22:5n-3 and 22:6n-3, in marine molluscs [54] as well as high level of monounsaturated fatty acids and arachidonic acid (20:4n-6) in freshwater molluscs [55]. Yet, the high variability of fatty acid content observed in both freshwater and marine mollusc species is primarily due to the factors of nutrition and ambient temperature [14, 15, 41–43, 45, 56–58]. Nonetheless, a lower level of n-6 polyunsaturated fatty acids was found in gills of marine bivalves acclimated to high salinity as compared to the individuals exposed to low salinity [46]. White Sea mussels acclimated to different seawater salinities also manifested considerable modifications of the lipid

fatty acid composition in gills and mantle edge. Thus, in gills of intertidal mussels, the level of n-6 polyunsaturated fatty acids (mainly owing to AA, 20:4n-6) increased in response to seawater salinity reduction (5 psu) and elevation (45 psu). At the same time, cultured mussels collected from aquaculture substrates responded with a decrease in n-3 polyunsaturated fatty acid (PUFA) content and an increase in saturated fatty acid level in gills both to a reduction (to 5 and 15 psu) and an elevation (to 35 and 45 psu) of seawater salinity. Remarkably, notwithstanding the considerably different lipid composition of gills in intertidal and aquaculture mussels, they both responded to critically low salinity (5 psu) with similar modifications of the lipid composition, indicative of non-specific defence reaction in bivalves—closure of shell valves, reduction of total metabolism and transition to anaerobic metabolic pathways [44]. Varied response of fatty acid composition to salinity effects was detected in the mantle edge of mussels from different tidal zones (intertidal and aquaculture) [51]. Thus, it was shown that the concentration of non-methylene-interrupted fatty acids (NMIFA) in mantle edge increased in the intertidal mussels exposed to 5, 35 and 45 psu (25 psu as a control), whereas in cultured mussels exposed to 5, 35 and 45 psu salinity, there was an increase in n-3 PUFA content. It is known that NMIFA can be synthesized in marine bivalve molluscs in the case of a lack of usual n-3 PUFA [53]. Probably, n-3 PUFA deficiency in intertidal mussels is the result of their utilization to generate energy required for mussel acclimation to different salinities, whereas additional synthesis of NMIFA is essential for maintaining the unsaturated state of membrane phospholipids as well as fluidity and permeability of membranes in mantle edge.

2.3. Short-term anoxia

Blue mussels, *M. edulis*, living in the marine coastal (intertidal) zone are facultative anaerobes tolerant of short-term anoxia during low tide [30, 59, 60]. The main sources of energy for bivalves in the anaerobic metabolism conditions are glycogen and proteins [1, 61], whereas lipids are utilized to provide for gametogenesis [61]. The role of storage lipids (chiefly triacylglycerols) in the adaptive reactions of mussels under anoxic conditions was demonstrated in our studies [62, 63]. In White Sea blue mussels, we observed a rise in the levels of cholesterol and PC within total lipids of soft tissues, which are known to have a stabilizing effect on membranes and thus reduce their permeability. It is known that anoxia may reduce the permeability of cell membranes, thus causing modifications in their lipid composition [64]. On the other hand, elevated concentrations of polyunsaturated fatty acids (in particular, arachidonic acid) and non-methylene-interrupted fatty acids within total lipids of soft tissues balance the stabilizing effect of membrane lipids and probably facilitate the functioning of membrane-bound proteins (enzymes, ion channels and receptors) [63]. Additional research is needed to determine organ-specific reaction of the lipid and fatty acid composition in blue mussels under short-term anoxia effect.

2.4. The nutrition factor

It is a known fact that lipid composition, especially the fatty acid profile of filter-feeding mussels, like in any consumers, is a trophic marker of the composition of their food and includes the biochemical markers of all seston components, namely, phytoplankton, zooplankton and bacteria (detritus) [14, 15, 56, 58, 65–68]. The study of modifications in the composition of lipids and their fatty acids in gills and digestive glands of White Sea mussels, *M. edulis*, induced by

their acclimation to laboratory conditions where they were fed with artificial feed ("Coraliquid", Sera) revealed organ-specific patterns in the assimilation and modification of lipids, primarily concerning their fatty acid profile [69]. It was shown that change of the food source caused alterations in the lipid and fatty acid composition, mainly in the digestive gland. Elevated content of high-energy lipids (triacylglycerols) enriched in saturated fatty acids (namely, myristic 14:0 acid), as well as vaccenic 18:1n-7 acid in the feed, promoted the storage of these lipids in mussel gills and digestive gland. At the same time, the fact that phospholipids in the feed lacked essential fatty acids, EPA and DHA, which are known to be derived from phytoplankton, told considerably on the fatty acid profile of triacylglycerols in the mussels [69].

Intertidal habitats are the most variable in terms of such abiotic environmental factors as temperature, salinity, aerial exposure and concentrations of suspended nutritive material [70]. Life under such heavily variable environmental conditions reflects both on the mussels' physiological (including growth rate) and metabolic processes and on the size and age structure of mussel beds (local populations), their abundance and biomass [27, 29, 30, 71]. Ecologo-biochemical monitoring during 2009–2014 years of two intertidal mussel beds located in different parts of the Gulf of Kandalaksha, White Sea, and differing in hydrological characteristics, including seawater salinity, revealed the features of the lipid composition in gills and digestive glands which reflect the chronic impact of salinity drops [38]. Frequent salinity drops in one of the investigated sites (Site 1) are due not only to discharge from streams but also to human activities (namely, unregulated freshwater discharges from hydropower plant). The effect of reduced seawater salinity (approximately 9.7–14.0 psu) on mussels from Site 1 appears not only in the level of some phospholipids (in particular, phosphatidylserine, phosphatidylethanolamine and phosphatidylcholine), and the ratio of n-3/n-6 polyunsaturated fatty acids in the molluscs' gills and digestive gland, but also in some ecological characteristics of the mussel beds, i.e. its size-age structure, abundance and biomass [38]. The elevated content of the named phospholipids and the prevalence of n-3 polyunsaturated fatty acids over n-6 polyenes apparently serve to keep membranes permeable to ions and maintain the functioning of membrane-bound enzymes involved in cellular volume regulation in response to low seawater salinity. These data agree with the results of previous aquarium experiments on the effect of low seawater salinity on the lipid composition of intertidal and cultured mussels [44], which suggest that the mussels' lipid and fatty acid composition is adapted to secure the survival of the molluscs under low seawater salinity. The elevated ratio of n-3/n-6 polyunsaturated fatty acids in the mussels chronically exposed to salinity drops may also be a result of high metabolic rate of n-6 polyunsaturated fatty acids (chiefly AA, 20:4n-6). Arachidonic acid is a precursor for the synthesis of physiologically active hormone-like molecules, eicosanoids (such as prostaglandins), which are known to build up bivalves' resistance to stress, including various seawater salinities [8, 72–75]. One must mention that intertidal mussels living in a habitat with relatively stable salinity conditions (away from freshwater discharges, Site 2 where seawater salinity is 20.1–22.5 psu) feature an elevated content of cholesterol and n-6 polyunsaturated acids within total lipids of both gills and digestive glands [38]. Our monitoring studies of the lipid and fatty acid composition in intertidal mussels from different habitats in the White Sea showed that the fatty acid composition of digestive glands, unlike their content in gills, reflects the adaptive features of the lipid metabolism in the mussels under chronic effect of a wide range of environmental factors [38].

There is a lot of research on the study of the differences in physiological (energetic, growth rate, clearance rate, ingestion rate, absorption rate, respiration rate) and biochemical indices between mussels collected in intertidal (rocky shore) and subtidal (aquaculture) environments [13–15, 76–78]. It is considered that these origin-related differences in physiological rates have to do with features of the energy distribution, namely, in intertidal mussels more energy is directed to the formation of a thicker shell, while in subtidal mussels the energy is spent on tissue growth [78]. Simultaneously, biochemical differences between these mussel groups are associated with various concentrations and quality of seston in the intertidal and subtidal zones [13, 14, 76]. In particular, frequent exposure to air (during low tide) has very high effect on the mussels' energy reserves including triacylglycerols, saturated fatty acids and some polyunsaturated fatty acids, similarly to the effect of starvation [13, 14]. We have also studied origin-related lipid composition differences in gills of littoral (intertidal) and cultured (sublittoral) mussels after 2 weeks of acclimation to laboratory conditions [44, 79]. It was demonstrated that gills of intertidal mussels differ from those kept under the fairly stable conditions of aquaculture (cultured mussels) in that the former have a higher level of lipids that stabilize membrane structure (cholesterol and saturated fatty acids), as well as n-6 polyunsaturated fatty acids (chiefly AA, 20:4n-6), which arguably contribute to the establishment of suitable membrane permeability and regulate the activity of membrane-bound enzymes, ion channels and receptors. High level of AA in the whole body as well as in gills of intertidal mussels appears to be due to selective retention of the fatty acid required for eicosanoid synthesis [14, 15]. In addition, unlike for mussels collected in August (where mussels were on reproductive stage IIIc or stage 0, resting), increased content of the fatty acid in the whole body of the mussels collected in June (where mussels were on reproductive stage IIIb, spawning) is needed for reproductive processes [79]. Although no differences were found in the sterol content in mussels originating from the two habitats (rocky shore and subtidal) [13], a significant excess in cholesterol level in the whole body and gills of intertidal mussels from the White Sea is probably due to the effect of severe fluctuations in temperatures (up to subzero temperatures). These features of the lipid composition are assumed to be one of the biochemical adaptation mechanisms providing for the phenotypic plasticity and survival of blue mussels in a frequently changing coastal environment. On the other hand, high level of triacylglycerols as well as elevated concentrations of n-3 polyunsaturated fatty acids, primarily of phytoplanktonic origin, EPA and DHA, in mussels collected from artificial substrates evidences high food availability (phytoplankton) and relatively stable environmental conditions in aquaculture [44, 79]. These origin-related differences of blue mussel lipid composition reflect the important role of lipids in adaptation to a changing environment.

3. Pollution effect

Natural habitats of marine aquatic organisms may also be negatively affected by human impact. Seawater is contaminated by organic and inorganic chemical substances (such as metals, petroleum hydrocarbons, pesticides) from municipal and industrial discharges. Since pollutants of various nature get either directly or indirectly involved in lipid peroxidation reactions [80–82], it is obvious that a characteristic sign of their impact on cell membranes is the disruption of lipid bilayer packing, which in its turn triggers modifications in the composition

of membrane lipid components (cholesterol, phospholipids and their fatty acids) [83]. Thus, mussels from contaminated sites had an elevated level of triacylglycerols and an increased triacylglycerol/phospholipid ratio, which implies a reduced rate of mobilization of triacylglycerols into the phospholipid pool with serious consequences for the structure and function of cell membranes. There was also a substantial decrease in phospholipids, apparently in connection with membrane destruction [84–86]. Some papers have reported the modifications in lipid and fatty acid composition of hydrobionts, including marine mussels, in response to organic and inorganic pollutants' effect [34, 37, 87–93]. Since the lipid metabolism plays an important role in living organism, it is believed that the lipid and fatty acid profile may be used to indicate the organism's health under stress conditions of pollutant effect. Exposure of blue mussels from the White Sea to various concentrations of oil products in an aquarium experiment led to an increase in the level of phospholipids and a reduction of cholesterol concentration in gills and mantle, i.e. the gateway organs for external impacts [25, 34]. These modifications in membrane lipids are believed to make cell membranes more permeable to oil products and create the conditions for their accumulation in these organs for further detoxification. A significant decrease in the level of membrane lipids—phospholipids (mainly at the expense of PC and PE) and cholesterol—simultaneously with an increase in triacylglycerols was observed in gills and digestive glands of mussels exposed to various concentrations of cadmium [37]. These modifications of the lipid profile reflect the destructive effect of cadmium on cell membranes realized through the activation of lipid peroxidation processes. It is worth noting that a significant decrease of the cholesterol concentration under the impact of oil products, mainly their high concentrations, was observed in all the studied organs (gills, mantle, mantle edge and foot) of *M. edulis* [25, 34], as well as under the impact of cobalt on *Mytilus galloprovincialis* [94]. One of the presumed examples of the toxic effect of oil products, as well as some heavy metals on bivalves, is the inhibition of cholesterol synthesis, leading to high membrane permeability. At the same time, the effect from exposure to copper as an essential metal was the opposite (significant increase of cholesterol concentration), probably meant to stabilize the membranes under the metal's oxidative action and to reduce their permeability. It was noted also that when exposed to cadmium and copper [37], as well as to relatively low concentrations of oil products [95], mussels demonstrated an elevated level of arachidonic acid. Apparently, AA involvement in the synthesis of eicosanoids ensures high resistance of the mussels to these xenobiotic impacts. On the other hand, when the concentrations of oil products were high, the level of this acid in the mussels decreased, probably due to inhibition of its biosynthesis, given the observed elevated concentrations of linoleic acid, its metabolic precursor [95]. The results of studies on the lipid composition of gills and digestive glands of intertidal blue mussels, *M. edulis*, collected from different sites in the Gulf of Kandalaksha, White Sea, prove that the composition of lipids and their fatty acids depends not only on the hydrological conditions in the habitat but also on the degree of human impact on it [38]. To wit, the fatty acid profile of the intertidal mussels living in habitats with high human impact is noted for the prevalence of oleic (18:1n-9) acid among total lipids of gills and digestive glands. A similar effect in the fatty acid composition of total lipids was observed in the mussels exposed to various doses of copper in an aquarium experiment [37]. Elevated content of non-essential oleic acid in bivalves may be associated with its additional synthesis under the toxic effect of pollutants and have the goal of binding and detoxifying xenobiotic substances. Unsaturated fatty acids are known

to be capable of forming complexes with metal ions and thus to contribute to the accumulation and detoxification of these xenobiotic substances in mussels [96].

4. Conclusions

The lipid profile of White Sea blue mussels, *M. edulis* L., is modified in response to various environmental factors in order to protect cell membranes, maintain or recover their homeostasis, replenish the cell's energy and metabolic resources and thus to secure the mussels' adaptation to the change in environmental conditions. Organ-specific distribution of lipids and fatty acids in White Sea blue mussels, as well as the dependence of the lipid and fatty acid composition response on the effect of various environmental factors on the studied organ, was detected. Modifications in the lipid composition predominantly in gills reflect the acute effect of environmental factors in aquarium experiment conditions, whereas changes in the lipid composition of digestive glands represent an adaptation of the lipid metabolism in response to chronic exposure to ambient factors (field monitoring). The composition of lipids and their fatty acids in intertidal mussels evidences their chronic exposure to abiotic environmental factors and human impact and is in agreement with data on the modifications of the lipid profile in White Sea blue mussels subjected to such environmental factors (namely, salinity, short-term anoxia, heavy metal and oil pollution) in aquarium experiments. The data discussed above prove that the lipid profile plays an important role in the adaptation of blue mussels, *M. edulis*, to new conditions in the habitat. Assessment of the lipid composition in intertidal and cultured mussels helps disclose the metabolic strategy to ensure resistance and adaptation of the organisms to environmental impacts of different nature and can be used as a biochemical marker for indicating the organism's physiological condition. This knowledge is necessary for environmental safety assessment under both natural and human impacts, as well as to predict an organism's and population's status in biomonitoring.

Acknowledgements

The results on the lipid composition of White Sea blue mussels were obtained using the facilities of the Equipment Sharing Centre of the Institute of Biology KarRC RAS (Petrozavodsk, Russia).

The authors are grateful to the administration and staff of the "Kartesh" White Sea Biological Research Station of the Zoological Institute RAS, especially Drs. V.Ja. Berger, V.V. Khalaman and A.A. Sukhotin, for the permission and help with aquarium experiments at the station, as well as to researchers from the Institute of Biology KarRC RAS I.N. Bakhmet, PhD, and I.V. Sukhovskaya, PhD, for the assistance in setting up the experiments and sampling biological material. The authors also acknowledge the help of the Kandalaksha Strict Nature Reserve staff, as well as the professor at the Petrozavodsk State University Dr. G.A. Shklyarevich in collecting material from the nature reserve territory.

The study was funded by the federal budget under state-ordered project № 0221-2014-0033.

Author details

Natalia N. Fokina*, Tatiana R. Ruokolainen and Nina N. Nemova

*Address all correspondence to: fokinann@gmail.com

Institute of Biology, Karelian Research Centre of Russian Academy of Sciences, Petrozavodsk, Russia

References

- [1] Hochachka PM, Somero GN. Biochemical Adaptation. Oxford: Princeton University Press; 2002. 480 p.
- [2] Vance DE, Vance JE, editors. Biochemistry of Lipids, Lipoproteins and Membranes. 4th ed. Amsterdam: Elsevier; 2002. 610 p.
- [3] Los DA, Murata N. Membrane fluidity and its role in the perception of environmental signals. *Biochimica et Biophysica Acta*. 2004;**1666**(1-2):142-157. DOI: <http://dx.doi.org/10.1016/j.bbamem.2004.08.002>
- [4] Lopez CS, Alice AF, Heras H, Rivas EA, Sanches-Rivas C. Role of anionic phospholipids in the adaptation of *Bacillus subtilis* to high salinity. *Microbiology*. 2006;**152**:605-616. DOI: 10.1099/mic.0.28345-0
- [5] Crockett EL. Cholesterol function in plasma membranes from ectotherms: membrane-specific roles in adaptation to temperature. *American Zoologist*. 1998;**38**(2):291-304. DOI: <http://dx.doi.org/10.1093/icb/38.2.291>
- [6] Fodor E, Jones RH, Buda C, Kitajka K, Dey I, Farkas T. Molecular architecture and biophysical properties of phospholipids during thermal adaptation in fish: an experimental and model study. *Lipids*. 1995;**30**(12):1119-1126. DOI: 10.1007/BF02536612
- [7] Logue JA, de Vries AL, Fodor E, Cossins AR. Lipid compositional correlates of temperature-adaptive interspecific differences in membrane physical structure. *Journal of Experimental Biology*. 2000;**203**:2105-2115.
- [8] Parrish CC. Essential fatty acids in aquatic food webs. In: Arts MT, Brett MT, Kainz M, editors. *Lipids in Aquatic Ecosystems*. New York: Springer; 2009. pp. 309-326. DOI: 10.1007/978-0-387-89366-2_13
- [9] Di Paolo G, De Camilli P. Phosphoinositides in cell regulation and membrane dynamics. *Nature*. 2006;**443**(7112):651-657. DOI: 10.1038/nature05185
- [10] Cossins AR. Homeoviscous adaptation of biological membranes and its functional significance. In: Cossins AR, editor. *Temperature Adaptation of Biological Membranes*. London: Portland Press; 1994. pp. 63-76.

- [11] Hazel JR, Williams EE. The role of alterations in membrane lipid-composition in enabling physiological adaptation of organisms to their physical environment. *Progress in Lipid Research*. 1990;**29**:167-227. DOI: 10.1016/0163-7827(90)90002-3
- [12] Sangiao-Alvarellos S, Laiz-Carrion R, Guzman JM, Martin del Rio MP, Miguez JM, Mancera JM, Soengas JL. Acclimation of *S. aurata* to various salinities alters energy metabolism of osmoregulatory and nonosmoregulatory organs. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2003;**285**:R897-R907. DOI: 10.1152/ajpregu.00161.2003
- [13] Freitas L, Fernandez-Reiriz MJ, Labarta U. Lipid classes of mussel seeds *Mytilus galloprovincialis* of subtidal and rocky shore origin. *Aquaculture*. 2002;**207**:97-116.
- [14] Freitas L, Fernandez-Reiriz MJ, Labarta U. Fatty acid profiles of *Mytilus galloprovincialis* (Lmk) mussel of subtidal and rocky shore origin. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*. 2002;**2**:453-461.
- [15] Freitas L, Labarta U, Fernández-Reiriz MJ. Evolution of fatty acid profiles of subtidal and rocky shore mussel seed (*Mytilus galloprovincialis*, Lmk.). Influence of environmental parameters. *Journal of Experimental Marine Biology and Ecology*. 2002;**268**(2):185-204. DOI: [http://dx.doi.org/10.1016/S0022-0981\(01\)00377-X](http://dx.doi.org/10.1016/S0022-0981(01)00377-X)
- [16] De Moreno JEA, Moreno VJ, Brenner RR. Lipid metabolism of the yellow clam, *Mesodesma mactroides*: 2-polyunsaturated fatty acid metabolism. *Lipids*. 1976;**11**:561-566. DOI: 10.1007/BF02532902
- [17] De Moreno JEA, Moreno VJ, Brenner RR. Lipid metabolism of the yellow clam, *Mesodesma mactroide*: 3-saturated fatty acids and acetate metabolism. *Lipids*. 1977;**12**:804-808. DOI: 10.1007/BF02533268
- [18] Delaunay F, Marty Y, Moal J, Samain JF. The effect of monospecific algal diets on growth and fatty acid composition of *Pecten maximus* (L) larvae. *Journal of Experimental Marine Biology and Ecology*. 1993;**173**:163-179. DOI: 10.1016/0022-0981(93)90051-O
- [19] Soudant P, Marty Y, Moal J, Masski H, Samain JF. Fatty acid composition of polar lipid classes during larval development of scallop *Pecten maximus* (L.). *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 1998;**121**(3):279-288. DOI: [http://dx.doi.org/10.1016/S1095-6433\(98\)10130-7](http://dx.doi.org/10.1016/S1095-6433(98)10130-7)
- [20] Goldberg ED. The mussel watch concept. *Environmental Monitoring and Assessment*. 1986;**7**(1):91-103. DOI: 10.1007/BF00398031
- [21] Widdows J, Donkin P. Mussels and environmental contaminants: bioaccumulation and physiological aspects. In: Gosling E, editor. *The Mussel Mytilus: Ecology, Physiology, Genetics and Aquaculture*. Amsterdam: Elsevier; 1992. pp. 383-424.
- [22] Gosling E. *The Mussel Mytilus: Ecology, Physiology, Genetics and Culture*. Amsterdam: Elsevier; 1992. 589 p.

- [23] Berger VJa, Kharazova AD. Mechanisms of salinity adaptations in marine molluscs. *Hydrobiologia*. 1997;**355**:115-126. DOI: 10.1007/978-94-017-1907-0_12
- [24] Bakhmet IN, Berger VJa, Khalaman VV. The effect of salinity change on the heart rate of *Mytilus edulis* specimens from different ecological zones. *Journal of Experimental Marine Biology and Ecology*. 2005;**318**(2):121-126. DOI: <http://dx.doi.org/10.1016/j.jembe.2004.11.023>
- [25] Bakhmet IN, Fokina NN, Nefedova ZA, Nemova NN. Physiological-biochemical properties of blue mussel *Mytilus edulis* adaptation to oil contamination. *Environmental Monitoring and Assessment*. 2009;**155**:581-591. DOI: 10.1007/s10661-008-0457-5
- [26] Bakhmet IN, Khalaman VV. Heart rate variation patterns in some representatives of Bivalvia. *Biology Bulletin*. 2006;**33**:276-280. DOI: 10.1134/S1062359006030101
- [27] Sukhotin AA, Abele D, Portner H-O. Growth, metabolism and lipid peroxidation in *Mytilus edulis*: age and size effects. *Marine Ecology Progress Series*. 2002;**226**:223-234. DOI: 10.3354/meps226223
- [28] Sukhotin AA, Lajus DL, Lesin PA. Influence of age and size on pumping activity and stress resistance in the marine bivalve *Mytilus edulis* L. *Journal of Experimental Marine Biology and Ecology*. 2003;**284**:129-144. DOI: [http://dx.doi.org/10.1016/S0022-0981\(02\)00497-5](http://dx.doi.org/10.1016/S0022-0981(02)00497-5)
- [29] Sukhotin AA, Portner H-O. Age-dependence of metabolism in mussels *Mytilus edulis* (L.) from the White Sea. *Journal of Experimental Marine Biology and Ecology*. 2001;**257**:53-72. DOI: [http://dx.doi.org/10.1016/S0022-0981\(00\)00325-7](http://dx.doi.org/10.1016/S0022-0981(00)00325-7)
- [30] Sukhotin AA, Pörtner HO. Habitat as a factor involved in the physiological response to environmental anaerobiosis of White Sea *Mytilus edulis*. *Marine Ecology Progress Series*. 1999;**184**:149-160. DOI: 10.3354/meps184149
- [31] Lesin PA, Khalaman VV, Flyachinskaya LP. Heterogeneity of the byssal threads in *Mytilus edulis* L. on different stages of attachment. *Proceedings of the Zoological Institute of the Russian Academy of Sciences*. 2006;**310**:107-112.
- [32] Lysenko LA, Kantserova NP, Käiväräinen EI, Krupnova MJu, Nemova NN. Osmotic balance in marine organisms: adaptation through protein degradation. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2012;**163**:S29-S30.
- [33] Khalaman VV, Lezin PA. Clumping behavior and byssus production as strategies for substrate competition in *Mytilus edulis*. *Invertebrate Biology*. 2015;**134**(1):38-47. DOI: 10.1111/ivb.12075
- [34] Fokina NN, Bakhmet IN, Shklyarevich GA, Nemova NN. Effect of seawater desalination and oil pollution on the lipid composition of blue mussels *Mytilus edulis* L. from the White Sea. *Ecotoxicology and Environmental Safety*. 2014;**110**:103-109. DOI: <http://dx.doi.org/10.1016/j.ecoenv.2014.08.010>
- [35] Fokina NN, Lysenko LA, Sukhovskaya IV, Vdovichenko EA, Borvinskaya EV, Kantserova NP, Krupnova MYu, Ruokolainen TR, Smirnov LP, Vysotskaya RU, Bakhmet IN, Nemova

- NN. Biochemical response of blue mussels *Mytilus edulis* L. from the White Sea to rapid changes in ambient temperature. *Journal of Evolutionary Biochemistry and Physiology*. 2015;**51**(5):378-387. DOI: 10.1134/S0022093015050038
- [36] Fokina NN, Ruokolainen TR, Bakhmet IN, Nemova NN. Lipid composition in response to temperature changes in blue mussels *Mytilus edulis* L. from the White Sea. *Journal of the Marine Biological Association of the United Kingdom*. 2015;**95**(08):1629-1634. DOI: <http://dx.doi.org/10.1017/S0025315415000326>
- [37] Fokina NN, Ruokolainen TR, Nemova NN, Bakhmet IN. Changes of blue mussels *Mytilus edulis* L. lipid composition under cadmium and copper toxic effect. *Biological Trace Element Research*. 2013;**154**(2):217-225. DOI: 10.1007/s12011-013-9727-3
- [38] Fokina NN, Shklyarevich GA, Ruokolainen TR, Nemova NN. Effect of low salinity on intertidal blue mussels, *Mytilus edulis* L., from the White Sea: lipids and their fatty acid composition as a biochemical marker. In: Snyder M, editor. *Aquatic Ecosystems: Influences, Interactions and Impact on the Environment*. New York: Nova Science Publishers Inc; 2016. pp. 87-124.
- [39] Dey I, Farkas T. Temperature shifts induce adaptive changes in the physical state of carp (*Cyprinus carpio* L.) erythrocyte plasma membranes in vitro. *Fish Physiology and Biochemistry*. 1992;**10**(4):347-355. DOI: 10.1007/BF00004484
- [40] Williams E, Somero G. Seasonal-, tidal-cycle- and microhabitat related variation in membrane order of phospholipid vesicles from gills of the intertidal mussel *Mytilus californianus*. *Journal of Experimental Biology*. 1996;**199**:1587-1596.
- [41] Parent GJ, Pernet F, Tremblay R, Sevigny JM, Ouellette M. Remodeling of membrane lipids in gills of adult hard clam *Mercenaria mercenaria* during declining temperature. *Aquatic Biology*. 2008;**3**(2):101-109. DOI: <http://doi.org/10.3354/ab00073>
- [42] Pernet F, Tremblay R, Comeau L, Guderley H. Temperature adaptation in two bivalve species from different thermal habitats: energetics and remodelling of membrane lipids. *Journal of Experimental Biology*. 2007;**210**(17):2999-3014. DOI: 10.1242/jeb.006007
- [43] Pernet F, Tremblay R, Gionet C, Landry T. Lipid remodeling in wild and selectively bred hard clams at low temperatures in relation to genetic and physiological parameters. *Journal of Experimental Biology*. 2006;**209**(23):4663-4675. DOI: 10.1242/jeb.02581
- [44] Nemova NN, Fokina NN, Nefedova ZA, Ruokolainen TR, Bakhmet IN. Modifications of gill lipid composition in littoral and cultured blue mussels *Mytilus edulis* L. under the influence of ambient salinity. *Polar Record*. 2013;**49**(03):272-277. DOI: <http://dx.doi.org/10.1017/S0032247412000629>
- [45] Kashin AG. Influence of Ambient Salinity on Lipid Composition of Some Water Invertebrates [dissertation]. Samara: Samara State University; 1997. 14 p.
- [46] Glemet HC, Ballantyne JS. Influences of environmental salinity on the structure and function of gill mitochondrial membranes of an osmoconforming invertebrate, *Crassostrea virginica*. *Marine Biology*. 1995;**121**:673-683. DOI: 10.1007/BF00349303

- [47] Fokina NN, Bakhmet IN, Nemova NN. Cooperative effect of crude oil and low salinity on the digestive glands lipid composition of the White Sea blue mussels *Mytilus edulis*. Proceedings ZIN. 2016;**320**(3):357-366.
- [48] Berger VJa. On the minimal terms of triggering the processes of phenotypic adaptation. Doklady Biological Sciences. 2005;**400**(1):57-60. DOI: 10.1007/s10630-005-0043-8
- [49] Pagliarani A, Bandiera P, Ventrella V, Trombetti F, Pirini M, Borgatti AR. Response to alkyltins of two Na⁺-dependent ATPase activities in *Tapes philippinarum* and *Mytilus galloprovincialis*. Toxicology in Vitro. 2006;**20**:1145-1153. DOI: <http://dx.doi.org/10.1016/j.tiv.2006.02.006>
- [50] Pierce SK. Invertebrate cell volume control mechanisms: a coordinated use of intracellular amino acids and inorganic ions as osmotic solute. The Biological Bulletin. 1982;**163**(3):405-419.
- [51] Fokina NN, Nefedova ZA, Ruokolainen TR, Bakhmet IN, Nemova NN. Effects of various salinity on the White Sea blue mussels *Mytilus edulis* lipid composition. Proceedings ZIN. Supplement 3. 2013;**3**:55-63.
- [52] Pollero RJ, Irazú CE, Brenner RR. Effect of sexual stages on lipids and fatty acids of *Diplodon delodontus*. Comparative Biochemistry and Physiology Part B: Comparative Biochemistry. 1983;**76**(4):927-931. DOI: 10.1016/0305-0491(83)90414-5
- [53] Zhukova NV. Non-methylene-interrupted fatty acids of marine bivalves: distribution in tissues and lipid classes. Journal of Evolutionary Biochemistry and Physiology. 1992;**28**(4):434-440.
- [54] Gardner D, Riley JP. The component fatty acids of the lipids of some species of marine and freshwater molluscs. Journal of the Marine Biological Association of the United Kingdom. 1972;**52**(04):827-838. DOI: <http://dx.doi.org/10.1017/S0025315400040571>
- [55] Fried B, Rao KS, Sherma J, Huffman JE. Fatty acid composition of *Goniobasis virginica*, *Physa* sp. and *Viviparus malleatus* (Mollusca: Gastropoda) from lake Musconetcong, New Jersey. Biochemical Systematics and Ecology. 1993;**21**(8):809-812.
- [56] Sushchik NN. Role of essential fatty acids in trophometabolic interactions in the freshwater ecosystems (a review). Zhurnal obshchei biologii. 2007;**69**(4):299-316.
- [57] Khardin AS, Aizdaicher NA, Latyshev NA. Changes in the fatty acid composition of hepatopáncreas of the mollusk *Mytilus trossulus* fed on microalgae. Russian Journal of Marine Biology. 2003;**29**(6):378-382. DOI: 10.1023/B:RUMB.0000011706.89867.ec
- [58] Alkanani T, Parrish CC, Thompson RJ, McKenzie CH. Role of fatty acids in cultured mussels, *Mytilus edulis*, grown in Notre Dame Bay, Newfoundland. Journal of Experimental Marine Biology and Ecology. 2007;**348**:33-45. DOI: <http://dx.doi.org/10.1016/j.jembe.2007.02.017>
- [59] Alyakrinskaya IO. Resistance to drying in aquatic molluscs. Biology Bulletin. 2004;**31**:299-309. DOI: 10.1023/b:bibu.0000030153.33353.77

- [60] Gromosova SA, Shapiro AZ. The Main Biochemical Patterns of Energy Metabolism in Mussels. Moscow: Legkaya I pishchevaya promyshlennost; 1984. 120 p.
- [61] Gabbott PA. Developmental and seasonal metabolic activities in marine molluscs. In: Hochachka P, editor. Mollusca Vol. 2 Environmental Biochemistry and Physiology. New York: Academic press Inc; 1983. pp. 165-217.
- [62] Fokina NN, Nefedova ZA, Nemova NN. Lipid Composition of *Mytilus edulis* L. Mussels from the White Sea. Effect of Some Environmental Factors. Petrozavodsk: Karelian Research Centre of RAS; 2010. 243 p. DOI: 10.13140/2.1.2154.8322
- [63] Fokina N, Nemova N, Nefedova Z. Fatty acid composition of mussels *Mytilus edulis* under short-term anoxia. Chemistry and Physics of Lipids. 2007;**149**:S60–S60.
- [64] Hochachka PW, Bick LT, Doll CJ, Land SC. Unifying theory of hypoxia tolerance: molecular/metabolic defense and rescue mechanisms for surviving oxygen lack. Proceedings of the National Academy of Sciences of the United States of America. 1996;**93**:9493-9498.
- [65] Zhukova NV, Imbs AB, Yi LF. Diet-induced changes in lipid and fatty acid composition of *Artemia salina*. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology. 1998;**120**(3):499-506. DOI: [http://dx.doi.org/10.1016/S0305-0491\(98\)10036-6](http://dx.doi.org/10.1016/S0305-0491(98)10036-6)
- [66] Dalsgaard J, John MS, Kattner G, Müller-Navarra D, Hagen W. Fatty acid trophic markers in the pelagic marine environment. Advances in Marine Biology. 2003;**46**:225-340. DOI: [http://dx.doi.org/10.1016/S0065-2881\(03\)46005-7](http://dx.doi.org/10.1016/S0065-2881(03)46005-7)
- [67] Kelly JR, Scheibling RE. Fatty acids as dietary tracers in benthic food webs. Marine Ecology Progress Series. 2012;**446**:1-22.
- [68] Bergé JP, Barnathan G. Fatty acids from lipids of marine organisms: molecular biodiversity, roles as biomarkers, biologically active compounds, and economical aspects. In: Le Gal Y, Ulber R, editors. Marine Biotechnology I. Berlin Heidelberg: Springer; 2005. pp. 49-125. DOI: 10.1007/b135782
- [69] Fokina NN, Ruokolainen TR, Nemova NN, Bakhmet IN. Alteration of lipid composition of blue mussels *Mytilus edulis* L. as a result of their acclimation to laboratory conditions. Proceedings of KarRC of RAS. 2015;**11**:76-84. DOI: 10.17076/eb235
- [70] Newell RIE. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North and Mid-Atlantic). Blue mussel. Biological Report. 1989;**82**(11.102) TR EL-82-4:1-34.
- [71] Gerasimova AV, Maximovich NV. Age–size structure of common bivalve mollusc populations in the White Sea: the causes of instability. Hydrobiologia. 2013;**706**(1):119-137. DOI: 10.1007/s10750-012-1415-3
- [72] Freas W, Grollman S. Ionic and osmotic influences on prostaglandin release from the gill tissue of a marine bivalve, *Modiolus demissus*. Journal of Experimental Biology. 1980;**84**:169-185.

- [73] Stanley-Samuelson DW. Physiological roles of prostaglandins and other eicosanoids in invertebrates. *The Biological Bulletin*. 1987;**173**(1):92-109.
- [74] Di Marzo V, Cimino G, Crispino A, Minardi C, Sodano G, Spinella A. A novel multifunctional metabolic pathway in a marine mollusc leads to unprecedented prostaglandin derivatives (prostaglandin 1, 15-lactones). *Biochemical Journal*. 1991;**273**:593-600. DOI: 10.1042/bj2730593
- [75] Bell JG, Sargent JR. Arachidonic acid in aquaculture feeds: current status and future opportunities. *Aquaculture*. 2003;**218**(1):491-499. DOI: [http://dx.doi.org/10.1016/S0044-8486\(02\)00370-8](http://dx.doi.org/10.1016/S0044-8486(02)00370-8)
- [76] Labarta U, Fernández-Reiriz MJ, Babarro JMF. Differences in physiological energetics between intertidal and raft cultivated mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*. 1997;**152**:167-173. DOI: 10.3354/meps152167
- [77] Babarro JMF, Fernández-Reiriz MJ, Labarta U. Metabolism of the mussel *Mytilus galloprovincialis* from two origins in the Ría de Arousa (north-west Spain). *Journal of the Marine Biological Association of the UK*. 2000;**80**(05):865-872.
- [78] Fernández-Reiriz MJ, Irisarri J, Labarta U. Flexibility of physiological traits underlying inter-individual growth differences in intertidal and subtidal mussels *Mytilus galloprovincialis*. *PLoS ONE*. 2016;**11**(2):e0148245. DOI: 10.1371/journal.pone.0148245
- [79] Fokina NN, Nefedova ZA, Nemova NN, Khalaman VV. Modulating role of lipids and their fatty acids in adaptation of the White Sea mussels *Mytilus edulis* L. to environmental salinity change. *Journal of Evolutionary Biochemistry and Physiology*. 2007;**43**(4):379-387. DOI: 10.1134/S0022093007030023
- [80] Vlahogianni TH, Valavandis A. Heavy-metal effects on lipid peroxidation and antioxidant defence enzymes in mussels *Mytilus galloprovincialis*. *Chemistry and Ecology*. 2007;**5**:361-371. DOI: <http://dx.doi.org/10.1080/02757540701653285>
- [81] Geret F, Serafim A, Barreira L, Bebianno MJ. Response of antioxidant systems to copper in the gills of the clam *Ruditapes decussatus*. *Marine Environmental Research*. 2002;**54**:413-417. DOI: [http://dx.doi.org/10.1016/S0141-1136\(02\)00164-2](http://dx.doi.org/10.1016/S0141-1136(02)00164-2)
- [82] Company R, Serafim A, Bebianno MJ, Cosson R, Shillito B, Fiala-Medioni A. Effect of cadmium, copper and mercury on antioxidant enzyme activities and lipid peroxidation in the gills of the hydrothermal vent mussel *Bathymodiulus azoricus*. *Marine Environmental Research*. 2004;**58**:377-381. DOI: <http://dx.doi.org/10.1016/j.marenvres.2004.03.083>
- [83] Thompson GA Jr. Metabolism and control of lipid structure modification. *Biochemistry and Cell Biology*. 1986;**64**(1):66-69. DOI: 10.1139/o86-010
- [84] Dzogbefia VP, Kling D, Gamble W. Polychlorinated biphenyls in vitro and in vitro modifications of phospholipid and glyceride biosynthesis. *Journal of Environmental Pathology and Toxicology*. 1978;**1**:841-856.
- [85] Capuzzo JM, Leavitt DF. Lipid composition of the digestive glands of *Mytilus edulis* and *Carcinus maenas* in response to pollutant gradients. *Marine Ecology Progress Series*. 1988;**46**:139-145.

- [86] Kato N, Kawai K, Yoshida A. Effects of dietary polychlorinated biphenyls and protein level on liver and serum lipid metabolism of rats. *Agricultural and Biological Chemistry*. 1982;**46**:703-708. DOI: <http://dx.doi.org/10.1080/00021369.1982.10865132>
- [87] Albergamo A, Rigano F, Purcaro G, Mauceri A, Fasulo S, Mondello L. Free fatty acid profiling of marine sentinels by nanoLC-ESI-MS for the assessment of environmental pollution effects. *Science of the Total Environment*. 2016;**571**:955-962. DOI: <http://dx.doi.org/10.1016/j.scitotenv.2016.07.082>
- [88] Filimonova V, Gonçalves F, Marques JC, De Troch M, Gonçalves AM. Fatty acid profiling as bioindicator of chemical stress in marine organisms: a review. *Ecological Indicators*. 2016;**67**:657-672. DOI: <http://dx.doi.org/10.1016/j.ecolind.2016.03.044>
- [89] Gladyshev MI, Anishchenko OV, Sushchnik NN, Kalacheva GS, Gribovskaya IV, Ageev AV. Influence of anthropogenic pollution on content of essential polyunsaturated fatty acids in links of food chain of river ecosystem. *Contemporary Problems of Ecology*. 2012;**5**:376-385. DOI: <http://dx.doi.org/10.1134/S1995425512040051>
- [90] Hannam ML, Bamber SD, Galloway TS, John Moody A, Jones MB. Effects of the model PAH phenanthrene on immune function and oxidative stress in the haemolymph of the temperate scallop *Pecten maximus*. *Chemosphere*. 2010;**78**:779-784. DOI: <http://dx.doi.org/10.1016/j.chemosphere.2009.12.049>ICRAM
- [91] Filimonova V, Gonçalves F, Marques JC, De Troch M, Gonçalves AM. Biochemical and toxicological effects of organic (herbicide Primextra® Gold TZ) and inorganic (copper) compounds on zooplankton and phytoplankton species. *Aquatic Toxicology*. 2016;**177**:33-43. DOI: <http://dx.doi.org/10.1016/j.aquatox.2016.05.008>
- [92] Rocchetta I, Pasquevich MY, Heras H, Ríos de Molina MdC, Luquet CM. Effects of sewage discharges on lipid and fatty acid composition of the Patagonian bivalve *Diplodon chilensis*. *Marine Pollution Bulletin*. 2014;**79**:211-219. DOI: <http://dx.doi.org/10.1016/j.marpolbul.2013.12.011>
- [93] Signa G, Di Leonardo R, Vaccaro A, Tramati CD, Mazzola A, Vizzini S. Lipid and fatty acid biomarkers as proxies for environmental contamination in caged mussels *Mytilus galloprovincialis*. *Ecological Indicators*. 2015;**57**:384-394. DOI: <http://dx.doi.org/10.1016/j.ecolind.2015.05.002>
- [94] Nechev J, Stefanov K, Popov S. Effect of cobalt ions on lipid and sterol metabolism in the marine invertebrates *Mytilus galloprovincialis* and *Actinia equina*. *Comparative Biochemistry and Physiology. Part A*. 2006;**144**:112-118. DOI: <http://dx.doi.org/10.1016/j.cbpa.2006.02.022>
- [95] Bakhmet IN, Fokina NN, Nefyodova ZA, Ruokolainen TR, Nemova NN. Blue mussels *Mytilus edulis* L. in the White Sea as bioindicators under diluted oil impact. *Proceedings of KarRC of RAS*. 2012;**2**:38-46.
- [96] Mal Reddy N, Venkateswara Rao P. Copper toxicity to the fresh water snail, *Lymnaea luteola*. *Bulletin of Environmental Contamination and Toxicology*. 1987;**39**(1):50-55. DOI: 10.1007/BF01691788

